

Person—microalbumin level (measured), total milligrams per 24 hour N[NNN].N

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Person—microalbumin level (measured), total milligrams per 24 hour N[NNN].N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Microalbumin level—milligrams per 24 hour (measured)
METEOR identifier:	270337
Registration status:	Health , Standard 01/03/2005
Definition:	A person's microalbumin level measured in milligrams per 24 hours.
Data Element Concept:	Person—microalbumin level
Value Domain:	Total milligrams per 24 hours N[NNN].N

Value domain attributes

Representational attributes

Representation class:	Total	
Data type:	Number	
Format:	N[NNN].N	
Maximum character length:	5	
	Value	Meaning
Supplementary values:	9999.9	Not stated/inadequately described
Unit of measure:	Milligram per 24-hour period (mg/24h)	

Data element attributes

Collection and usage attributes

Collection methods:	<p>Measurement of microalbumin levels should be carried out by laboratories, or practices, which have been accredited to perform these tests by the National Association of Testing Authority.</p> <p>Microalbumin is not detected by reagent strips for urinary proteins, and requires immunoassay.</p> <p>As urinary albumin varies with posture and exercise it is important to collect the urine under very standard conditions; short-term (2 hours) during rest, overnight (approximately 8 hours) or an early morning sample. For screening purposes an early morning urine specimen is adequate.</p> <p>Test for albuminuria by measuring microalbumin in timed or first morning urine sample.</p> <p>The results considered elevated are</p> <ul style="list-style-type: none">• spot urine 30 to 300mg/L; or• timed urine (24 hr collection) 20 to 200 ug/min.
----------------------------	---


Source and reference attributes


Submitting organisation:	National Diabetes Data Working Group
---------------------------------	--------------------------------------

Origin: National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary

Relational attributes

Related metadata references:

Is re-engineered from  [Microalbumin - units, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005.pdf](#) (16.3 KB)
No registration status

Is re-engineered from  [Microalbumin/protein - measured, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005.pdf](#) (16.5 KB)
No registration status

See also [Laboratory standard—upper limit of normal range for microalbumin, total milligrams per 24 hour N\[NN\].N Health, Standard 01/03/2005](#)

Implementation in Data Set Specifications: [Diabetes \(clinical\) DSS](#)
[Health, Superseded 21/09/2005](#)

DSS specific information:

A small amount of protein (albumin) in the urine (microalbuminuria) is an early sign of kidney damage. Microalbuminuria is a strong predictor of macrovascular disease and diabetic nephropathy. Incipient diabetic nephropathy can be detected by urine testing for microalbumin. Incipient diabetic nephropathy is suspected when microalbuminuria is detected in two of three samples collected over a six-month period in patients in whom other causes of an increased urinary album excretion have been excluded.

Diagnosis of microalbuminuria is established if 2 of the 3 measurements are abnormal.

According to the Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus a test for microalbuminuria is to be performed:

- at diagnosis and then every 12 months for patients with Type 2 diabetes,
- 5 years post diagnosis and then every 12 months for patients with Type 1 diabetes,
- if microalbuminuria is present, perform up to two additional measurements in the next 6 weeks.

[Diabetes \(clinical\) NBPDS](#)
[Health, Standard 21/09/2005](#)

DSS specific information:

A small amount of protein (albumin) in the urine (microalbuminuria) is an early sign of kidney damage. Microalbuminuria is a strong predictor of macrovascular disease and diabetic nephropathy. Incipient diabetic nephropathy can be detected by urine testing for microalbumin. Incipient diabetic nephropathy is suspected when microalbuminuria is detected in two of three samples collected over a six-month period in patients in whom other causes of an increased urinary album excretion have been excluded.

Diagnosis of microalbuminuria is established if 2 of the 3 measurements are abnormal.

According to the Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus a test for microalbuminuria is to be performed:

- at diagnosis and then every 12 months for patients with Type 2 diabetes,
- 5 years post diagnosis and then every 12 months for patients with Type 1 diabetes,
- if microalbuminuria is present, perform up to two additional measurements in the next 6 weeks.