

# Cancer treatment—lung cancer surgical margin qualifier, code N[N]

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Cancer treatment—lung cancer surgical margin qualifier, code N[N]

## Identifying and definitional attributes

<b>Metadata item type:</b>	Data Element
<b>Short name:</b>	Surgical margin qualifier (lung cancer)
<b>METEOR identifier:</b>	433052
<b>Registration status:</b>	<a href="#">Health</a> , Standard 08/05/2014
<b>Definition:</b>	The orientation of the surgical margin that is closest to the invasive or in situ carcinoma after surgical treatment for lung cancer, as represented by a code.
<b>Data Element Concept:</b>	<a href="#">Cancer treatment—surgical margin qualifier</a>
<b>Value Domain:</b>	<a href="#">Lung cancer surgical margin qualifier code N[N]</a>

## Value domain attributes

## Representational attributes

<b>Representation class:</b>	Code
<b>Data type:</b>	Number
<b>Format:</b>	N[N]
<b>Maximum character length:</b>	2

	Value	Meaning
<b>Permissible values:</b>	1	Bronchial
	2	Mediastinal
	3	Vascular
	4	Parenchymal
	5	Parietal pleural
	6	Chest wall
	88	Other
<b>Supplementary values:</b>	97	Not applicable-surgery was not performed
	98	Unknown whether margin involvement was present
	99	Margin involvement present but not qualified

## Collection and usage attributes

<b>Guide for use:</b>	Record the code number for the surgical margin closest to the invasive or in situ carcinoma.
-----------------------	--

## Source and reference attributes

<b>Submitting organisation:</b>	Cancer Australia
<b>Reference documents:</b>	College of American Pathologists (CAP) 2009. Protocol for the examination of specimens from patients with primary non-small cell carcinoma, small cell carcinoma, or carcinoid tumour of the lung. Viewed 7 June 2011, <a href="http://www.cap.org/apps/docs/committees/cancer/cancer_protocols/2011/Lung_11protocol.pdf">http://www.cap.org/apps/docs/committees/cancer/cancer_protocols/2011/Lung_11protocol.pdf</a>

# Data element attributes

## Collection and usage attributes

<b>Guide for use:</b>	<p>Surgical margins represent sites that have either been cut or bluntly dissected by the surgeon to resect the specimen.</p> <p>The presence of tumour at a surgical margin is an important finding because there is the potential for residual tumour remaining in the patient in the area surrounding a positive margin.</p> <p>Record the code for the margin described as the closest surgical margin from the invasive or in situ carcinoma. Where two or more margins are reported, only the closest should be recorded.</p> <p>Record only for the most definitive surgical procedure performed. For instance, if a surgical procedure to remove a portion of tumour at the primary site is followed by additional surgery to remove the remainder of the tumour at that site, code the closest surgical margin for the final surgical procedure.</p> <p>Record for the primary tumour site only, not for metastatic sites.</p>
<b>Collection methods:</b>	<p>This information should be sought from the patient's pathology report under microscopic findings.</p> <p>Collect this item when a person undergoes surgery for the treatment of lung cancer.</p>
<b>Comments:</b>	<p>Identifying the margins involved by in situ or invasive carcinoma is useful for surgical audit. Margin involvement may influence treatment decisions and is a prognostic indicator.</p>

## Source and reference attributes

**Submitting organisation:** Cancer Australia

## Relational attributes

<b>Related metadata references:</b>	<p>See also <a href="#">Cancer treatment—distance of closest surgical margin, total millimetres N[N]</a> <a href="#">Health</a>, Standard 08/05/2014</p> <p>See also <a href="#">Cancer treatment—surgical margin status, positive/negative/unknown code N</a> <a href="#">Health</a>, Standard 14/05/2015</p>
<b>Implementation in Data Set Specifications:</b>	<p><a href="#">Lung cancer (clinical) DSS</a> <a href="#">Health</a>, Superseded 14/05/2015</p> <p><b>Conditional obligation:</b> Collect when a person with cancer has undergone surgery during their initial course of cancer treatment for the purpose of removing lung cancer (either invasive or in situ).</p> <p><a href="#">Lung cancer (clinical) NBPDS</a> <a href="#">Health</a>, Standard 14/05/2015</p> <p><b>Conditional obligation:</b></p> <p>Collect when a person with cancer has undergone surgery during their initial course of cancer treatment for the purpose of removing lung cancer (either invasive or in situ).</p>