

# National Health Performance Authority, Healthy Communities, Percentage of adults who are daily smokers, 2011–12

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# National Health Performance Authority, Healthy Communities, Percentage of adults who are daily smokers, 2011–12

## Identifying and definitional attributes

<b>Metadata item type:</b>	Indicator
<b>Indicator type:</b>	Indicator
<b>Short name:</b>	Adults who smoke daily
<b>METEOR identifier:</b>	527654
<b>Registration status:</b>	<a href="#">National Health Performance Authority (retired)</a> , Retired 01/07/2016
<b>Description:</b>	Percentage of adults who are daily smokers.
<b>Indicator set:</b>	<a href="#">National Health Performance Authority: Healthy Communities: 2011–</a> <a href="#">National Health Performance Authority (retired)</a> , Retired 01/07/2016

## Collection and usage attributes

<b>Population group age from:</b>	18 years
<b>Computation description:</b>	<p>A current daily smoker was defined as a person who smokes one or more cigarettes, roll-your-own cigarettes, cigars or pipes at least once a day. Chewing tobacco and the smoking of non-tobacco products were excluded.</p> <p>This indicator was calculated using data from two components of the ABS Australian Health Survey (AHS) 2011–13, the ABS National Health Survey (NHS) 2011–12 and the ABS National Nutrition and Physical Activity Survey (NNPAS) 2011–12.</p>
<b>Computation:</b>	$100 \times (\text{Numerator} \div \text{Denominator})$
<b>Numerator:</b>	<p>Number of adults who smoke tobacco every day.</p> <ul style="list-style-type: none"><li>The numerator was calculated as the sum of calibrated sample weights for adults who responded that they currently smoke tobacco, and who were enumerated within the particular Medicare local catchment.</li></ul>

### Numerator data elements:

#### Data Element / Data Set

[Person—tobacco smoking status, code N](#)

#### Data Source

[ABS Australian Health Survey \(AHS\), 2011-13](#)

#### Guide for use

Data source type: Survey

**Denominator:**

Total population of adults

- The denominator was calculated as the sum of calibrated sample weights for adults who were enumerated within the Medicare Local catchment.
- Person level survey weights were calibrated to independent estimates of the population of interest, referred to as 'benchmarks'. Weights calibrated against population benchmarks ensure that the survey estimates conform to independently estimated distributions of the population, rather than to the distribution within the sample itself. These benchmarks account for the distribution of people across state and territory, age group, and sex categories. Note: These benchmarks have not been calibrated for Medicare Local geography.

**Denominator data elements:****Data Element / Data Set**[Person—age, total years N\[NN\]](#)**Data Source**[ABS Australian Health Survey \(AHS\), 2011-13](#)**Guide for use**

Data source type: Survey

**Disaggregation:**

Medicare Local catchments and Medicare Local peer groups.

- Medicare Locals are primary health care organisations established to improve responsiveness, coordinate primary health care delivery and tackle local health care needs and service gaps. A Medicare Local catchment refers to the geographical area of a Medicare local.
- Medicare Local peer groups are the grouping of the 61 Medicare Locals into seven peer groups. This grouping was undertaken by the National Health Performance Authority using statistical cluster analysis of 2006 Census based socioeconomic status, for each Medicare Local and 2006 Census based Remoteness Area categories (ABS cat. no. 2039.0). Additional information on the average distance to the closest large city and major hospital (A1 public hospital peer group classification, 2010–11) contributed to the assignment of Medicare Locals to peer groups. Groupings were tested for face validity with a stakeholder group. The National Health Performance Authority has created 7 peer groups: Metro 1 to 3, Regional 1 and 2 and Rural 1 and 2.

**Disaggregation data elements:****Data Element / Data Set**[Address—Australian postcode, Australian postcode code \(Postcode datafile\) {NNNN}](#)**Data Source**[ABS Australian Health Survey \(AHS\), 2011-13](#)

**Comments:**

National Health Performance Authority suppression protocols:

- Additional suppression rules were developed and applied by the National Health Performance Authority to ensure robust reporting of these data at small areas.
- These suppression rules are based on limits for Relative Standard Error<sup>1</sup> and Confidence Interval width of 30%, with additional cross-validation for estimates close to these limits, that is plus or minus 3% of the limits. If an estimate was marginal<sup>2</sup> with respect to Relative Standard Error, the Confidence Interval width was used as the deciding factor. If an estimate was marginal<sup>2</sup> with respect to Confidence Interval width, then Relative Standard Error is used as the deciding factor.
- Data were suppressed based on the following rules:
  - Relative Standard Error of 33% or greater, or
  - Confidence Interval (95%) width of 33% or greater, or
  - Relative Standard Error between 27% and 33%, with significantly<sup>3</sup> wider Confidence Interval width than the average for that indicator, or
  - Confidence Interval width between 27% and 33%, with significantly<sup>3</sup> wider Relative Standard Error than the average for that indicator.

1. For a dichotomous proportion, Relative Standard Error can be defined as the ratio of the standard error and the minimum of the estimate and its complement (100%-estimate).
2. In this context, marginal is defined as within 10% of the 30% limit, or +/- 3%.
3. In this context, statistical significance is defined as at least two standard deviations above average.

## Representational attributes

**Representation class:** Percentage

**Data type:** Real

**Unit of measure:** Person

**Format:** N[NN].N

## Indicator conceptual framework

**Framework and dimensions:** [PAF-Responsiveness](#)

## Data source attributes

**Data sources:**

**Data Source**

[ABS Australian Health Survey \(AHS\), 2011-13](#)

**Frequency**

Every 3 years

**Data custodian**

Australian Bureau of Statistics

## Accountability attributes

**Reporting requirements:** National Healthcare Agreement

**Organisation responsible for providing data:** Australian Bureau of Statistics

**Further data development / collection required:** Specification: Long-term

## Source and reference attributes

**Submitting organisation:** NHPA  
**Origin:** Healthy Communities  
**Reference documents:** NHR\_PAF 2011