

6.7 Number of heart failure admissions to hospital per 100,000 people aged 40 years and over, 2012–13

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

6.7 Number of heart failure admissions to hospital per 100,000 people aged 40 years and over, 2012–13

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Heart failure hospital admissions 40 years and over, 2012-13
METEOR identifier:	611789
Registration status:	Australian Commission on Safety and Quality in Health Care , Standard 23/11/2016 National Health Performance Authority (retired) , Retired 01/07/2016
Description:	Number of admissions to hospital for heart failure per 100,000 people aged 40 years and over, age standardised. Data are disaggregated by the area in which the person lives.
Indicator set:	Australian Atlas of Healthcare Variation Australian Commission on Safety and Quality in Health Care , Standard 23/11/2016 National Health Performance Authority (retired) , Retired 01/07/2016

Collection and usage attributes

Population group age from: 40 years and over

Computation description:

Presented as a number per 100,000 people, age standardised, by Statistical Area Level 3.

Analysis by Statistical Area Level 3 (SA3) is based on postcode of usual residence of the patient. The postcode is based on that which is provided by the patient to the hospital at which they are admitted.

SA3s where the total population was less than 2,500 or where the population of people aged 40 years and over was less than 200 are excluded.

Suppression protocol for calculating age standardised rates:

Numerator	Total admissions less than 20 (unrounded)
Denominator	ERP for one or more five year age groups less than 30

Computation: $100,000 \times (\text{Numerator} \div \text{Denominator})$, age standardised

For more information about age-standardisation, see

</content/index.phtml/itemId/327276>

Numerator: Number of heart failure admissions to hospital for people aged 40 years and over in 2012-13.

Numerator data elements:

Data Element / Data Set

[Person—date of birth, DDMMYYYY](#)

NMDS / DSS

[Admitted patient care NMDS 2012-13](#)

Guide for use

People aged 40 years and over in 2012-13 who were admitted to hospital with a diagnosis as described above were included in the analysis of this item.

Data Element / Data Set

[Episode of care—principal diagnosis, code \(ICD-10-AM 7th edn\) ANN{.N\[N\]}](#)

Data Source

[National Hospital Morbidity Database \(NHMD\)](#)

NMDS / DSS

[Admitted patient care NMDS 2012-13](#)

Guide for use

Inclusion codes and description for principal diagnosis only:

ICD-10-AM (7th edn) code	Description
I150	Congestive heart failure
I110	Hypertensive heart disease with (congestive) heart failure
J81	Pulmonary oedema

Patients with a principal diagnosis of I150, I110, J81 AND the following procedure blocks/codes are excluded

Procedure blocks	Description
[600-606]	Heart - Atrium procedures
[608-650]	Heart - Ventricle procedures
[653-657]	Insertion, adjustment or removal of cardiac pacemaker or defibrillator
[660-664], [666]	Transplantation of heart or lung, other repair, reconstruction or revision procedures on other sites of the heart
[669-682]	Excision, repair, bypass procedures on coronary arteries
[684-691], [693]	Repair procedures on aorta (aortic arch and thoracic) and closure of patent ductus arteriosus
[705-707]	Resection of lesion of carotid artery and patch graft of artery
[717]	Repair procedures on pulmonary artery for congenital heart disease
Procedure codes	
33172-00[715]	Replace other major arterial aneurysm with graft
33827-01[733]	Repair jugular vein by direct anastomosis
34800-00[726]	Interruption of vena cava
35412-00[11]	Endovas occl cerebral aneur / AV malform

38721-01[733]	Rep vena cava by direct anastomosis
90217-02[734]	Repair vena cava by interpstn gft
90215-02[732]	Direct closure of vena cava

Data source type: Administrative by-product data

Data Element / Data Set

[Hospital service—care type, code N\[N\]](#)

NMDS / DSS

[Admitted patient care NMDS 2012-13](#)

Guide for use

Include all care types except:

Code	Description
7.3	Unqualified newborn
10.0	Hospital border
9.0	Organ procurement

Denominator:

Total population aged 40 years and over.

Denominator data elements:

Data Element / Data Set

Data Element / Data Set

[Person—estimated resident population of Australia, total people N\[N\(7\)\]](#)

Data Source

[ABS Estimated resident population \(total population\)](#)

Guide for use

ABS Estimated resident population as at June 2001

Data source type: Census based plus administrative by-product data

Disaggregation:

Statistical Area Level 3, which have been derived from [Postcode—Australian \(person\)](#):

- Statistical Area Level 3 (SA3s) are geographic areas defined in the ABS Australian Statistical Geography Standard (ASGS). The aim of SA3s is to create a standard framework for the analysis of ABS data at the regional level through clustering groups of SA2s that have similar regional characteristics. There are 333 spatial SA3s covering the whole of Australia without gaps or overlaps. They are designed to provide a regional breakdown of Australia. SA3s generally have a population of between 30,000 and 130,000 people. There are approximately 50 with fewer than 30,000 people and 35 with more than 130,000 as at 30 June 2011. For further information see the ABS publication, Australian Statistical Geography Standard (ASGS): Volume 1 – Main Structure and Greater Capital City Statistical Areas, July 2011 (cat. no. 1270.0.55.001).

Disaggregation data elements:

Data Element / Data Set

[Address—Australian postcode, Australian postcode code \(Postcode datafile\) {NNNN}](#)

Data Source

[National Hospital Morbidity Database \(NHMD\)](#)

Guide for use

Postcode is based on that which is provided by the patient to the hospital at which they are admitted.

Data source type: Administrative by-product data

Used for disaggregation by Statistical Area Level 3.

Data Element / Data Set

[Address—statistical area, level 3 \(SA3\) code \(ASGS 2011\) NNNNN](#)

Representational attributes

Representation class: Rate
Data type: Integer
Unit of measure: Episode
Format: NN[NNNN]

Data source attributes

Data sources:

Data Source

[National Hospital Morbidity Database \(NHMD\)](#)

Frequency

Annual

Data custodian

Australian Institute of Health and Welfare

Data Source

[ABS Estimated resident population \(total population\)](#)

Frequency

Quarterly

Data quality statement

[ABS Estimated resident population \(total population\), QS](#)

Data custodian

Australian Bureau of Statistics

Accountability attributes

Reporting requirements: Australian Commission of Safety and Quality in Health Care's Atlas of Healthcare Variation, released November 2015

**Organisation responsible
for providing data:** Australian Institute of Health and Welfare

Accountability: Australian Commission of Safety and Quality in Health Care

Source and reference attributes

Submitting organisation: National Health Performance Authority