

Hip fracture care clinical care standard indicators: 5b- Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Hip fracture care clinical care standard indicators:

5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery
METEOR identifier:	628094
Registration status:	Health , Standard 12/09/2016
Description:	Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery.
Rationale:	Restricted weight bearing can reduce and delay patients' functional recovery and return to independent living (ANZHFR Steering Group 2014; Ariza-Vega et al. 2014).
Indicator set:	Clinical care standard indicators: hip fracture Australian Commission on Safety and Quality in Health Care , Superseded 18/06/2018 Health , Standard 12/09/2016
Outcome area:	Mobilisation and weight-bearing Health , Standard 12/09/2016

Collection and usage attributes

Computation description:	<p>For the numerator, unrestricted weight bearing status refers to the fact that the patient can be mobilised with full use of the affected limb, to weight bear as pain allows, immediately following surgery for hip fracture. This is contrasted with patients with restricted/non weight bearing status immediately post surgery, which means that there is a specific instruction that prevents the patient being allowed to fully utilise the leg irrespective of the degree of pain. Restricted weight bearing includes terms such as partial weight bear, touch weight bear and non weight bear.</p> <p>In the phrase "unrestricted weight bearing immediately post surgery", <i>immediately</i> refers to what should be aimed for with a patient as soon as practical after the hip fracture surgery. It is usually no later than the next calendar day following the surgery, when it is expected that the patient is first mobilised.</p> <p>Both the numerator and the denominator exclude patients where Episode of admitted patient care—separation mode, code N = 8 Died, and the date of death was the date of the hip fracture surgery or the day following the surgery.</p> <p>Presented as a percentage.</p>
Computation:	$(\text{Numerator} \div \text{denominator}) \times 100$
Numerator:	Number of patients admitted to hospital with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery.
Denominator:	Number of patients admitted to hospital with a hip fracture who underwent surgery for their hip fracture.
Comments:	For hospitals collecting the Australian and New Zealand Hip Fracture Registry (ANZHFR) data set (ANZHFR Steering Group 2013), the variable <i>Full weight bear</i> can be used for the numerator of this indicator.

Representational attributes

Representation class: Percentage
Data type: Real
Unit of measure: Service event
Format: N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: ANZHFR (Australian and New Zealand Hip Fracture Registry) Steering Group 2013. Data dictionary. Sydney: ANZHFR. Viewed 5 May 2016, <http://www.anzhfr.org/images/resources/Data%20Dictionary%20v8%20Dec%202013.pdf>.

ANZHFR Steering Group 2014. Australian and New Zealand guideline for hip fracture care: improving outcomes in hip fracture management of adults. Sydney: ANZHFR.

Ariza-Vega P, Jimenez-Moleon JJ & Kristensen MT 2014. Non-weight-bearing status compromises the functional level up to 1 yr after hip fracture surgery. American Journal of Physical Medicine & Rehabilitation / Association of Academic Physiatrists 93(8):641-8.