National Healthcare Agreement: PI 05–Levels of risky alcohol consumption, 2017

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY4.0 (CC BY4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

National Healthcare Agreement: PI 05–Levels of risky alcohol consumption, 2017

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Progress measure
Short name:	PI05-Levels of risky alcohol consumption, 2017
METEOR identifier:	629999
Registration status:	Health, Superseded 30/01/2018
Description:	Proportion of adults at risk of long-term harm from alcohol.
Indicator set:	National Healthcare Agreement (2017) Health, Superseded 30/01/2018
Outcome area:	Prevention <u>Health</u> , Standard 07/07/2010
Data quality statement:	National Healthcare Agreement: PI 05-Levels of risky alcohol consumption, 2017 QS Health, Standard 31/01/2017

Collection and usage attributes

Population group age from:	18 years
Computation description:	Data on persons 'at risk of long term alcohol related harm' is based on the 2009 National Health and Medical Research Council (NHMRC) guideline 'for healthy men and women, drinking no more than two standard drinks on any day reduces the lifetime risk of harm from alcohol-related disease or injury' and has been operationalised as: for both males and females, an average of more than 2 standard drinks per day in the last week.
	Rates are directly age-standardised to the 2001 Australian population.
	Analysis by remoteness and Socio-Economic Indexes for Areas (SEIFA) Index of Relative Socio-economic Disadvantage (IRSD) is based on usual residence of person.
	Presented as a percentage.
	95% confidence intervals and relative standard errors are calculated for rates.
Computation:	100 × (Numerator ÷ Denominator)
Numerator:	Number of persons aged 18 years or over assessed as having an alcohol consumption pattern that puts them at risk of long term alcohol-related harm.
Numerator data elements:	Data Element / Data Set
	Data Element
	Person-age
	Data Source
	ABS 2014-15 National Health Survey (NHS)
	Guide for use
	Data source type: Survey
	Data Element / Data Set

Data Element

Person-alcohol consumption amount

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

-Data Element / Data Set

Data Element

Person—alcohol consumption frequency

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

-Data Element / Data Set-

Data Element

Person-age

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Guide for use

Data source type: Survey

-Data Element / Data Set-

Data Element

Person-alcohol consumption amount

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Guide for use

Data source type: Survey

Denominator:

Population aged 18 years or over.

Denominator data elements:	Data Element / Data Set
	Data Element
	Person—age
	Data Source
	ABS 2014-15 National Health Survey (NHS)
	Guide for use
	Data source type: Survey
	Data Element / Data Set
	Data Element
	Person—age
	Data Source
	ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)
	Guide for use
	Data source type: Survey
Disaggregation:	2014–15—State and territory, by:
	 sex by age (not reported) Indigenous status (non-Indigenous not reported) remoteness (Australian Statistical Geography Standard (ASGS) Remoteness Structure) 2011 SEIFA IRSD quintiles disability status (not reported) 2014–15—Nationally, by (all not reported): sex by remoteness (ASGS Remoteness Structure).
	 2011 SEIFA IRSD deciles. remoteness (ASGS Remoteness Structure) by 2011 SEIFA IRSD deciles.
	Some disaggregation may result in numbers too small for publication.
Disaggregation data	Data Element / Data Set
elements:	Data Element
	Person—age
	Data Source
	ABS 2014-15 National Health Survey (NHS)
	Guide for use
	Data source type: Survey
	Data Element / Data Set
	Data Element
	Person—area of usual residence
	Data Source
	ABS 2014-15 National Health Survey (NHS)
	Guide for use

Data source type: Survey Used for disaggregation by state/territory, remoteness and SEIFA IRSD

-Data Element / Data Set-

Data Element

Person-disability status

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

- Data Element / Data Set-

Data Element

Person-Indigenous status

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

Data Element / Data Set-

Data Element

Person-area of usual residence

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Guide for use

Data source type: Survey Used for disaggregation by state/territory

Comments:

Most recent data available for 2017 National Healthcare Agreement performance reporting: 2014–15 (total population, non-Indigenous: NHS); 2014–15 (Indigenous only: NATSISS).

National Aboriginal and Torres Strait Islander Social Survey (NATSISS) or National Aboriginal and Torres Strait Islander Health Survey (NATSIHS) data may be used for analysis, dependent upon which survey is most recent.

Data on persons 'at risk of long term alcohol related harm' had previously been defined according to the 2001 NHMRC guidelines: for males, an average of more than 4 standard drinks per day in the last week, and for females, an average of more than 2 standard drinks per day in the last week.

Representational attributes

Representation class:	Percentage
Data type:	Real
Unit of measure:	Person
Format:	N[N].N

Indicator conceptual framework

Framework and <u>Health behaviours</u> dimensions:

Data source attributes

Data sources:

Data Source
ABS 2014-15 National Health Survey (NHS)
Frequency
Every 3 years
Data custodian
Australian Bureau of Statistics
- Data Source
ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)
Frequency
Every 6 years
Data custodian
Australian Bureau of Statistics

Accountability attributes

Organisation responsible Australian Bureau of Statistics for providing data:

Further data development / Specification: Final, the measure meets the intention of the indicator. **collection required:**

Relational attributes

Supersedes <u>National Healthcare Agreement: PI05–Levels of risky alcohol</u> consumption, 2016

Health, Superseded 31/01/2017

Has been superseded by <u>National Healthcare Agreement: P105–Levels of risky</u> alcohol consumption, 2018

Health, Superseded 19/06/2019

See also National Healthcare Agreement: PB e–Better health: by 2018, reduce the national smoking rate to 10 per cent of the population and halve the Indigenous smoking rate over the 2009 baseline, 2017 Health, Superseded 30/01/2018

See also <u>National Healthcare Agreement: PI03–Prevalence of overweight and</u> obesity, 2017

Health, Superseded 30/01/2018

See also <u>National Healthcare Agreement: PI04–Rates of current daily smokers,</u> 2017

Health, Superseded 30/01/2018

See also <u>National Healthcare Agreement: PI 16–Potentially avoidable deaths</u>, <u>2017</u>

Health, Superseded 30/01/2018

See also <u>National Indigenous Reform Agreement: PI 04-Levels of risky alcohol</u> consumption, 2018

Indigenous, Superseded 31/07/2018