

Osteoarthritis of the knee clinical care standard indicators: 1a-Local arrangements to ensure that patients newly diagnosed with knee osteoarthritis have a comprehensive assessment

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Osteoarthritis of the knee clinical care standard indicators: 1a-Local arrangements to ensure that patients newly diagnosed with knee osteoarthritis have a comprehensive assessment

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 1a-Local arrangements to ensure that patients newly diagnosed with knee osteoarthritis have a comprehensive assessment
METEOR identifier:	644252
Registration status:	Health , Standard 02/08/2017
Description:	Documented evidence of local arrangements to ensure that patients who are newly diagnosed with knee osteoarthritis have a comprehensive assessment undertaken.
Rationale:	Assessment of a comprehensive range of factors is essential to clinical decision-making (Pencharz & MacLean 2004).
Indicator set:	Clinical care standard indicators: osteoarthritis of the knee Health , Standard 02/08/2017

Collection and usage attributes

Computation description:	<p>The local arrangements should specify that all newly diagnosed patients with knee osteoarthritis have a comprehensive assessment undertaken.</p> <p>A 'comprehensive assessment' involves a clinician taking a detailed history of the patient that includes the presenting symptoms and other health conditions, a physical examination (including measurement of height and weight to calculate body mass index), and a psychosocial evaluation, to identify factors (especially level of pain) that may impact on the patient's ability to do their usual daily activities and participate in work, leisure and social activities. A comprehensive assessment should use a clinically validated assessment tool. A range of clinically validated assessment tools is available (see list in the <i>Comments</i> section).</p> <p>'Newly diagnosed' means that the patient was diagnosed or referred with knee osteoarthritis in the last three months.</p>
Computation:	<p>Healthcare settings where care is provided to patients with knee osteoarthritis that have documented evidence of local arrangements, should record 'Yes'.</p> <p>Otherwise, the healthcare setting should record 'No'.</p>

Comments:	<p>A comprehensive assessment should use clinically validated assessment tools, such as:</p> <ul style="list-style-type: none"> • Disease-specific - Knee injury and Osteoarthritis Outcome Score (KOOS) (Roos et al. 1998), see survey instrument http://www.koos.nu; OsteoArthritis Questionnaire (OA-Quest) (Busija, Buchbinder & Osborne 2016), see survey instrument http://iha.acu.edu.au/osteo-arthritis/ • Pain and function - Timed Up and Go Test and 30-second Chair Stand Test (OARSI 2013), see survey instrument https://www.oarsi.org/research/physical-performance-measures • Pain - Verbal Numerical Rating Scale (VNRS) and Visual Analogue Scale (VAS) (NSW Agency for Clinical Innovation Pain Management Network 2017a, 2017b), see survey instrument http://www.aci.health.nsw.gov.au/chronic-pain/health-professionals/assessment • Depression - Kessler (K10) Psychological Distress Scale (Kessler et al. 2016), see survey instrument http://www.gpcare.org/outcome%20measures/outcomemeasures.html and Depression, Anxiety and Stress Scale 21 (DASS 21) (Henry & Crawford 2005) see survey instrument http://www.depression-test.net/support-files/dass21-dmz.pdf • Activities and quality of life - Workplace Activity Limitations Scale (WALS) (Gignac 2005; Gignac, Sutton & Badley 2007), see survey instrument http://www.acreu.ca/pdf/WALS.pdf; Assessment of Quality of Life (Richardson, Iezzi, Khan et al. 2014), see survey instrument http://www.aqol.com.au/. <p>This indicator has been adapted from the <i>Osteoarthritis quality standard</i> (NICE 2015).</p>
------------------	---

Representational attributes

Representation class:	Count
Data type:	Real
Unit of measure:	Service event
Format:	Yes/No

Accountability attributes

Other issues caveats:	Applicable setting: all healthcare settings where care is provided to patients with knee osteoarthritis, including primary care, specialist care, hospitals and community settings.
------------------------------	---

Source and reference attributes

Submitting organisation:	Australian Commission on Safety and Quality in Health Care
---------------------------------	--

Reference documents:

- Busija L, Buchbinder R & Osborne RH 2016. Development and preliminary evaluation of the OsteoArthritis Questionnaire (OA-Quest): a psychometric study. *Osteoarthritis and Cartilage* 24(8):1357-66.
- Gignac MA 2005. Arthritis and employment: an examination of behavioral coping efforts to manage workplace activity limitations. *Arthritis & Rheumatism* 53(3):328-36.
- Gignac MA, Sutton D, & Badley EM 2007. Arthritis symptoms, the work environment, and the future: Measuring perceived job strain among employed persons with arthritis. *Arthritis & Rheumatism* 57(5):738-47.
- Henry JD & Crawford JR 2005. The 21-item version of the Depression Anxiety Stress Scales (DASS-21): Normative data and psychometric evaluation in a large non-clinical sample. *British Journal of Clinical Psychology* 44:227-239.
- Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand S-LT et al. 2016. Anxiety and depression checklist (K10). Melbourne: beyondblue. Viewed 9 April 2017, <https://www.beyondblue.org.au/the-facts/anxiety-and-depression-checklist-k10>.
- NICE (National Institute for Health and Care Excellence) 2015. Osteoarthritis Quality standard 87. London: NICE.
- NSW Agency for Clinical Innovation Pain Management Network 2017a. Verbal numerical rating scale (VNRS). Sydney: NSW ACI. Viewed 9 April 2017, http://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0017/212912/Verbal_Numerical_Rating_Scale.pdf.
- NSW Agency for Clinical Innovation Pain Management Network 2017b. Visual analogue scale (VAS). Sydney: NSW ACI. Viewed 9 April 2017, http://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0018/212913/Visual_Analogue_Scale.pdf.
- OARSI (Osteoarthritis Research Society International) 2013. Physical performance measures. Viewed 9 April 2017, <https://www.oarsi.org/research/physical-performance-measures>.
- Pencharz JN & MacLean CH 2004. Measuring quality in arthritis care: the Arthritis Foundation's Quality Indicator set for osteoarthritis. *Arthritis & Rheumatism* 51(4): 538-548.
- Richardson J, Iezzi A, Khan MA & Maxwell A 2014. Validity and reliability of the Assessment of Quality of Life (AQoL-8D) multi attribute utility instrument. *Patient* 7(1):85-96.
- Roos EM, Roos HP, Lohmander LS, Ekdahl C & Beynnon BD 1998. Knee Injury and Osteoarthritis Outcome Score (KOOS) - development of a self-administered outcome measure. *Journal of Orthopaedic & Sports Physical Therapy* 28:88-96.