

Osteoarthritis of the knee clinical care standard indicators: 7a-Number of patients undergoing arthroscopic procedures for knee osteoarthritis

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Osteoarthritis of the knee clinical care standard indicators: 7a-Number of patients undergoing arthroscopic procedures for knee osteoarthritis

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 7a-Number of patients undergoing arthroscopic procedures for knee osteoarthritis
METEOR identifier:	644337
Registration status:	Health , Standard 02/08/2017
Description:	Number of patients undergoing arthroscopic procedures for knee osteoarthritis .
Rationale:	Several large, high-quality randomized controlled trials have failed to show any clinical benefit of arthroscopic procedures for knee osteoarthritis (Herrlin et al. 2013; Kirkley et al. 2008; Kise et al. 2016; Moseley et al. 2002). There may also be some harms related to this procedure (Thorlund et al. 2015).
Indicator set:	Clinical care standard indicators: osteoarthritis of the knee Health , Standard 02/08/2017

Collection and usage attributes

Population group age from:	45 years
Computation description:	<p>Count of patients with a diagnosis of knee osteoarthritis undergoing arthroscopic procedures. Exclude from the count patients with:</p> <ul style="list-style-type: none">• true mechanical locking• septic arthritis. <p>Data from hospital information systems can be used to identify patients with knee osteoarthritis undergoing arthroscopic procedures. A chart audit is recommended for any cases identified, to assess whether these are due to errors in the data or documentation errors in the first instance. Any cases remaining following the audit should be assessed clinically to determine whether they are appropriate.</p> <p>Note: Different codes may apply if the data is outside of the period 1 July 2015 to 30 June 2017.</p> <p>Patients with knee osteoarthritis are those with a principal or additional diagnosis code (ICD-10-AM 9th edn.) of one of the following:</p> <ul style="list-style-type: none">• M17.0 Primary gonarthrosis, bilateral [arthrosis of the knee]• M17.1 Other primary gonarthrosis (includes: not otherwise specified, unilateral) [arthrosis of the knee]• M23.2- Derangement of meniscus due to old tear or injury• S83.2 Tear of meniscus, current. <p>However, exclude from the count patients with any one of the following codes as a principal or additional diagnosis coded in addition to one of the codes for osteoarthritis listed above:</p> <ul style="list-style-type: none">• M23.8 Other internal derangements of knee (includes: laxity of ligament of knee, snapping knee)• M00.96 Pyogenic arthritis, unspecified, lower leg (includes: infective arthritis not otherwise <p>Note: '-' refers to the fifth character subdivision indicating the site of involvement (e.g. anterior cruciate ligament or anterior horn of medial meniscus or posterior</p>

cruciate ligament or posterior horn of medial meniscus).

Arthroscopic procedures include the following procedure codes (ACHI 9th edn.)

ACHI 9 th edition procedure code	Description
49557-00 [1501]	Arthroscopy of knee
49560-00 [1501]	Arthroscopic removal of loose body of knee
49560-02 [1501]	Arthroscopic lateral release of knee
49557-01 [1502]	Arthroscopic biopsy of knee
49558-00 [1503]	Arthroscopic debridement of knee
49560-01 [1503]	Arthroscopic trimming of ligament of knee
49557-02 [1503]	Arthroscopic excision of meniscal margin or plica of knee
49560-03 [1503]	Arthroscopic meniscectomy of knee
49561-02 [1511]	Arthroscopic removal of loose body of knee with debridement, osteoplasty or chondroplasty
49562-02 [1511]	Arthroscopic removal of loose body of knee with chondroplasty and multiple drilling or implant (includes: carbon fibre implant, debridement, osteoplasty)
49561-00 [1514]	Arthroscopic lateral release of knee with debridement, osteoplasty or chondroplasty
49562-00 [1514]	Arthroscopic lateral release of knee with chondroplasty and multiple drilling or implant (includes: carbon fibre implant, debridement, osteoplasty)
49561-01 [1517]	Arthroscopic meniscectomy of knee with debridement, osteoplasty or chondroplasty
49562-01 [1517]	Arthroscopic meniscectomy of knee with chondroplasty and multiple drilling or implant (includes: carbon fibre implant, debridement, osteoplasty)
49558-01 [1520]	Arthroscopic chondroplasty of knee (includes: debridement, implant (carbon fibre), multiple drilling, osteoplasty)
49558-02 [1520]	Arthroscopic osteoplasty of knee

Presented as a number.

Computation: Number of patients with a diagnosis of knee osteoarthritis undergoing arthroscopic procedures.

Comments: This indicator has been adapted from the *Osteoarthritis quality standard* (NICE 2015).

Representational attributes

Representation class: Count
Data type: Real
Unit of measure: Service event
Format: N[NNN]

Accountability attributes

Other issues caveats: Applicable setting: hospital or any setting in which arthroscopic procedures are undertaken (e.g. day procedure centre).

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: Herrlin SV, Wange PO, Lapidus G, Hallander M, Werner S & Weidenhielm L 2013. Is arthroscopic surgery beneficial in treating non-traumatic, degenerative medial meniscal tears? A five year follow-up. *Knee Surgery, Sports Traumatology, Arthroscopy* 21(2): 358-364.

Kirkley A et al. 2008. A randomized trial of arthroscopic surgery for osteoarthritis of the knee. *New England Journal of Medicine* 359(11): 1097-1107.

Kise NJ, Risberg MA, Stensrud S, Ranstam J, Engebretsen L & Roos EM 2016. Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomised controlled trial with two year follow-up. *BMJ* 354: i3740.

Moseley JB, O'Malley K, Petersen NJ, Menke TJ, Brody BA, Kuykendall DH et al. 2002. A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *New England Journal of Medicine* 347(2): 81-88.

NICE (National Institute for Health and Care Excellence) 2015. Osteoarthritis Quality standard 87. London: NICE.

Thorlund JB, Juhl CB, Roos EM & Lohmander LS 2015. Arthroscopic surgery for degenerative knee: systematic review and meta-analysis of benefits and harms. *BMJ* 350:h2747.