Acute stroke clinical care standard indicators: 2c-Proportion of patients with a final diagnosis of ischaemic stroke provided thrombolysis who received the therapy within 60 minutes of presentation to hospital, 2019-

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute stroke clinical care standard indicators: 2c-Proportion of patients with a final diagnosis of ischaemic stroke provided thrombolysis who received the therapy within 60 minutes of presentation to hospital, 2019-

Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

Short name: Indicator 2c-Proportion of patients with a final diagnosis of ischaemic stroke

provided thrombolysis who received the therapy within 60 minutes of presentation

to hospital

METEOR identifier: 719050

Registration status: Australian Commission on Safety and Quality in Health Care, Qualified 09/09/2019

Description: Proportion of <u>acute stroke</u> patients provided thrombolysis who received

thrombolysis within 60 minutes of presentation to hospital.

Indicator set: Clinical care standard indicators: acute stroke

Australian Commission on Safety and Quality in Health Care, Standard

03/11/2020

Collection and usage attributes

Computation description: Both the numerator and the denominator include patients with a final diagnosis

of ischaemic stroke who were provided thrombolysis.

Presented as a percentage.

Computation: (Numerator ÷ denominator) x 100

Numerator: Number of patients with a final diagnosis of ischaemic stroke who were provided

thrombolysis who received the therapy within 60 minutes of presentation to hospital.

Denominator: Number of patients with a final diagnosis of ischaemic stroke who were provided

thrombolysis.

Representational attributes

Representation class: Percentage

Data type: Real

Unit of measure: Service event

Format: N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care