

Acute stroke clinical care standard indicators: 6a- Proportion of patients with a final diagnosis of acute stroke whose carer(s) received a formal needs assessment prior to separation from hospital, 2019-

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute stroke clinical care standard indicators: 6a- Proportion of patients with a final diagnosis of acute stroke whose carer(s) received a formal needs assessment prior to separation from hospital, 2019-

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 6a- Proportion of patients with a final diagnosis of acute stroke whose carer(s) received a formal needs assessment prior to separation from hospital
METEOR identifier:	719068
Registration status:	Australian Commission on Safety and Quality in Health Care , Qualified 09/09/2019
Description:	Proportion of patients with a final diagnosis of acute stroke whose carer(s) have had a documented formal needs assessment prior to separation from hospital.
Rationale:	Assessment of carer support needs allows for referral to appropriate services.
Indicator set:	Clinical care standard indicators: acute stroke Australian Commission on Safety and Quality in Health Care , Standard 03/11/2020

Collection and usage attributes

Computation description:	<p>Both the numerator and the denominator include patients with a final diagnosis of acute stroke. The final diagnosis is made at the hospital where the patient is admitted for the acute phase of management of their stroke.</p> <p>Both the numerator and the denominator only include patients separated to their usual residence, own accommodation or welfare institution following the acute episode of care (i.e. where Episode of admitted patient care—separation mode, code N = 9 Other). Welfare institutions includes prisons, hostels and group homes providing primarily welfare services.</p> <p>Both the numerator and denominator exclude:</p> <ul style="list-style-type: none">• Acute stroke patients separated from hospital who do not have a carer.• Acute stroke patients or carer(s) that refuse a formal assessment. <p>Presented as a percentage.</p>
Computation:	$(\text{Numerator} \div \text{denominator}) \times 100$
Numerator:	Number of patients with a final diagnosis of acute stroke whose carer(s) have had a documented formal needs assessment.
Denominator:	Number of patients with a final diagnosis of acute stroke who have a carer upon separation from hospital.

Representational attributes

Representation class:	Percentage
Data type:	Real
Unit of measure:	Episode
Format:	N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care