Third and Fourth Degree Perineal Tears Clinical Care Standard: 4a-Proportion of women who sustained a perineal tear during birth who received a genito-anal examination to assess the grade of the perineal tear



### © Australian Institute of Health and Welfare 2024

This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

# Third and Fourth Degree Perineal Tears Clinical Care Standard: 4a-Proportion of women who sustained a perineal tear during birth who received a genito-anal examination to assess the grade of the perineal tear

# Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

**Short name:** 4a-Proportion of women who sustained a perineal tear during birth who received a

genito-anal examination to assess the grade of the perineal tear

METEOR identifier: 728506

Registration status: Australian Commission on Safety and Quality in Health Care, Standard 20/04/2021

**Description:** The proportion of women who sustained a perineal tear during birth who received a

genito-anal examination to assess the grade of the perineal tear.

Indicator set: Clinical care standard indicators: Third and Fourth Degree Perineal Tears

Australian Commission on Safety and Quality in Health Care, Standard

20/04/2021

# Collection and usage attributes

Population group age

from:

18 yrs or older

Computation description: For the numerator, 'received' means that it is documented in the medical record

that the woman was provide with information about a suspected perineal tear and received a genito-anal examination to determine the degree of perineal injury or

declined the examination.

Presented as a percentage.

**Computation:** (Numerator ÷ Denominator) x 100

**Numerator:** The number of women who sustained a perineal tear who received a genito-anal

examination after birth or declined the offer of a genito-anal examination after birth.

**Denominator:** The number of women who sustained a perineal tear.

Include separations where an ICD code for perineal laceration was recorded (Table

A).

### Table A: ICD-10-AM codes for perineal laceration

Code	Description
O70.0	First degree perineal laceration during delivery
O70.1	Second degree perineal laceration during delivery
070.2	Third degree perineal laceration during delivery
O70.3	Fourth degree perineal laceration during delivery

# Representational attributes

Representation class: Percentage

Data type:RealUnit of measure:PersonFormat:N[NN]

### Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: Australian Commission on Safety and Quality in Health Care 2017. National Safety

and Quality Health Service Standards. 2nd ed. Australian Commission on Safety

and Quality in Health Care, Sydney.

Australian Commission on Safety and Quality in Health Care 2021. Third and Fourth Degree Perineal Tears clinical care standard. Australian Commission on

Safety and Quality in Health Care, Sydney.