

Antimicrobial stewardship: 8c-The proportion of patients who are administered prophylactic antimicrobials within the recommended time perioperatively

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Antimicrobial stewardship: 8c-The proportion of patients who are administered prophylactic antimicrobials within the recommended time perioperatively

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	8c-The proportion of patients who are administered prophylactic antimicrobials within the recommended time perioperatively
METEOR identifier:	736898
Registration status:	Australian Commission on Safety and Quality in Health Care, Standard 24/12/2020
Description:	The proportion of patients who are administered prophylactic antimicrobials within the recommended time perioperatively.
Indicator set:	Clinical care standard indicators: antimicrobial stewardship 2020 Australian Commission on Safety and Quality in Health Care, Standard 24/12/2020

Collection and usage attributes

Computation description:	<p>The numerator and denominator exclude patients undergoing surgery or a procedure for which there are no documented guidelines for the administration of prophylactic antimicrobials or where prophylaxis is not indicated due to the patient's current antimicrobial therapy. The numerator and denominator also exclude patients who have not received antimicrobials perioperatively.</p> <p>For the numerator, 'in accordance' means that the antimicrobial is administered within the recommended time period as per the current <i>Therapeutic Guidelines</i> or evidence-based, locally endorsed guidelines .</p> <p>If the patient undergoing surgery or a procedure receives multiple prophylactic antimicrobials, then the time of administration of all prophylactic antimicrobials should be in accordance with the guidelines in order for the patient to be counted in the numerator.</p> <p>Presented as a percentage.</p>
Computation:	$(\text{Numerator} \div \text{Denominator}) \times 100$
Numerator:	Number of patients who receive perioperative prophylactic antimicrobial within the recommended time period, in accordance with the current <i>Therapeutic Guidelines</i> or evidence-based, locally endorsed guidelines.
Denominator:	Number of patients undergoing surgery or a procedure for which there are documented guidelines on the use of prophylactic antimicrobials.

Comments: As per the *Antimicrobial Stewardship Clinical Care Standard* (ACSQHC 2020), the term 'locally endorsed guidelines' should meet the following specific requirements:

Therapeutic Guidelines: Antibiotic provides national, evidence-based guidelines for antimicrobial use in Australia. However, in some cases, evidence-based and formally endorsed local adaptations may be more relevant to local patterns of antimicrobial resistance or specific patient populations requiring a localised approach or with highly specialised diagnoses out of the scope of *Therapeutic Guidelines*. Ensure that local adaptations of guidelines are based on *Therapeutic Guidelines* and, where they deviate, a clear rationale based on published clinical evidence and local epidemiology is documented and endorsed by a governing body. Endorsing bodies may include peer review at a drug and therapeutics, antimicrobial stewardship or medicines advisory committee, or equivalent.

See the *Antimicrobial Stewardship Clinical Care Standard* (ACSQHC 2020) for key considerations to optimise the use of antimicrobials in surgical prophylaxis.

Services may select specific procedures to focus on as part of the audit process, based on their mix of patients and priorities.

Representational attributes

Representation class: Percentage
Data type: Real
Unit of measure: Service event
Format: N[NN]

Accountability attributes

Other issues caveats: Applicable settings:

All hospital settings, including public and private hospitals, subacute facilities, outpatient and day procedure services

General practice and other community and primary healthcare settings such as Hospital in the Home, Aboriginal community controlled health services, dental practices and community pharmacies

Residential aged care services.

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care
Reference documents: Australian Commission on Safety and Quality in Health Care. *Antimicrobial Stewardship Clinical Care Standard*. Sydney: ACSQHC; 2020.