

Delirium clinical care standard indicators: 5b- Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Delirium clinical care standard indicators: 5b- Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium

Identifying and definitional attributes

| | |
|-----------------------------|---|
| Metadata item type: | Indicator |
| Indicator type: | Indicator |
| Short name: | 5b-Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium |
| METEOR identifier: | 745820 |
| Registration status: | Australian Commission on Safety and Quality in Health Care, Standard 09/09/2021 |
| Description: | Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium. |
| Indicator set: | Clinical care standard indicators: delirium 2021 Australian Commission on Safety and Quality in Health Care, Standard 09/09/2021 |

Collection and usage attributes

| | |
|---------------------------------|--|
| Computation description: | For the numerator and denominator identify patients with delirium using the ICD codes specified in the Hospital-Acquired Complications (HACs) List - Specifications , worksheet 1, where complication group = 11. <i>Delirium</i> . For the numerator, the multicomponent interventions to treat and manage delirium must be documented in the patient's medical record and include tailored interventions to treat any identified cause/s of delirium. Presented as a percentage. |
| Computation: | $(\text{Numerator} \div \text{Denominator}) \times 100$ |
| Numerator: | Number of patients with delirium who received multicomponent interventions to treat and manage delirium. |
| Denominator: | The number of patients with delirium. |
| Comments: | This indicator is based on the Ontario Senior Friendly Hospital Strategy Delirium and Functional Decline Indicators (Wong et al, 2012). |

Representational attributes

| | |
|------------------------------|------------|
| Representation class: | Percentage |
| Data type: | Real |
| Unit of measure: | Person |
| Format: | N[NN] |

Source and reference attributes

| | |
|---------------------------------|---|
| Submitting organisation: | Australian Commission on Safety and Quality in Health Care |
| Reference documents: | Australian Commission on Safety and Quality in Health Care. Delirium Clinical Care Standard. Sydney: ACSQHC; 2021. Wong K, Tsang A, Liu B & Schwartz R. The Ontario Senior Friendly Hospital Strategy Delirium and Functional Decline Indicators. Toronto: Ontario Local Health Integration Network; 2012. |

