
Female—number of previous pregnancies (induced abortion), total NN



parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Female—number of previous pregnancies (induced abortion), total NN

Identifying and definitional attributes

Metadata item type: Data Element

Short name: Previous pregnancies—induced abortion

METEOR identifier: 269935

Registration status: Health, Standard 01/03/2005

Definition: The total number of previous pregnancies of a female resulting in induced abortion

(termination of pregnancy before 20 weeks' gestation).

Data Element Concept: Female—number of previous pregnancies

Value Domain: <u>Total pregnancies N[N]</u>

Value domain attributes

Representational attributes

Representation class: Total

Data type: Number

Format: N[N]

Maximum character length: 2

Value Meaning

Supplementary values: 99 Not stated

Unit of measure: Pregnancy

Data element attributes

Collection and usage attributes

Guide for use: A pregnancy resulting in multiple births should be counted as once pregnancy.

In multiple pregnancies with more than one type of outcome, the pregnancies should be recorded in the following order:

· all live births

• stillbirth

· spontaneous abortion

induced abortion

ectopic pregnancy

Where the outcome was one stillbirth and one live birth, count as stillbirth.

If a previous pregnancy was a hydatidiform mole, code as spontaneous or induced

abortion (or rarely, ectopic pregnancy), depending on the outcome.

Comments: The number of previous pregnancies is an important component of the woman's

reproductive history. Parity may be a risk factor for adverse maternal and perinatal

outcomes.

A previous history of induced abortion may increase the risk of some outcomes in

subsequent pregnancies.

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Related metadata Is re-engineered from Previous pregnancies, version 1, DE, NHDD, NHIMG,

references: Superseded 01/03/2005.pdf (15.1 KB)

No registration status