

Female—number of previous pregnancies (induced abortion), total NN

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Previous pregnancies—induced abortion
METEOR identifier:	269935
Registration status:	<ul style="list-style-type: none">• Health, Standard 01/03/2005
Definition:	The total number of previous pregnancies of a female resulting in induced abortion (termination of pregnancy before 20 weeks' gestation).

Data element concept attributes

Identifying and definitional attributes

Data element concept:	Female—number of previous pregnancies
METEOR identifier:	269454
Registration status:	<ul style="list-style-type: none">• Health, Standard 01/03/2005
Definition:	The total number of previous pregnancies.
Object class:	Female
Property:	Number of previous pregnancies

Value domain attributes

Identifying and definitional attributes

Value domain:	Total pregnancies N[N]
METEOR identifier:	270603
Registration status:	<ul style="list-style-type: none">• ACT Health (retired), Candidate 14/08/2018• Health, Standard 01/03/2005• Tasmanian Health, Standard 20/12/2016
Definition:	Total number of pregnancies.

Representational attributes

Representation class:	Total				
Data type:	Number				
Format:	N[N]				
Maximum character length:	2				
Supplementary values:	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99</td><td>Not stated</td></tr></tbody></table>	Value	Meaning	99	Not stated
Value	Meaning				
99	Not stated				

Unit of measure: Pregnancy

Data element attributes

Collection and usage attributes

Guide for use: A pregnancy resulting in multiple births should be counted as once pregnancy.

In multiple pregnancies with more than one type of outcome, the pregnancies should be recorded in the following order:

- all live births
- stillbirth
- spontaneous abortion
- induced abortion
- ectopic pregnancy

Where the outcome was one stillbirth and one live birth, count as stillbirth.

If a previous pregnancy was a hydatidiform mole, code as spontaneous or induced abortion (or rarely, ectopic pregnancy), depending on the outcome.


Comments: The number of previous pregnancies is an important component of the woman's reproductive history. Parity may be a risk factor for adverse maternal and perinatal outcomes.

A previous history of induced abortion may increase the risk of some outcomes in subsequent pregnancies.

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Related metadata references: Supersedes  [Previous pregnancies, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005.pdf](#) (15.1 KB) *No registration status*

© Australian Institute of Health and Welfare 2015–2022

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 3.0 (CC BY 3.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build upon this work. However, you must attribute the AIHW as the copyright holder of the work in compliance with our attribution policy available at www.aihw.gov.au/copyright. The full terms and conditions of this licence are available at <http://creativecommons.org/licenses/by3.0/au/>.

Enquiries relating to copyright should be addressed to the Head of the Communications, Media and Marketing Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.