

# Child—body mass index (self-reported), ratio NN[N].N[N]

## Identifying and definitional attributes

<b>Metadata item type:</b>	Data Element
<b>Short name:</b>	Body mass index—child (self-reported)
<b>METEOR identifier:</b>	270087
<b>Registration status:</b>	<ul style="list-style-type: none"><li>• <a href="#">Health</a>, Standard 01/03/2005</li></ul>
<b>Definition:</b>	A measure of a child's weight (body mass) relative to height used to assess the extent of weight excess where at least one of the measures is self reported.
<b>Data Element Concept:</b>	<a href="#">Child—body mass index</a>

## Value domain attributes

## Representational attributes

<b>Representation class:</b>	Ratio						
<b>Data type:</b>	Number						
<b>Format:</b>	NN[N].N[N]						
<b>Maximum character length:</b>	5						
<b>Supplementary values:</b>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>888.8</td><td>Unknown</td></tr><tr><td>999.9</td><td>Not reported</td></tr></tbody></table>	Value	Meaning	888.8	Unknown	999.9	Not reported
Value	Meaning						
888.8	Unknown						
999.9	Not reported						

## Data element attributes

## Collection and usage attributes

**Collection methods:** NN.N for BMI calculated from either self-reported height and/or self-reported weight.

BMI calculated from measured height and weight should be distinguished from BMI calculated from self-reported height and/or weight. When either self-reported height or self-reported weight is used in the calculation, BMI should be recorded as self-reported BMI. Self-reported or parentally reported height and weight for children and adolescents should be used cautiously if at all.

BMI should be derived after the data entry of weight and height. It should be stored on the raw data set as a continuous variable and should not be aggregated or rounded.

This metadata item applies to persons aged 2 years or older. It is recommended

## Comments:

This metadata item applies to persons aged 2 years or older. It is recommended for use in population surveys and health care settings for adults and population surveys only for children and adolescents. It is recommended that calculated BMI for children and adolescents be compared with a suitable growth reference such as the United States Centers for Disease Control 2000 BMI-for-age chart be used for in health care settings such as hospitals, clinics and in general practice. A BMI greater than the 85th percentile would be classified as overweight, while a BMI greater than the 95th percentile would be classified as obese. These percentiles are arbitrary and do not relate to morbidity as the BMI cut-points do in adults.

BMI is relatively easy to determine, and has been validated against more direct measures of adiposity such as Magnetic Resonance Imaging and Dual X-ray Absorptiometry.

BMI is a low cost technique, with low respondent and investigator burden. In addition, it offers low inter-observer and intra-observer error, therefore offering good reliability.

Overweight and obesity, as defined by the World Health Organisation (WHO) for the interpretation of BMI (WHO 2000), are exceedingly common in Australia and their prevalence is increasing.

It is recommended that in population surveys, sociodemographic data including ethnicity should be collected, as well as other risk factors including physiological status (e.g. pregnancy), physical activity, smoking and alcohol consumption. Summary statistics may need to be adjusted for these variables.

National health metadata items currently exist for sex, date of birth, country of birth, Indigenous status and smoking. Metadata items are being developed for physical activity.

Presentation of data:

Means, 95% confidence intervals, medians and centiles should be reported to one decimal place. Where the sample permits, population estimates should be presented by sex and 5-year age groups. Estimates based on sample surveys may need to take into account sampling weights.

For consistency with conventional practice, and for current comparability with international data sets, recommended centiles are 5, 10, 15, 25, 50, 75, 85, 90 and 95. To estimate the 5th and 95th centiles a sample size of at least 200 is recommended for each group for which the centiles are being specified.

Body mass index can be calculated from measured height and weight, or self-reported height and weight, however for children and adolescents, self-reported or parentally reported data should be used cautiously if at all.

For adults, body mass index tends to be underestimated when based on self-reported, rather than measured, height and weight. This is due to the fact that, on average, height tends to be overestimated and weight tends to be underestimated when self-reported by respondents.

There are many individuals for whom BMI is an inappropriate measure of body fatness. These are individuals whose high body mass is due to excess muscle rather than fat (e.g. body builders or others in whom the level of physical activity promotes an increase in muscle mass); or in those with osteoporosis who will have a lower than usual BMI; or those who have a different body build (e.g. individuals with unusually long or short legs or a different body fat distribution) (WHO Expert Committee 1995).

This is particularly important when assessing individuals but should also be taken into account in interpreting data from populations in which there are sub-groups with genetic or environmental differences in body build, composition, skeletal proportions or body fat distribution. As such, both BMI and a measure of fat distribution (waist circumference or waist: hip ratio) are important in calculating the risk of obesity comorbidities.

Epidemiological research shows that there is a strong association between BMI and health risk. Excess adipose tissue in adults is associated with excess morbidity and mortality from conditions such as hypertension, unfavourable blood lipid concentrations, diabetes mellitus, coronary heart disease, some cancers, gall

bladder disease, and osteoarthritis. It may also lead to social and economic disadvantage as well as psychosocial problems. It is a major public health issue in most industrialised societies.

Thinness (low BMI) is also an indicator of health risk, often being associated with general illness, anorexia, cigarette smoking, drug addiction and alcoholism. Low BMI is consistently associated with increased risk of osteoporosis and fractures in the elderly.

## Source and reference attributes


**Submitting organisation:** The Commonwealth Department of Health and Ageing based on the work of the consortium to develop an Australian standard definition of child/adolescent overweight and obesity; based at the Children Hospital at Westmead.

**Origin:** Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation. 2000. World Health Organization.

Cole TJ, Bellizzi MC, Flegal KM, Bietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *British Medical Journal* 2000; 320: 1240-1243

## Relational attributes

### Related metadata references:

Supersedes  [Body mass index, version 2, Derived DE, NHDD, NHIMG, Superseded 01/03/2005.pdf](#) (25.7 KB) *No registration status*

Is formed using [Person—height \(measured\), total centimetres NN\[N\].N](#)

- [Health](#), Standard 01/03/2005

Is formed using [Person—height \(self-reported\), total centimetres NN\[N\]](#)

- [Health](#), Standard 01/03/2005
- [Tasmanian Health](#), Standard 20/12/2016

Is formed using [Person—weight \(measured\), total kilograms N\[NN\].N](#)

- [Health](#), Superseded 12/12/2018

Is formed using [Person—weight \(self-reported\), total kilograms NN\[N\]](#)

- [Health](#), Standard 14/07/2005
- [Tasmanian Health](#), Standard 20/12/2016

See also [Person—body mass index \(classification\), code N\[.N\]](#)

- [Health](#), Standard 01/03/2005
- [Indigenous](#), Standard 13/03/2015

© Australian Institute of Health and Welfare 2015–2022

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 3.0 (CC BY 3.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build upon this work. However, you must attribute the AIHW as the copyright holder of the work in compliance with our attribution policy available at [www.aihw.gov.au/copyright](http://www.aihw.gov.au/copyright). The full terms and conditions of this licence are available at <http://creativecommons.org/licenses/by3.0/au/>.

Enquiries relating to copyright should be addressed to the Head of the Communications, Media and Marketing Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.