

Laboratory standard—upper limit of normal range of glycosylated haemoglobin, percentage N[N].N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Glycosylated haemoglobin—upper limit of normal range (percentage)
METEOR identifier:	270333
Registration status:	<ul style="list-style-type: none">Health, Standard 01/03/2005
Definition:	Laboratory standard for the value of glycosylated haemoglobin (HbA1c) measured as a percentage that is the upper boundary of the normal range.
Data Element Concept:	Laboratory standard—upper limit of normal range of glycosylated haemoglobin

Value domain attributes

Representational attributes

Representation class:	Percentage				
Data type:	Number				
Format:	N[N].N				
Maximum character length:	3				
Supplementary values:	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99.9</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	99.9	Not stated/inadequately described
Value	Meaning				
99.9	Not stated/inadequately described				

Data element attributes

Collection and usage attributes

Guide for use:	Record the upper limit of the HbA1c normal reference range from the laboratory result.
Collection methods:	This value is usually notified in patient laboratory results and may vary for different laboratories.
Comments:	HbA1c results vary between laboratories; use the same laboratory for repeated testing.

Source and reference attributes

Submitting organisation:	National Diabetes Data Working Group
Origin:	National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary.

Relational attributes

Related metadata references:

Supersedes  [Glycosylated Haemoglobin \(HbA1c\) - upper limit of normal range, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005.pdf](#) (15.9 KB) *No registration status*

See also [Person—glycosylated haemoglobin level \(measured\), percentage N\[N\].N](#)

- [Health](#), Standard 01/03/2005

Implementation in Data Set Specifications:

[Acute coronary syndrome \(clinical\) DSSHealth](#), Superseded 01/09/2012

[Acute coronary syndrome \(clinical\) DSSHealth](#), Superseded 02/05/2013

[Acute coronary syndrome \(clinical\) NBPDS Health](#), Recorded 15/05/2017

[Acute coronary syndrome \(clinical\) NBPDS 2013-Health](#), Standard 02/05/2013

Implementation start date: 01/07/2013

[Diabetes \(clinical\) DSSHealth](#), Superseded 21/09/2005

[Diabetes \(clinical\) NBPDSHealth](#), Standard 21/09/2005

DSS specific information:

HbA1c is a measurement of long-term blood glucose control and is used to assess the effectiveness of treatment. It is a convenient way to obtain an integrated assessment of antecedent glycaemia over an extended period under real life conditions and is used as a standard for assessing overall blood glucose control. The target is to achieve an HbA1c within 1% of the upper limit of normal or achieve control as near to this target as possible without producing unacceptable hypoglycaemia as recommended from the Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus.

If HbA1c is 2% above the upper limit of normal, explore reasons for unsatisfactory control such as diet, intercurrent illness, appropriateness of medication, concurrent medication, stress, and exercise and review management:

- review and adjust treatment
- consider referral to diabetes educator
- consider referral to dietitian
- consider referral to endocrinologist or physician or diabetes centre.

[Diabetes \(clinical\) NBPDSHealth](#), Recorded 15/05/2017