

# Concurrent acute coronary syndrome clinical condition code NN

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Concurrent acute coronary syndrome clinical condition code NN

## Identifying and definitional attributes

<b>Metadata item type:</b>	Value Domain
<b>METEOR identifier:</b>	284887
<b>Registration status:</b>	<a href="#">Health</a> , Superseded 01/10/2008
<b>Definition:</b>	A code set describing medical conditions pertinent to the risk stratification and treatment of acute coronary syndrome.

## Representational attributes

<b>Representation class:</b>	Code
<b>Data type:</b>	Number
<b>Format:</b>	NN
<b>Maximum character length:</b>	2

	Value	Meaning
<b>Permissible values:</b>	11	Angina for more than last two weeks
	12	Angina only in the last two weeks
	21	Chronic lung disease
	31	Heart failure
	41	Hypertension
	51	Ischaemic: non-haemorrhagic cerebral infarction
	52	Haemorrhagic: intracerebral haemorrhage
	61	Peripheral artery disease
	62	Aortic aneurysm
	63	Renal artery stenosis
<b>Supplementary values:</b>	71	Sleep apnoea
	99	not stated/inadequately described

## Collection and usage attributes

<b>Guide for use:</b>	Angina:
	CODE 11 Angina for more than last two weeks
	This code is used where there are symptoms, which can be described as chest pain or pressure, jaw pain, arm pain, or other equivalent discomfort suggestive of cardiac ischaemia, for more than the last two weeks.
	CODE 12 Angina only in the last two weeks
	This code is used where there are symptoms, which can be described as chest pain or pressure, jaw pain, arm pain, or other equivalent discomfort suggestive of cardiac ischaemia, only in the last two weeks.
	Chronic lung disease:
	CODE 21 Chronic lung disease
	This code is used where there is a history or symptoms suggestive of chronic lung

disease.

Heart failure:

CODE 31 Heart failure

This code is used where a patient has past or current symptoms of heart failure (typically breathlessness or fatigue), either at rest or during exercise and/or signs of pulmonary or peripheral congestion suggestive of cardiac dysfunction.

Hypertension:

CODE 41 Hypertension

This code is used where there is current use of pharmacotherapy for hypertension and/or clinical evidence of high blood pressure.

Stroke:

CODE 51 Ischaemic: non-haemorrhagic cerebral infarction

This code is used if there is history of stroke or cerebrovascular accident (CVA) resulting from an ischaemic event where the patient suffered a loss of neurological function with residual symptoms remaining for at least 24 hours.

CODE 52 Haemorrhagic: intracerebral haemorrhage

This code is used if there is history of stroke or cerebrovascular accident (CVA) resulting from a haemorrhagic event where the patient suffered a loss of neurological function with residual symptoms remaining for at least 24 hours.

Peripheral arterial disease:

CODE 61 Peripheral artery disease

This code is used where there is history of either chronic or acute occlusion or narrowing of the arterial lumen in the aorta or extremities.

CODE 62 Aortic aneurysm

This code is used where there is a history of aneurysmal dilatation of the aorta (thoracic and or abdominal).

CODE 63 Renal artery stenosis

This code is used where there is history of functional stenosis of one or both renal arteries.

Sleep Apnoea syndrome:

CODE 71 Sleep apnoea

This code is used where there is evidence of sleep apnoea syndrome (SAS) on history.

## Source and reference attributes

**Submitting organisation:** Australian Institute of Health and Welfare

## Relational attributes

**Related metadata references:** Has been superseded by [Acute coronary syndrome related medical history code NN](#)  
[Health](#), Standard 01/10/2008

**Data elements implementing this value domain:** [Person—acute coronary syndrome concurrent clinical condition, code NN](#)  
[Health](#), Superseded 01/10/2008

