Episode of community based palliative care—phase of care, code N

| Exported from METEOR (AIHW's Metadata Online Registry) |
|--|
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| © Australian Institute of Health and Welfare 2024 |
| This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties. |
| You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our |

attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Page 1 of 4

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at

meteor@aihw.gov.au.

Metadata 287942

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Episode of community based palliative care—phase of care, code N

Identifying and definitional attributes

Metadata item type: Data Element

Short name: Phase of palliative care

METEOR identifier: 287942

Registration status: Health, Recorded 13/05/2008

Definition: The client's stage of illness/situation in terms of four recognised phases of palliative

care, as represented by a code.

Context: Palliative care delivered in community settings

Data Element Concept: Episode of community based palliative care—phase of care

Value Domain: Phase of palliative care code N

Value domain attributes

Representational attributes

Representation class: Code

Data type: Number

Format: N

Maximum character length: 1

ValueMeaningPermissible values:1Stable2Unstable3Deteriorating4Terminal

Collection and usage attributes

Guide for use: CODE 1 Stable phase

The patient's symptoms are adequately controlled by established management. Further interventions to maintain symptom control and quality of life have been planned.

The situation of the carer(s)/family/friends is relatively stable and no new issues are apparent. Any needs are met by the established plan of care.

CODE 2 Unstable phase

The patient experiences the development of a new problem or a rapid increase in the severity of existing problems, either of which require an urgent change in management or emergency treatment.

The carer(s)/family/friends experience a sudden change in their situation requiring urgent intervention by members of the multidisciplinary team.

CODE 3 Deteriorating phase

The patient experiences a gradual worsening of existing symptoms or the development of new but expected problems. These require the application of specific plans of care and regular review but not urgent or emergency treatment.

The carer(s)/family/friends experience gradually worsening distress and other difficulties, including social and practical difficulties, as a result of the illness of the patient. This requires a planned support program and counselling is necessary.

CODE 4 Terminal care phase

Death is likely in a matter of days and no acute intervention is planned or required. The typical features of a person in this phase may include the following:

- · Profoundly weak
- · Essentially bed bound
- Drowsy for extended periods
- · Disorientated for time and has a severely limited attention span
- Increasingly disinterested in food and drink
- · Finding it difficult to swallow medication
- This requires the use of frequent, usually daily, interventions aimed at physical, emotional and spiritual issues.

The carer(s)/family/friends recognise that death is imminent and care is focussed on emotional and spiritual issues as a prelude to bereavement.

There is a fifth phase, the bereavement phase, that is usually included as one of the recognised phases of care. As the bereavement phase applies to the period after the person with the life-limiting illness has died, this phase is not relevant to current

agreed data collections.

agre

Source and reference attributes

Submitting organisation: Palliative Care Intergovernmental Forum

Origin: Eagar K et al (1997) The Australian National Sub-Acute and Non-Acute Patient

(AN-SNAP) Casemix Classification: report of the National Sub-Acute and Non-Acute Casemix Classification Study. Centre for Health Service Development,

University of Wollongong.

Data element attributes

Collection and usage attributes

Guide for use:

Comments:

Collection methods: Record the client's phase of care at the first assessment by the agency which

initiates the episode of community-based palliative care.

Source and reference attributes

Submitting organisation: Palliative Care Intergovernmental Forum