

Person—C-reactive protein level (measured), total milligrams per litre N[NN].N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	C-reactive protein level (measured)
Synonymous names:	CRP measured
METEOR identifier:	338256
Registration status:	<ul style="list-style-type: none">• Health, Standard 01/10/2008
Definition:	A person's serum C-reactive protein (CRP) level, measured in milligrams per litre.
Data Element Concept:	Person—C-reactive protein level (measured)

Value domain attributes

Representational attributes

Representation class:	Total				
Data type:	Number				
Format:	N[NN].N				
Maximum character length:	4				
Supplementary values:	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>999.9</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	999.9	Not stated/inadequately described
Value	Meaning				
999.9	Not stated/inadequately described				
Unit of measure:	Milligram per litre (mg/L)				
Unit of measure precision:	1				

Data element attributes

Collection and usage attributes

Comments:	<p>The value should be recorded on a high sensitivity assay.</p> <p>CRP is used in the assessment of acute phase reaction in inflammatory, infective and neoplastic disorders.</p>
------------------	--

Source and reference attributes

Reference documents:	<p>The Royal College of Pathologists of Australia Version 4.0 12th March 2004 (last accessed 12May 2006). http://www.rcpamannual.edu.au/sections/pathologytest.asp?s=33&i=468</p>
-----------------------------	--

Relational attributes

Related metadata references:

See also [Person—C-reactive protein level measured date, DDMMYYYY](#)

- [Health](#), Standard 01/10/2008

See also [Person—C-reactive protein level measured time, hhmm](#)

- [Health](#), Standard 01/10/2008

Implementation in Data Set Specifications:

[Acute coronary syndrome \(clinical\) DSSHealth](#), Superseded 01/09/2012

DSS specific information:

The C-reactive protein (CRP) level should be measured as early as possible following presentation to the hospital.

[Acute coronary syndrome \(clinical\) DSSHealth](#), Superseded 02/05/2013

DSS specific information:

The C-reactive protein (CRP) level should be measured as early as possible following presentation to the hospital.

[Acute coronary syndrome \(clinical\) NBPDS Health](#), Recorded 15/05/2017

DSS specific information:

The C-reactive protein (CRP) level should be measured as early as possible following presentation to the hospital.

[Acute coronary syndrome \(clinical\) NBPDS 2013-Health](#), Standard 02/05/2013

Implementation start date: 01/07/2013

DSS specific information:

The C-reactive protein (CRP) level should be measured as early as possible following presentation to the hospital.