

Person—C-reactive protein level (measured), total milligrams per litre N[NN].N

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Person—C-reactive protein level (measured), total milligrams per litre N[NN].N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	C-reactive protein level (measured)
Synonymous names:	CRP measured
METEOR identifier:	338256
Registration status:	Health , Standard 01/10/2008
Definition:	A person's serum C-reactive protein (CRP) level, measured in milligrams per litre.
Data Element Concept:	Person—C-reactive protein level (measured)
Value Domain:	Total milligrams per litre N[NN].N

Value domain attributes

Representational attributes

Representation class:	Total	
Data type:	Number	
Format:	N[NN].N	
Maximum character length:	4	
	Value	Meaning
Supplementary values:	999.9	Not stated/inadequately described
Unit of measure:	Milligram per litre (mg/L)	

Data element attributes

Collection and usage attributes

Comments:	The value should be recorded on a high sensitivity assay. CRP is used in the assessment of acute phase reaction in inflammatory, infective and neoplastic disorders.
------------------	---

Source and reference attributes

Reference documents:	The Royal College of Pathologists of Australia Version 4.0 12th March 2004 (last accessed 12May 2006). http://www.rcpamanual.edu.au/sections/pathologytest.asp?s=33&i=468
-----------------------------	--

Relational attributes

Related metadata references:	See also Person—C-reactive protein level measured date, DDMMYYYY Health , Standard 01/10/2008 See also Person—C-reactive protein level measured time, hhmm Health , Standard 01/10/2008
-------------------------------------	--

Implementation in Data Set Specifications:

[Acute coronary syndrome \(clinical\) DSS](#)

[Health](#), Superseded 01/09/2012

DSS specific information: The C-reactive protein (CRP) level should be measured as early as possible following presentation to the hospital.

[Acute coronary syndrome \(clinical\) DSS](#)

[Health](#), Superseded 02/05/2013

DSS specific information: The C-reactive protein (CRP) level should be measured as early as possible following presentation to the hospital.

[Acute coronary syndrome \(clinical\) NBPDS 2013-](#)

[Health](#), Standard 02/05/2013

Implementation start date: 01/07/2013

DSS specific information:

The C-reactive protein (CRP) level should be measured as early as possible following presentation to the hospital.