



National Health Data Dictionary volume 6

Data elements S to Y (by short name)
Generated on 05/01/2007

© Australian Institute of Health and Welfare 2006

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced without prior written permission from the Australian Institute of Health and Welfare. Requests and enquiries concerning reproduction and rights should be directed to the Head, Business Promotion and Media, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.

Any enquiries about or comments on this publication should be directed to:

National Data Development and Standards Unit
Australian Institute of Health and Welfare
GPO Box 570
Canberra ACT 2601
Email: datadevelopment@aihw.gov.au
Phone: (02) 6244 1222 Fax: (02) 6244 1166

List of metadata items

Data Elements	9
Satisfaction with participation	10
Scheduled admission date	13
Separation date	14
Separation time	18
Separations	19
Service contact date	21
Service mode (non-admitted patient)	23
Service type (non-admitted patient)	25
Sex	29
Source of referral to alcohol and other drug treatment service	34
Source of referral to public psychiatric hospital	37
Specialised mental health service program type	39
Specialised mental health service setting	42
Specialised mental health service target population	44
Specialised mental health service – hours staffed	47
Specialised mental health service – supported public housing places	49
Specialised service indicators	51
Specialised service indicators – acquired immune deficiency syndrome unit	52
Specialised service indicators – acute renal dialysis unit	54
Specialised service indicators – acute spinal cord injury unit	56
Specialised service indicators – alcohol and drug unit	58
Specialised service indicators – bone marrow transplantation unit	60
Specialised service indicators – burns unit (level III)	62
Specialised service indicators – cardiac surgery unit	64
Specialised service indicators – clinical genetics unit	66
Specialised service indicators – comprehensive epilepsy centre	68
Specialised service indicators – coronary care unit	70
Specialised service indicators – diabetes unit	72
Specialised service indicators – domiciliary care service	74
Specialised service indicators – geriatric assessment unit	76
Specialised service indicators – heart, lung transplantation unit	78
Specialised service indicators – hospice care unit	80
Specialised service indicators – in-vitro fertilisation unit	82
Specialised service indicators – infectious diseases unit	84
Specialised service indicators – intensive care unit (level III)	86
Specialised service indicators – liver transplantation unit	88
Specialised service indicators – maintenance renal dialysis centre	90
Specialised service indicators – major plastic/reconstructive surgery unit	92
Specialised service indicators – neonatal intensive care unit (level III)	94
Specialised service indicators – neuro surgical unit	96
Specialised service indicators – nursing home care unit	98
Specialised service indicators – obstetric/maternity	100
Specialised service indicators – oncology unit, cancer treatment	102
Specialised service indicators – pancreas transplantation unit	104
Specialised service indicators – psychiatric unit/ward	106
Specialised service indicators – rehabilitation unit	108
Specialised service indicators – renal transplantation unit	110
Specialised service indicators – sleep centre	112

Specialised service indicators – specialist paediatric.....	114
Specialist private sector rehabilitation care indicator.....	116
Staging basis of cancer	118
Staging scheme source	120
Staging scheme source edition number.....	122
State/Territory of birth.....	124
Status of the baby	126
Street name (person)	128
Street name (service provider organisation).....	130
Street suffix code (person).....	132
Street suffix code (service provider organisation)	134
Street type code (person).....	136
Street type code (service provider organisation)	138
Suburb/town/locality name (person).....	140
Suburb/town/locality name (service provider organisation)	142
Surgical specialty	144
Surgical treatment procedure for cancer	146
Systemic therapy agent name	148
Teaching status	150
Telephone number	152
Telephone number type.....	154
Time creatine kinase MB isoenzyme measured	156
Time of first angioplasty balloon inflation or stenting.....	158
Time of intravenous fibrinolytic therapy	160
Time of triage	162
Time patient presents.....	164
Time troponin measured	166
Tobacco smoking status.....	167
Tobacco smoking status (diabetes mellitus)	170
Tobacco smoking – consumption/quantity (cigarettes)	172
Tobacco smoking – duration (daily smoking).....	175
Tobacco smoking – ever daily use.....	177
Tobacco smoking – frequency.....	179
Tobacco smoking – product	181
Tobacco smoking – quit age (daily smoking)	183
Tobacco smoking – start age (daily smoking)	185
Tobacco smoking – time since quitting (daily smoking).....	187
Total contract patient days.....	189
Total hours worked by a medical practitioner	191
Total leave days	193
Total number of group sessions	196
Total occasions of service	198
Total psychiatric care days.....	200
Treatment delivery setting for alcohol and other drugs	204
Triglyceride level (measured).....	206
Troponin assay type	210
Troponin assay – upper limit of normal range (micrograms per litre)	212
Troponin level (measured).....	214
Tumour size at diagnosis (solid tumours)	216
Tumour thickness at diagnosis (melanoma).....	218

Type and sector of employment establishment.....	219
Type of accommodation	222
Type of augmentation of labour	225
Type of labour induction.....	227
Type of usual accommodation.....	228
Type of visit to emergency department.....	230
Urgency of admission	232
Vascular history	235
Vascular procedures.....	237
Visual acuity (left eye)	239
Visual acuity (right eye).....	242
Waist circumference (measured).....	245
Waist circumference risk indicator - adults	249
Waist-to-hip ratio.....	252
Waiting list category	255
Waiting time at a census date	260
Waiting time at removal from elective surgery waiting list.....	263
Weight (self-reported).....	266
Weight in grams (measured).....	269
Weight in kilograms (measured).....	271
Year insulin started	277
Year of arrival in Australia.....	279
Year of diagnosis of diabetes mellitus	281

Data Element Technical Names

Admitted patient care waiting list episode – scheduled admission date, DDMMYYYY	13
Admitted patient hospital stay – number of patient days (of contracted care), total N[NN]	189
Adult – waist circumference risk indicator, Caucasian adult code N	249
Adult – waist-to-hip ratio, N.NN	252
Birth event – labour augmentation type, code N	225
Birth event – labour induction type, code N	227
Birth event – state/territory of birth, code N	124
Birth – birth status, code N	126
Cancer staging – cancer staging scheme source edition number, code N[N]	122
Cancer staging – cancer staging scheme source, code N	120
Cancer staging – staging basis of cancer, code A	118
Cancer treatment – surgical procedure for cancer, procedure code (ACHI 5th edn) NNNNNN-NN	146
Cancer treatment – systemic therapy agent name (primary cancer), antineoplastic drug code (Self-Instructional Manual for Tumour Registrars Book 8 3rd edn) X[X(39)]	148
Elective care waiting list episode – elective care type, code N	255
Elective surgery waiting list episode – surgical specialty (of scheduled doctor), code NN	144
Elective surgery waiting list episode – waiting time (at a census date), total days N[NNN]	260
Elective surgery waiting list episode – waiting time (at removal), total days N[NNN]	263
Episode of admitted patient care – admission urgency status, code N	232
Episode of admitted patient care – number of leave days, total N[NN]	193
Episode of admitted patient care – referral source, public psychiatric hospital code NN	37
Episode of admitted patient care – separation date, DDMMYYYY	14
Episode of admitted patient care – separation time, hhmm	18
Episode of care – number of psychiatric care days, total N[NNNN]	200
Episode of care – specialist private sector rehabilitation care indicator, code N	116
Episode of treatment for alcohol and other drugs – referral source, code NN	34
Episode of treatment for alcohol and other drugs – service delivery setting, code N	204
Establishment – number of group sessions, total N[NNNNN]	196
Establishment – number of occasions of service, total N[NNNNNN]	198
Establishment – number of separations (financial year), total N[NNNNN]	19
Establishment – specialised service indicator (acquired immune deficiency syndrome unit), yes/no code N	52
Establishment – specialised service indicator (acute renal dialysis unit), yes/no code N	54
Establishment – specialised service indicator (acute spinal cord injury unit), yes/no code N	56
Establishment – specialised service indicator (alcohol and drug unit), yes/no code N	58
Establishment – specialised service indicator (bone marrow transplantation unit), yes/no code N	60
Establishment – specialised service indicator (burns unit (level III)), yes/no code N	62
Establishment – specialised service indicator (cardiac surgery unit), yes/no code N	64
Establishment – specialised service indicator (clinical genetics unit), yes/no code N	66
Establishment – specialised service indicator (comprehensive epilepsy centre), yes/no code N	68
Establishment – specialised service indicator (coronary care unit), yes/no code N	70
Establishment – specialised service indicator (diabetes unit), yes/no code N	72
Establishment – specialised service indicator (domiciliary care service), yes/no code N	74
Establishment – specialised service indicator (geriatric assessment unit), yes/no code N	76
Establishment – specialised service indicator (heart, lung transplantation unit), yes/no code N	78
Establishment – specialised service indicator (hospice care unit), yes/no code N	80
Establishment – specialised service indicator (infectious diseases unit), yes/no code N	84
Establishment – specialised service indicator (intensive care unit (level III)), yes/no code N	86

Establishment – specialised service indicator (in-vitro fertilisation unit), yes/no code N	82
Establishment – specialised service indicator (liver transplantation unit), yes/no code N.....	88
Establishment – specialised service indicator (maintenance renal dialysis centre), yes/no code N ..	90
Establishment – specialised service indicator (major plastic/reconstructive surgery unit), yes/no code N	92
Establishment – specialised service indicator (neonatal intensive care unit (level III)), yes/no code N	94
Establishment – specialised service indicator (neuro surgical unit), yes/no code N	96
Establishment – specialised service indicator (nursing home care unit), yes/no code N.....	98
Establishment – specialised service indicator (obstetric/maternity), yes/no code N	100
Establishment – specialised service indicator (oncology unit) (cancer treatment), yes/no code N..	102
Establishment – specialised service indicator (pancreas transplantation unit), yes/no code N	104
Establishment – specialised service indicator (psychiatric unit/ward), yes/no code N.....	106
Establishment – specialised service indicator (rehabilitation unit), yes/no code N.....	108
Establishment – specialised service indicator (renal transplantation unit), yes/no code N.....	110
Establishment – specialised service indicator (sleep centre), yes/no code N	112
Establishment – specialised service indicator (specialist paediatric), yes/no code N	114
Establishment – specialised service indicator (transplantation unit), yes/no code N.....	51
Establishment – teaching status (university affiliation), code N.....	150
Health professional – establishment type (employment), industry code NN	219
Health service event – presentation time, hhmm.....	164
Laboratory standard – upper limit of normal range for troponin assay, total micrograms per litre N[NNN].....	212
Medical practitioner – hours worked, total NNN.....	191
Non-admitted patient emergency department service episode – triage time, hhmm	162
Non-admitted patient emergency department service episode – type of visit to emergency department, code N.....	230
Non-admitted patient service event – service event type (clinical), code N[N]	25
Non-admitted patient service event – service mode, hospital code N{N}.....	23
Patient – diagnosis date (diabetes mellitus), YYYY	281
Patient – insulin start date, YYYY	277
Person (address) – street name, text [A(30)]	128
Person (address) – street suffix, code A[A].....	132
Person (address) – street type, code A[AAA].....	136
Person (address) – suburb/town/locality name, text [A(50)].....	140
Person (overseas born) – year of first arrival in Australia, date YYYY	279
Person (telephone) – telephone number type, code A	154
Person with cancer – melanoma thickness (at diagnosis), total millimetres NNN.NN.....	218
Person with cancer – solid tumour size (at diagnosis), total millimetres NNN	216
Person – accommodation type (prior to admission), code N	228
Person – accommodation type (usual), code N[N]	222
Person – creatine kinase myocardial band isoenzyme measured time, hhmm	156
Person – first angioplasty balloon inflation or stenting time, hhmm.....	158
Person – intravenous fibrinolytic therapy time, hhmm	160
Person – level of satisfaction with participation in a life area, code N.....	10
Person – number of cigarettes smoked (per day), total N[N].....	172
Person – regular tobacco smoking indicator (last 3 months), code N	170
Person – sex, code N.....	29
Person – telephone number, text [X(40)]	152
Person – time since quitting tobacco smoking (daily smoking), code NN.....	187
Person – tobacco product smoked, code N	181

Person—tobacco smoking daily use status, code N	177
Person—tobacco smoking duration (daily smoking), total years N[N].....	175
Person—tobacco smoking frequency, code N	179
Person—tobacco smoking quit age (daily smoking), total years NN	183
Person—tobacco smoking start age (daily smoking), total years NN.....	185
Person—tobacco smoking status, code N	167
Person—triglyceride level (measured), total millimoles per litre N[N].N	206
Person—troponin assay type, code N.....	210
Person—troponin level (measured), total micrograms per litre NN.NN	214
Person—troponin level measured time, hhmm	166
Person—vascular condition status (history), code NN	235
Person—vascular procedures (history), code NN	237
Person—visual acuity (left eye), code NN	239
Person—visual acuity (right eye), code NN	242
Person—waist circumference (measured), total centimetres NN[N].N.....	245
Person—weight (measured), total grams NNNN.....	269
Person—weight (measured), total kilograms N[NN].N	271
Person—weight (self-reported), total kilograms NN[N]	266
Service contact—service contact date, DDMMYYYY	21
Service provider organisation (address)—street name, text [A(30)]	130
Service provider organisation (address)—street suffix, code A[A].....	134
Service provider organisation (address)—street type, code A[AAA].....	138
Service provider organisation (address)—suburb/town/locality name, text [A(50)].....	142
Specialised mental health service—admitted patient care program type, code N.....	39
Specialised mental health service—number of hours staffed, average hours NN	47
Specialised mental health service—number of supported public housing places, total N[N(5)]	49
Specialised mental health service—service setting, code N	42
Specialised mental health service—target population group, code N	44

Data Elements

Satisfaction with participation

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person—level of satisfaction with participation in a life area, code N
<i>METeOR identifier:</i>	320216
<i>Registration status:</i>	NHIG, Standard 29/11/2006 NCSIMG, Standard 16/10/2006
<i>Definition:</i>	The degree to which a person is satisfied with their involvement in a specified life area, as represented by a code.
<i>Context:</i>	Human functioning and disability

Data element concept attributes

<i>Data element concept:</i>	Person—level of satisfaction with participation in a life area
<i>Definition:</i>	A person's level of satisfaction with their involvement in a life area, in relation to their current life goals.
<i>Object class:</i>	Person
<i>Property:</i>	Level of satisfaction with participation in a life area

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																		
<i>Data type:</i>	Number																		
<i>Format:</i>	N																		
<i>Maximum character length:</i>	1																		
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>0</td><td>High satisfaction with participation</td></tr><tr><td>1</td><td>Moderate satisfaction with participation</td></tr><tr><td>2</td><td>Neither satisfied nor dissatisfied with participation</td></tr><tr><td>3</td><td>Moderate dissatisfaction with participation</td></tr><tr><td>4</td><td>Extreme dissatisfaction with participation</td></tr><tr><td>5</td><td>Complete restriction and dissatisfaction</td></tr><tr><td>8</td><td>Not specified</td></tr><tr><td>9</td><td>Not applicable</td></tr></tbody></table>	Value	Meaning	0	High satisfaction with participation	1	Moderate satisfaction with participation	2	Neither satisfied nor dissatisfied with participation	3	Moderate dissatisfaction with participation	4	Extreme dissatisfaction with participation	5	Complete restriction and dissatisfaction	8	Not specified	9	Not applicable
Value	Meaning																		
0	High satisfaction with participation																		
1	Moderate satisfaction with participation																		
2	Neither satisfied nor dissatisfied with participation																		
3	Moderate dissatisfaction with participation																		
4	Extreme dissatisfaction with participation																		
5	Complete restriction and dissatisfaction																		
8	Not specified																		
9	Not applicable																		
<i>Supplementary values:</i>																			

Collection and usage attributes

<i>Guide for use:</i>	<p>This metadata item contributes to the definition of the concept 'Disability' and gives an indication of the experience of disability for a person.</p> <p>In the context of health, participation is involvement in a life situation. Participation restrictions are problems an individual may experience in involvement in life situations.</p> <p>This metadata item gives a rating of the person's degree of satisfaction with participation in a domain of life, in relation to their current life goals. Satisfaction with participation</p>
-----------------------	---

corresponds to the person's own perspective on their participation, and reflects their attitude to their participation in the various life areas. It is essentially a summary measure in which are embedded the concepts of choice, opportunity and importance.

CODE 0 High satisfaction with participation

Used if a person is involved in the specified life situation as he or she wishes to fulfil his or her current life goals in terms of duration, frequency, manner and outcome.

CODE 1 Moderate satisfaction with participation

Used if the person is reasonably satisfied with their participation in this life situation, in terms of duration, frequency, manner and outcome. This could occur if one of the criteria (duration, frequency, manner or outcome) is not fulfilled and that criterion is not critical to the person's goals. For example, the person does not participate in the specified life situation as frequently as wished, but the other criteria are met and the frequency is not so affected that it is critical to the person's satisfaction.

CODE 2 Neither satisfied nor dissatisfied with participation

Used if the person is neither satisfied nor dissatisfied with their participation in this life situation, in terms of duration, frequency, manner and outcome.

CODE 3 Moderate dissatisfaction with participation

Used if two or three criteria (duration, frequency, manner or outcome) are not fulfilled, but are not so badly affected, in relation to the person's goals in that life area, that the person is extremely dissatisfied. For example, a person is able to participate in work, but is placed in supported employment rather than employment in the open labour market. This is not in line with the person's goals, so that the manner and outcome of the participation are not fulfilled.

CODE 4 Extreme dissatisfaction with participation

Used when all criteria (duration, frequency, manner and outcome) are not fulfilled for the specified life situation, or where any of the criteria are so badly affected in relation to the person's goals that they consider themselves to be extremely dissatisfied with this life area. An example of the latter would arise when a person is extremely dissatisfied with participation in interpersonal activities because his/her goal in terms of duration of social visits is never fulfilled, although other criteria (frequency and manner) may be fulfilled.

CODE 5 Complete restriction and dissatisfaction

Used when the person does not participate in this life situation in line with his or her own goals, i.e. in an area where they wish to participate and is completely dissatisfied with not participating in this life situation.

CODE 9 Not applicable

Used when participation in a life situation is not relevant, such as employment of an infant or where there is no participation and the person has no desire to participate in this area. For example, a personal preference not to participate in specific areas of community, social and civic life such as sport or hobbies. The area may not be applicable to the person's current life goals.

Source and reference attributes

<i>Submitting organisation:</i>	Australian Institute of Health and Welfare (AIHW) which is the Australian Collaborating Centre for the World Health Organization Family of International Classifications.
<i>Origin:</i>	WHO 2001. ICF: International Classification of Functioning, Disability and Health. Geneva: WHO AIHW 2003. ICF Australian User Guide Version 1.0. Canberra: AIHW
<i>Reference documents:</i>	Further information on the ICF, including more detailed codes, can be found in the ICF itself and the ICF Australian User Guide (AIHW 2003), at the following websites: <ul style="list-style-type: none">• WHO ICF website http://www.who.int/classifications/icf/en/• Australian Collaborating Centre ICF website http://www.aihw.gov.au/disability/icf/index.html

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Satisfaction with participation should be coded from the perspective of the person. This data element should be coded in conjunction with the Person – activities and participation life area, code (ICF 2001) AN[NNN] data element. For example, a person's 'moderate satisfaction with participation in exchange of information'.
-----------------------	---

Source and reference attributes

<i>Submitting organisation:</i>	Australian Institute of Health and Welfare (AIHW) which is the Australian Collaborating Centre for the World Health Organization Family of International Classifications.
---------------------------------	---

Relational attributes

<i>Implementation in Data Set Specifications:</i>	Activities and Participation cluster NHIG, Standard 29/11/2006 NCSIMG, Standard 16/10/2006
---	---

Scheduled admission date

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Admitted patient care waiting list episode – scheduled admission date, DDMMYYYY
<i>METeOR identifier:</i>	269978
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The date on which it is proposed that a patient on the waiting list will be admitted for an episode of care.

Data element concept attributes

<i>Data element concept:</i>	Admitted patient care waiting list episode – scheduled admission date
<i>Definition:</i>	The date on which it is proposed that a patient on the waiting list will be admitted for an episode of care.
<i>Object class:</i>	Admitted patient care waiting list episode
<i>Property:</i>	Scheduled admission date

Value domain attributes

Representational attributes

<i>Representation class:</i>	Date
<i>Data type:</i>	Date/Time
<i>Format:</i>	DDMMYYYY
<i>Maximum character length:</i>	8

Data element attributes

Collection and usage attributes

<i>Comments:</i>	<p>If this metadata item were to be used to compare different hospitals or geographical locations, it would be necessary to specify when the scheduled date is to be allocated (for example, on addition to the waiting list).</p> <p>This metadata item is required for the purposes of hospital management - allocation of beds, operating theatre time and other resources.</p>
------------------	--

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Scheduled admission date, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005
-------------------------------------	--

Separation date

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of admitted patient care – separation date, DDMMYYYY
<i>METeOR identifier:</i>	270025
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Date on which an admitted patient completes an episode of care.

Data element concept attributes

<i>Data element concept:</i>	Episode of admitted patient care – separation date
<i>Definition:</i>	Date on which an admitted patient completes an episode of care.
<i>Context:</i>	Required to identify the period in which an admitted patient hospital stay or episode occurred and for derivation of length of stay.
<i>Object class:</i>	Episode of admitted patient care
<i>Property:</i>	Separation date

Value domain attributes

Representational attributes

<i>Representation class:</i>	Date
<i>Data type:</i>	Date/Time
<i>Format:</i>	DDMMYYYY
<i>Maximum character length:</i>	8

Data element attributes

Collection and usage attributes

<i>Comments:</i>	There may be variations amongst jurisdictions with respect to the recording of separation date. This most often occurs for patients who are statistically separated after a period of leave (and who do not return for further hospital care). In this case, some jurisdictions may record the separation date as the date of statistical separation (and record intervening days as leave days) while other jurisdictions may retrospectively separate patients on the first day of leave. Despite the variations in recording of separation date for this group of patients, the current practices provide for the accurate recording of length of stay.
------------------	---

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Separation date, version 5, DE, NHDD, NHIMG,
-------------------------------------	---

Superseded 01/03/2005

Is used in the formation of Episode of admitted patient care – length of stay (excluding leave days), total N[NN] NHIG, Standard 01/03/2005

Is used in the formation of Establishment – number of separations (financial year), total N[NNNNN] NHIG, Standard 01/03/2005

Is used in the formation of Episode of care – number of psychiatric care days, total N[NNNN] NHIG, Standard 01/03/2005

Is used in the formation of Episode of admitted patient care – major diagnostic category, code (AR-DRG v5.1) NN NHIG, Standard 01/03/2005

Is used in the formation of Episode of admitted patient care – length of stay (including leave days), total N[NN] NHIG, Standard 01/03/2005

Is used in the formation of Episode of admitted patient care – diagnosis related group, code (AR-DRG v5.1) ANNA NHIG, Standard 01/03/2005

Is used in the formation of Episode of admitted patient care (postnatal) – length of stay (including leave days), total N[NN] NHIG, Standard 01/03/2005

Implementation in Data Set Specifications:

Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Admitted patient care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this field must:

- be ≤ last day of financial year
- be ≥ first day of financial year
- be ≥ Admission date

Admitted patient care NMDS 2007-2008 NHIG, Standard 29/11/2006

Implementation start date: 01/07/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this field must:

- be ≤ last day of financial year
- be ≥ first day of financial year
- be ≤ Admission date

Admitted patient mental health care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this field must:

- be \leq last day of financial year
- be \geq first day of financial year
- be \geq Admission date

Admitted patient mental health care NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this field must:

- be \leq last day of financial year
- be \geq first day of financial year
- be \geq Admission date

Admitted patient mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this field must:

- be \leq last day of financial year
- be \geq first day of financial year
- be \geq Admission date

Admitted patient palliative care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this field must:

- be \leq last day of financial year
- be \geq first day of financial year
- be \geq Admission date

Admitted patient palliative care NMDS 2006-2007 NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this field must:

- be \leq last day of financial year
- be \geq first day of financial year

- be >= Admission date

Admitted patient palliative care NMDS 2007-08 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Information specific to this data set:

For the provision of state and territory hospital data to
Commonwealth agencies this field must:

- be <= last day of financial year
- be >= first day of financial year
- be >= Admission date

Perinatal NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

For the provision of state and territory hospital data to
Commonwealth agencies this field must:

- be <= last day of financial year
- be >= first day of financial year
- be >= Admission date

Perinatal NMDS NHIG, Superseded 06/09/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

For the provision of state and territory hospital data to
Commonwealth agencies this field must:

- be <= last day of financial year
- be >= first day of financial year
- be >= Admission date

Perinatal NMDS 2007-2008 NHIG, Standard 06/09/2006

Implementation start date: 01/07/2007

Information specific to this data set:

For the provision of state and territory hospital data to
Commonwealth agencies this field must:

- be <= last day of financial year
- be >= first day of financial year
- be >= Admission date

Separation time

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of admitted patient care – separation time, hhmm
<i>METeOR identifier:</i>	270026
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Time at which an admitted patient completes an episode of care.

Data element concept attributes

<i>Data element concept:</i>	Episode of admitted patient care – separation time
<i>Definition:</i>	Time at which an admitted patient completes an episode of care.
<i>Context:</i>	Admitted patient care.
<i>Object class:</i>	Episode of admitted patient care
<i>Property:</i>	Separation time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Time
<i>Data type:</i>	Date/Time
<i>Format:</i>	hhmm
<i>Maximum character length:</i>	4

Source and reference attributes

<i>Reference documents:</i>	ISO 8601:2000 : Data elements and interchange formats - Information interchange - Representation of dates and times
-----------------------------	---

Data element attributes

Collection and usage attributes

<i>Comments:</i>	Required to identify the time of completion of the episode or hospital stay, for calculation of length of stay.
------------------	---

Relational attributes

<i>Related metadata references:</i>	Supersedes Separation time, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
-------------------------------------	---

Separations

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment – number of separations (financial year), total N[NNNNN]
<i>Synonymous names:</i>	Discharge
<i>METeOR identifier:</i>	270407
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The total number of separations occurring during the reference period. This includes both formal and statistical separations.

Data element concept attributes

<i>Data element concept:</i>	Establishment – number of separations
<i>Definition:</i>	The total number of separations occurring during the reference period. This includes both formal and statistical separations.
<i>Context:</i>	Admitted patient care: Needed as the basic count of the number of separations from care for an establishment.
<i>Object class:</i>	Establishment
<i>Property:</i>	Number of separations

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NNNNN]
<i>Maximum character length:</i>	6
<i>Unit of measure:</i>	Separation

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	May be calculated at: <ul style="list-style-type: none">• individual establishment level; or• system (ie. state/territory) level ie. the sum of the number of establishments. The sum of the number of separations where the separation date has a value: <ul style="list-style-type: none">• >= the beginning of the reference period (typically a financial year); and• <= the end of the reference period.
-----------------------	---

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	<p>Is formed using Episode of admitted patient care – separation date, DDMMYYYY NHIG, Standard 01/03/2005</p> <p>Supersedes Separations, version 2, Derived DE, NHDD, NHIMG, Superseded 01/03/2005</p>
<i>Implementation in Data Set Specifications:</i>	<p>Community mental health establishments NMDS 2004-2005 NHIG, Superseded 08/12/2004</p> <p style="padding-left: 2em;"><i>Implementation start date:</i> 01/07/2004</p> <p style="padding-left: 2em;"><i>Implementation end date:</i> 30/06/2005</p> <p>Mental health establishments NMDS 2005-2006 NHIG, Superseded 07/12/2005</p> <p style="padding-left: 2em;"><i>Implementation start date:</i> 01/07/2005</p> <p>Mental health establishments NMDS 2005-2006 NHIG, Superseded 21/03/2006</p> <p style="padding-left: 2em;"><i>Implementation start date:</i> 01/07/2005</p> <p style="padding-left: 2em;"><i>Implementation end date:</i> 30/06/2006</p> <p style="padding-left: 2em;"><i>Information specific to this data set:</i></p> <p style="padding-left: 4em;">For the Mental health establishments national minimum data set, separations are only to be reported for specialised mental health services provided by private hospitals that receive state or territory government funding. That is, separations from public hospitals need not be reported.</p> <p>Mental health establishments NMDS 2006-2007 NHIG, Superseded 23/10/2006</p> <p style="padding-left: 2em;"><i>Implementation start date:</i> 01/07/2006</p> <p style="padding-left: 2em;"><i>Implementation end date:</i> 30/06/2007</p> <p style="padding-left: 2em;"><i>Information specific to this data set:</i></p> <p style="padding-left: 4em;">For the Mental health establishments national minimum data set, separations are only to be reported for specialised mental health services provided by private hospitals that receive state or territory government funding. That is, separations from public hospitals need not be reported.</p> <p>Mental health establishments NMDS 2007-2008 NHIG, Standard 23/10/2006</p> <p style="padding-left: 2em;"><i>Implementation start date:</i> 01/07/2007</p> <p style="padding-left: 2em;"><i>Information specific to this data set:</i></p> <p style="padding-left: 4em;">For the Mental health establishments national minimum data set, separations are only to be reported for specialised mental health services provided by private hospitals that receive state or territory government funding. That is, separations from public hospitals need not be reported.</p>

Service contact date

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Service contact – service contact date, DDMMYYYY
<i>METeOR identifier:</i>	270122
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The date of service contact between a health service provider and patient/client.

Data element concept attributes

<i>Data element concept:</i>	Service contact – service contact date
<i>Definition:</i>	The date of each service contact between a health service provider and patient/client.
<i>Context:</i>	Community-based mental health care and clinical settings: The service contact is required for clinical audit and other quality assurance purposes.
<i>Object class:</i>	Service contact
<i>Property:</i>	Service contact date

Value domain attributes

Representational attributes

<i>Representation class:</i>	Date
<i>Data type:</i>	Date/Time
<i>Format:</i>	DDMMYYYY
<i>Maximum character length:</i>	8

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Requires services to record the date of each service contact, including the same date where multiple visits are made on one day (except where the visits may be regarded as a continuation of the one service contact). Where an individual patient/client participates in a group activity, a service contact date is recorded if the person's participation in the group activity results in a dated entry being made in the patient's/client's record.
<i>Collection methods:</i>	For collection from community based (ambulatory and non-residential) agencies.

Relational attributes

<i>Related metadata references:</i>	Supersedes Service contact date, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005 Is used in the formation of Person – number of service contact dates, total N[NN] NHIG, Standard 01/03/2005
<i>Implementation in Data Set</i>	Cardiovascular disease (clinical) DSS NHIG, Superseded

Specifications:

15/02/2006

Cardiovascular disease (clinical) DSS NHIG, Standard

15/02/2006

Community mental health care 2004-2005 NHIG, Superseded

08/12/2004

Implementation start date: 01/07/2004

Implementation end date: 30/06/2005

Information specific to this data set:

Collection of the date of each service contact with health service providers allows a description or profile of service utilisation by a person or persons during an episode of care.

Diabetes (clinical) DSS NHIG, Superseded 21/09/2005

Diabetes (clinical) DSS NHIG, Standard 21/09/2005

Service mode (non-admitted patient)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Non-admitted patient service event – service mode, hospital code N{.N}
<i>METeOR identifier:</i>	270096
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Relative physical location of the patient, provider and the hospital campus of the provider of a non-admitted patient service event, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Non-admitted patient service event – service mode
<i>Definition:</i>	Relative physical location of the patient, provider and the hospital campus of the provider of a non-admitted patient service event.
<i>Context:</i>	Hospital non-admitted patient care.
<i>Object class:</i>	Non-admitted patient service event
<i>Property:</i>	Service mode

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code														
<i>Data type:</i>	Number														
<i>Format:</i>	N{.N}														
<i>Maximum character length:</i>	2														
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Patient and provider in the same physical location</td></tr><tr><td>1.1</td><td>On the hospital campus of the provider</td></tr><tr><td>1.2</td><td>Not on the hospital campus of the provider</td></tr><tr><td>2</td><td>Patient and provider not in the same physical location, and communicating via:</td></tr><tr><td>2.1</td><td>Telephone</td></tr><tr><td>2.2</td><td>Telemedicine</td></tr></tbody></table>	Value	Meaning	1	Patient and provider in the same physical location	1.1	On the hospital campus of the provider	1.2	Not on the hospital campus of the provider	2	Patient and provider not in the same physical location, and communicating via:	2.1	Telephone	2.2	Telemedicine
Value	Meaning														
1	Patient and provider in the same physical location														
1.1	On the hospital campus of the provider														
1.2	Not on the hospital campus of the provider														
2	Patient and provider not in the same physical location, and communicating via:														
2.1	Telephone														
2.2	Telemedicine														

Collection and usage attributes

<i>Guide for use:</i>	<p>CODE 1.1 On the hospital campus of the provider Patient and provider in the same physical location refers to face to face contacts. If this occurs at the hospital campus of the provider, use code 1.1.</p> <p>CODE 1.2 Not on the hospital campus of the provider If the service event does not occur on the hospital campus of the provider (hospital-based outreach services), use code 1.2. Hospital-based outreach service events occur when the patient is treated by hospital staff in a location that is not part of the</p>
-----------------------	--

hospital campus (such as in the patient's home or place of work).

Patient and provider not in the same physical location refers to service events delivered via a telephone call or video link (telemedicine). The provider may or may not be physically present on their hospital campus.

A service event delivered via a telephone call is included if

- it is a substitute for a face-to-face service event, and
- it is pre-arranged, and
- a record of the service event is included in the patient's medical record

A service event can be counted at each site participating via a video link.

Data element attributes

Source and reference attributes

Origin: National Health Data Committee

Relational attributes

Related metadata references: Supersedes Non-admitted patient service mode, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Service type (non-admitted patient)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Non-admitted patient service event – service event type (clinical), code N[N]
<i>METeOR identifier:</i>	270090
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The type of clinical service provided to a non-admitted patient in a non-admitted patient service event, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Non-admitted patient service event – service event type (clinical)
<i>Definition:</i>	The type of clinical service provided to a non-admitted patient in a non-admitted patient service event.
<i>Context:</i>	Hospital non-admitted patient care: This definition applies to non-admitted hospital patients and is not intended to apply to community-based services.
<i>Object class:</i>	Non-admitted patient service event
<i>Property:</i>	Service event type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																								
<i>Data type:</i>	Number																								
<i>Format:</i>	N[N]																								
<i>Maximum character length:</i>	2																								
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Allied health and/or clinical nurse specialist</td></tr><tr><td>2</td><td>Dental</td></tr><tr><td>3</td><td>Imaging</td></tr><tr><td>4</td><td>Medical</td></tr><tr><td>5</td><td>Obstetrics and gynaecology</td></tr><tr><td>6</td><td>Paediatrics</td></tr><tr><td>7</td><td>Pathology</td></tr><tr><td>8</td><td>Pharmacy</td></tr><tr><td>9</td><td>Psychiatric</td></tr><tr><td>10</td><td>Surgical</td></tr><tr><td>11</td><td>Emergency department</td></tr></tbody></table>	Value	Meaning	1	Allied health and/or clinical nurse specialist	2	Dental	3	Imaging	4	Medical	5	Obstetrics and gynaecology	6	Paediatrics	7	Pathology	8	Pharmacy	9	Psychiatric	10	Surgical	11	Emergency department
Value	Meaning																								
1	Allied health and/or clinical nurse specialist																								
2	Dental																								
3	Imaging																								
4	Medical																								
5	Obstetrics and gynaecology																								
6	Paediatrics																								
7	Pathology																								
8	Pharmacy																								
9	Psychiatric																								
10	Surgical																								
11	Emergency department																								

Collection and usage attributes

<i>Guide for use:</i>	The following provides a guide to types of clinical services that are included in each of the categories in the data domain.
-----------------------	--

for use should be classified as one of the groups in the data domain on the basis of the type of clinical professional staff involved in providing the service event.

In paediatric hospitals, the full range of service types should be used. That is, paediatric medical should be reported as medical and paediatric surgical should be reported as surgical.

Clinical service type	Clinical service examples
Allied health and/or clinical nurse specialist	Audiology
	Clinical pharmacy
	Diabetes education
	Neuropsychology
	Nutrition/dietetics
	Occupational therapy
	Optometry
	Orthoptics
	Orthotics
	Physiotherapy
	Podiatry
	Prosthetics
	Psychology
	Social work
	Speech pathology
	Stomal therapy
	Wound management
Dental	Dental
Imaging	Medical imaging
Medical	Aged care
	Alcohol and other drug
	Allergy
	Anti-coagulant
	Asthma
	Cardiology
	Clinical measurement
	Dermatology
	Dementia
	Developmental disabilities
	Diabetes
	Endocrine
	Epilepsy
	Falls
	Gastroenterology
	General internal medicine

	Genetic
	Haematology
	Hepatobiliary
	Hypertension
	Hyperbaric medicine
	Immunology
	Infectious diseases
	Medical oncology
	Metabolic bone
	Nephrology
	Neurology
	Occupational medicine
	Palliative care
	Pain management
	Pulmonary
	Radiation oncology
	Rehabilitation
	Respiratory
	Rheumatology
	Spinal
	Transplants
Obstetrics and gynaecology	Family planning
	Gynaecology
	Gynaecology oncology
	Obstetrics
	Assisted reproductive technology
Pathology	Pathology
Paediatrics	Adolescent health
	Neonatal
	Paediatric medicine
	Paediatric surgery
Pharmacy	Dispensing pharmacy
Psychiatric	Psychiatry
Surgical	Breast
	Burns
	Cardiac surgery
	Colorectal
	Craniofacial
	Ear, nose and throat
	Fracture
	General surgery
	Neurosurgery

	Ophthalmology
	Orthopaedics
	Plastic surgery
	Pre-admission
	Pre-anaesthesia
	Thoracic surgery
	Urology
	Vascular surgery
Emergency department	Emergency department

An emergency department provides triage, assessment, care and/or treatment for patients suffering from medical condition/s and/or injury.

Data element attributes

Relational attributes

Related metadata references:

Supersedes Non-admitted patient service type, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Sex

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – sex, code N
<i>METeOR identifier:</i>	287316
<i>Registration status:</i>	NHIG, Standard 04/05/2005 NCSIMG, Standard 25/08/2005 NHDAMG, Standard 10/02/2006
<i>Definition:</i>	The biological distinction between male and female, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – sex
<i>Definition:</i>	Sex is the biological distinction between male and female. Where there is an inconsistency between anatomical and chromosomal characteristics, sex is based on anatomical characteristics.
<i>Context:</i>	Sex is a core metadata item in a wide range of social, labour and demographic statistics.
<i>Object class:</i>	Person
<i>Property:</i>	Sex

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code										
<i>Data type:</i>	Number										
<i>Format:</i>	N										
<i>Maximum character length:</i>	1										
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Male</td></tr><tr><td>2</td><td>Female</td></tr><tr><td>3</td><td>Intersex or indeterminate</td></tr><tr><td>9</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	1	Male	2	Female	3	Intersex or indeterminate	9	Not stated/inadequately described
Value	Meaning										
1	Male										
2	Female										
3	Intersex or indeterminate										
9	Not stated/inadequately described										
<i>Supplementary values:</i>											

Collection and usage attributes

<i>Guide for use:</i>	Diagnosis and procedure codes should be checked against the national ICD-10-AM sex edits, unless the person is undergoing, or has undergone a sex change or has a genetic condition resulting in a conflict between sex and ICD-10-AM code. CODE 3 Intersex or indeterminate Intersex or indeterminate, refers to a person, who because of a genetic condition, was born with reproductive organs or sex chromosomes that are not exclusively male or female or whose sex has not yet been determined for whatever reason. Intersex or indeterminate, should be confirmed if reported for people aged 90 days or greater.
-----------------------	--

Comments: The definition for Intersex in Guide for use is sourced from the ACT Legislation (Gay, Lesbian and Transgender) Amendment Act 2003.

Source and reference attributes

Origin: Australian Capital Territory 2003. Legislation (Gay, Lesbian and Transgender) Amendment Act 2003

Reference documents: Legislation (Gay, Lesbian and Transgender) Amendment Act 2003. See <http://www.legislation.act.gov.au/a/2003-14/20030328-4969/pdf/2003-14.pdf>.

Data element attributes

Collection and usage attributes

Collection methods: Operationally, sex is the distinction between male and female, as reported by a person or as determined by an interviewer. When collecting data on sex by personal interview, asking the sex of the respondent is usually unnecessary and may be inappropriate, or even offensive. It is usually a simple matter to infer the sex of the respondent through observation, or from other cues such as the relationship of the person(s) accompanying the respondent, or first name. The interviewer may ask whether persons not present at the interview are male or female.

A person's sex may change during their lifetime as a result of procedures known alternatively as sex change, gender reassignment, transsexual surgery, transgender reassignment or sexual reassignment. Throughout this process, which may be over a considerable period of time, the person's sex could be recorded as either Male or Female.

In data collections that use the ICD-10-AM classification, where sex change is the reason for admission, diagnoses should include the appropriate ICD-10-AM code(s) that clearly identify that the person is undergoing such a process. This code(s) would also be applicable after the person has completed such a process, if they have a procedure involving an organ(s) specific to their previous sex (e.g. where the patient has prostate or ovarian cancer).

CODE 3 Intersex or indeterminate

Is normally used for babies for whom sex has not been determined for whatever reason.

Should not generally be used on data collection forms completed by the respondent.

Should only be used if the person or respondent volunteers that the person is intersex or where it otherwise becomes clear during the collection process that the individual is neither male nor female.

CODE 9 Not stated/inadequately described

Is not to be used on primary collection forms. It is primarily for use in administrative collections when transferring data from data sets where the item has not been collected.

Source and reference attributes

Origin: Australian Institute of Health and Welfare (AIHW) National

Mortality Database 1997/98 AIHW 2001 National Diabetes Register, Statistical Profile, December 2000 (Diabetes Series No. 2.)

Reference documents:

Australian Bureau of Statistics

AS4846 Health Care Provider Identification, 2004, Sydney: Standards Australia

AS5017 Health Care Client Identification, 2002, Sydney: Standards Australia

In AS4846 and AS5017 alternative codes are presented. Refer to the current standard for more details.

Relational attributes

Related metadata references:

Supersedes Person—sex, code N NHIG, Superseded 04/05/2005, NCSIMG, Superseded 31/08/2005

Is used in the formation of Episode of admitted patient care—major diagnostic category, code (AR-DRG v5.1) NN NHIG, Standard 01/03/2005

Is used in the formation of Episode of admitted patient care—diagnosis related group, code (AR-DRG v5.1) ANNA NHIG, Standard 01/03/2005

Implementation in Data Set Specifications:

Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Admitted patient care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient care NMDS 2007-2008 NHIG, Standard 29/11/2006

Implementation start date: 01/07/2007

Admitted patient mental health care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient mental health care NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Admitted patient mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Admitted patient palliative care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient palliative care NMDS 2006-2007 NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Admitted patient palliative care NMDS 2007-08 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Alcohol and other drug treatment services NMDS NHIG,
Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Alcohol and other drug treatment services NMDS NHIG,
Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Alcohol and other drug treatment services NMDS 2007-2008
NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Cancer (clinical) DSS NHIG, Standard 07/12/2005

Cancer (clinical) DSS NHIG, Superseded 07/12/2005

Cardiovascular disease (clinical) DSS NHIG, Superseded
15/02/2006

Cardiovascular disease (clinical) DSS NHIG, Standard
15/02/2006

Community mental health care 2004-2005 NHIG, Superseded
08/12/2004

Implementation start date: 01/07/2004

Implementation end date: 30/06/2005

Community mental health care NMDS 2005-2006 NHIG,
Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Community mental health care NMDS 2006-2007 NHIG,
Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Community mental health care NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Computer Assisted Telephone Interview demographic module
DSS NHIG, Standard 04/05/2005

Information specific to this data set:

For data collections using the Computer Assisted
Telephone Interview demographic module DSS, only
Codes 1 and 2 are collected.

For data collection using Computer Assisted Telephone
Interviewing (CATI) it is recommended that sex be
ascertained by the interviewer on the basis of information
already collected or other cues such as voice or name. If
the interviewer is unsure of the respondent's sex the
following question may be used:

As this interview is being conducted over the phone, I need to ask are you male or female?

Male

Female

(Interviewer note: If respondent provides any response that can't be categorised to male or female, terminate interview.)

Diabetes (clinical) DSS NHIG, Superseded 21/09/2005

Diabetes (clinical) DSS NHIG, Standard 21/09/2005

Health care client identification DSS NHIG, Standard 04/05/2005

NCSIMG, Standard 03/10/2006

Health care provider identification DSS NHIG, Standard 04/05/2005

Non-admitted patient emergency department care NMDS NHIG, Standard 24/03/2006

Implementation start date: 01/07/2006

Non-admitted patient emergency department care NMDS NHIG, Superseded 07/12/2005

Non-admitted patient emergency department care NMDS NHIG, Superseded 24/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Perinatal NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Perinatal NMDS NHIG, Superseded 06/09/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Perinatal NMDS 2007-2008 NHIG, Standard 06/09/2006

Implementation start date: 01/07/2007

Residential mental health care NMDS 2005-2006 NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Residential mental health care NMDS 2006-2007 NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Residential mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Source of referral to alcohol and other drug treatment service

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of treatment for alcohol and other drugs – referral source, code NN
<i>METeOR identifier:</i>	269946
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The source from which the person was transferred or referred to the alcohol and other drug treatment service, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Episode of treatment for alcohol and other drugs – referral source
<i>Definition:</i>	The source from which the person was transferred or referred to the alcohol and other drug treatment service.
<i>Context:</i>	Alcohol and other drug treatment services.
<i>Object class:</i>	Episode of treatment for alcohol and other drugs
<i>Property:</i>	Referral source

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																								
<i>Data type:</i>	String																								
<i>Format:</i>	NN																								
<i>Maximum character length:</i>	2																								
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>Self</td></tr><tr><td>02</td><td>Family member/friend</td></tr><tr><td>03</td><td>Medical practitioner</td></tr><tr><td>04</td><td>Hospital</td></tr><tr><td>05</td><td>Mental health care service</td></tr><tr><td>06</td><td>Alcohol and other drug treatment service</td></tr><tr><td>07</td><td>Other community/health care service</td></tr><tr><td>08</td><td>Correctional service</td></tr><tr><td>09</td><td>Police diversion</td></tr><tr><td>10</td><td>Court diversion</td></tr><tr><td>98</td><td>Other</td></tr></tbody></table>	Value	Meaning	01	Self	02	Family member/friend	03	Medical practitioner	04	Hospital	05	Mental health care service	06	Alcohol and other drug treatment service	07	Other community/health care service	08	Correctional service	09	Police diversion	10	Court diversion	98	Other
Value	Meaning																								
01	Self																								
02	Family member/friend																								
03	Medical practitioner																								
04	Hospital																								
05	Mental health care service																								
06	Alcohol and other drug treatment service																								
07	Other community/health care service																								
08	Correctional service																								
09	Police diversion																								
10	Court diversion																								
98	Other																								
<i>Supplementary values:</i>	99 Not stated/inadequately described																								

Collection and usage attributes

<i>Guide for use:</i>	CODE 03 Medical practitioner Includes medical specialists, vocationally registered general
-----------------------	---

practitioners, vocationally registered general practitioner trainees and other primary-care medical practitioners in private practice.

CODE 04 Hospital

Includes public and private hospitals, hospitals specialising in dental, ophthalmic aids and other specialised medical or surgical care, satellite units managed and staffed by a hospital, emergency departments of hospitals, and mothercraft hospitals. Excludes psychiatric hospitals, psychiatric units and drug and alcohol units located within or operating from hospitals, and outpatient clinics (see codes 05-07).

CODE 05 Mental health care service

Includes both residential and non-residential services. Includes psychiatric hospitals and psychiatric units within and outside of hospitals.

CODE 06 Alcohol and other drug treatment service

Includes both residential and non-residential services. Includes drug and alcohol units within and outside of hospitals.

CODE 07 Other community/health care service

Includes outpatient clinics and aged care facilities.

CODE 09 Police diversion

This code should be used when a person detained for a minor drug offence is formally referred to treatment by the police in order to divert the offender from the criminal justice pathway.

CODE 10 Court diversion

This code refers to the diversion of an offender into drug education, assessment and treatment at the discretion of a magistrate. This may occur at the point of bail or prior to sentencing.

CODE 98 Other

Includes persons referred under a legislative act (other than *Drug Diversion Act*) e.g. *Mental Health Act*.

Data element attributes

Collection and usage attributes

Comments: Source of referral is important in assisting in the analyses of inter-sectoral patient/client flow and for health care planning.

Source and reference attributes

Submitting organisation: Intergovernmental Committee on Drugs National Minimum Data Set Working Group

Relational attributes

Related metadata references: Supersedes Source of referral to alcohol and other drug treatment service, version 3, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Alcohol and other drug treatment services NMDS NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Alcohol and other drug treatment services NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Alcohol and other drug treatment services NMDS 2007-2008

NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Source of referral to public psychiatric hospital

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of admitted patient care – referral source, public psychiatric hospital code NN
<i>METeOR identifier:</i>	269947
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Source from which the person was transferred/referred to the public psychiatric hospital, as represented by a code.
<i>Context:</i>	To assist in analyses of intersectoral patient flow and health care planning.

Data element concept attributes

<i>Data element concept:</i>	Episode of admitted patient care – referral source
<i>Definition:</i>	The source from which a patient is referred for an episode of admitted patient care.
<i>Object class:</i>	Episode of admitted patient care
<i>Property:</i>	Referral source

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																						
<i>Data type:</i>	String																						
<i>Format:</i>	NN																						
<i>Maximum character length:</i>	2																						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>Private psychiatric practice</td></tr><tr><td>02</td><td>Other private medical practice</td></tr><tr><td>03</td><td>Other public psychiatric hospital</td></tr><tr><td>04</td><td>Other health care establishment</td></tr><tr><td>05</td><td>Other private hospital</td></tr><tr><td>06</td><td>Law enforcement agency</td></tr><tr><td>07</td><td>Other agency</td></tr><tr><td>08</td><td>Outpatient department</td></tr><tr><td>09</td><td>Other</td></tr><tr><td>10</td><td>Unknown</td></tr></tbody></table>	Value	Meaning	01	Private psychiatric practice	02	Other private medical practice	03	Other public psychiatric hospital	04	Other health care establishment	05	Other private hospital	06	Law enforcement agency	07	Other agency	08	Outpatient department	09	Other	10	Unknown
Value	Meaning																						
01	Private psychiatric practice																						
02	Other private medical practice																						
03	Other public psychiatric hospital																						
04	Other health care establishment																						
05	Other private hospital																						
06	Law enforcement agency																						
07	Other agency																						
08	Outpatient department																						
09	Other																						
10	Unknown																						
<i>Supplementary values:</i>																							

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Source of referral to public psychiatric hospital,
-------------------------------------	---

Implementation in Data Set Specifications:

version 3, DE, NHDD, NHIMG, Superseded 01/03/2005

Admitted patient care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient care NMDS 2006-2007 NHIG, Superseded 29/11/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Admitted patient care NMDS 2007-2008 NHIG, Standard 29/11/2006

Implementation start date: 01/07/2007

Admitted patient mental health care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient mental health care NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Admitted patient mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised mental health service program type

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Specialised mental health service – admitted patient care program type, code N
<i>METeOR identifier:</i>	288889
<i>Registration status:</i>	NHIG, Standard 08/12/2004
<i>Definition:</i>	Type of admitted patient care program provided by a specialised mental health service, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Specialised mental health service – admitted patient care program type
<i>Definition:</i>	Type of admitted patient care program provided by a specialised mental health service.
<i>Context:</i>	Specialised mental health services.
<i>Object class:</i>	Specialised mental health service
<i>Property:</i>	Admitted patient care program type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Acute care</td></tr><tr><td>2</td><td>Other</td></tr></tbody></table>	Value	Meaning	1	Acute care	2	Other
Value	Meaning						
1	Acute care						
2	Other						

Collection and usage attributes

<i>Guide for use:</i>	<p>The categorisation of the admitted patient program is based on the principal purpose(s) of the program rather than the classification of the individual patients.</p> <p>CODE 1 Acute care</p> <p>Programs primarily providing specialist psychiatric care for people with acute episodes of mental disorder. These episodes are characterised by recent onset of severe clinical symptoms of mental disorder, that have potential for prolonged dysfunction or risk to self and/or others. The key characteristic of acute services is that this treatment effort is focused on short-term treatment. Acute services may be focused on assisting people who have had no prior contact or previous psychiatric history, or individuals with a continuing mental disorder for whom there has been an acute exacerbation of symptoms. This category applies only to services with a mental health service setting of overnight admitted patient care or residential care.</p> <p>CODE 2 Other</p>
-----------------------	--

Refers to all other programs primarily providing admitted patient care.

Includes programs providing rehabilitation services that have a primary focus on intervention to reduce functional impairments that limit the independence of patients. Rehabilitation services are focused on disability and the promotion of personal recovery.

They are characterised by an expectation of substantial improvement over the short to mid-term. Patients treated by rehabilitation services usually have a relatively stable pattern of clinical symptoms.

Also includes programs providing extended care services that primarily provide care over an indefinite period for patients who have a stable but severe level of functional impairment and an inability to function independently, thus requiring extensive care and support. Patients of extended care services present a stable pattern of clinical symptoms, which may include high levels of severe unremitting symptoms of mental disorder. Treatment is focused on preventing deterioration and reducing impairment; improvement is expected to occur slowly.

Data element attributes

Collection and usage attributes

Guide for use:

This data element is used to disaggregate data on beds, activity, expenditure and staffing for admitted patient settings in mental health service units (see Specialised mental health service – service setting, code N data element).

Relational attributes

Implementation in Data Set Specifications:

Mental health establishments NMDS 2005-2006 NHIG,
Superseded 07/12/2005

Implementation start date: 01/07/2005

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Mental health establishments NMDS 2005-2006 NHIG,
Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Mental health establishments NMDS 2006-2007 NHIG,
Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Mental health establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Specialised mental health service setting

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Specialised mental health service—service setting, code N
<i>METeOR identifier:</i>	288899
<i>Registration status:</i>	NHIG, Standard 08/12/2004
<i>Definition:</i>	The setting for care provided by a specialised mental health service, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Specialised mental health service—service setting
<i>Definition:</i>	The setting for care provided by a specialised mental health service.
<i>Context:</i>	Specialised mental health services.
<i>Object class:</i>	Specialised mental health service
<i>Property:</i>	Service delivery setting

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code								
<i>Data type:</i>	Number								
<i>Format:</i>	N								
<i>Maximum character length:</i>	1								
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Admitted patient care setting</td></tr><tr><td>2</td><td>Residential care setting</td></tr><tr><td>3</td><td>Ambulatory care setting</td></tr></tbody></table>	Value	Meaning	1	Admitted patient care setting	2	Residential care setting	3	Ambulatory care setting
Value	Meaning								
1	Admitted patient care setting								
2	Residential care setting								
3	Ambulatory care setting								
<i>Supplementary values:</i>	9 Unknown/not stated/inadequately described								

Collection and usage attributes

<i>Guide for use:</i>	<p>CODE 1 Admitted patient care setting The component of specialised mental health services that provides admitted patient care. These are specialised psychiatric hospitals and specialist psychiatric units located within hospitals that are not specialised psychiatric hospitals. Excludes hospital outpatient clinics.</p> <p>CODE 2 Residential care setting The component of specialised mental health services that provides residential care within residential mental health services. Excludes components that provide ambulatory care to patients or clients who are not residents.</p> <p>CODE 3 Ambulatory care setting The component of specialised mental health services that provides ambulatory care (service contacts). They include hospital outpatient clinics and non-hospital community mental health services, such as crisis or mobile assessment and</p>
-----------------------	---

treatment services, day programs, outreach services and consultation/liaison services.

Data element attributes

Collection and usage attributes

Guide for use:

A single mental health service unit may provide care in more than one setting. This data element is intended to allow staffing, resource and expenditure data related to these settings to be identified and reported separately.

Relational attributes

Related metadata references:

Supersedes Specialised mental health service – service delivery setting, code N NHIG, Superseded 08/12/2004

Implementation in Data Set Specifications:

Community mental health establishments NMDS 2004-2005 NHIG, Superseded 08/12/2004

Implementation start date: 01/07/2004

Implementation end date: 30/06/2005

Information specific to this data set:

Only permissible values 2, 3 and 9 are to be used for the Community Mental Health Establishments National Minimum Data Set (NMDS). Domain value 1 is not applicable.

Mental health establishments NMDS 2005-2006 NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Mental health establishments NMDS 2005-2006 NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Mental health establishments NMDS 2006-2007 NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Mental health establishments NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised mental health service target population

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Specialised mental health service – target population group, code N
<i>METeOR identifier:</i>	288957
<i>Registration status:</i>	NHIG, Standard 08/12/2004
<i>Definition:</i>	The population group primarily targeted by a specialised mental health service, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Specialised mental health service – specialised mental health service target population group
<i>Definition:</i>	The population group primarily targeted by a specialised mental health service.
<i>Context:</i>	Specialised mental health services.
<i>Object class:</i>	Specialised mental health service
<i>Property:</i>	Specialised mental health service target population group

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code										
<i>Data type:</i>	String										
<i>Format:</i>	N										
<i>Maximum character length:</i>	1										
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Child and adolescent</td></tr><tr><td>2</td><td>Older person</td></tr><tr><td>3</td><td>Forensic</td></tr><tr><td>4</td><td>General</td></tr></tbody></table>	Value	Meaning	1	Child and adolescent	2	Older person	3	Forensic	4	General
Value	Meaning										
1	Child and adolescent										
2	Older person										
3	Forensic										
4	General										

Collection and usage attributes

<i>Guide for use:</i>	<p>CODE 1 Child and adolescent</p> <p>These services principally target children and young people under the age of 18 years. The classification of a service into this category requires recognition by the regional or central funding authority of the special focus of the service. These services may include a forensic component.</p> <p>CODE 2 Older person</p> <p>These services principally target people in the age group of 65 years and over. The classification of a service into this category requires recognition by the regional or central funding authority of the special focus of the service. These services may include a forensic component.</p> <p>CODE 3 Forensic</p> <p>Health services that provide services primarily for people</p>
-----------------------	---

whose health condition has led them to commit, or be suspected of, a criminal offence or make it likely that they will reoffend without adequate treatment or containment. This includes prison-based services, but excludes services that are primarily for children and adolescents and for older people even where they include a forensic component.

CODE 4 General

These services principally target the general adult population (aged 18–64 years) but may also provide services to children, adolescents or older people. These services are those services that cannot be described as specialist child and adolescent services or services for older people. It excludes forensic services.

Data element attributes

Collection and usage attributes

Guide for use:

This data element is used to disaggregate data on beds, activity, expenditure and staffing for admitted patient settings in mental health service units (see service setting data element).

The order of priority for coding is:

- where the forensic services are for children/ adolescents or older persons these services should be coded to the category for that age group; and
- where the forensic services are for adults these services should be coded to forensic.

Source and reference attributes

Submitting organisation:

Australian Institute of Health and Welfare

Relational attributes

Implementation in Data Set Specifications:

Mental health establishments NMDS 2005-2006 NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Mental health establishments NMDS 2005-2006 NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Mental health establishments NMDS 2006-2007 NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Mental health establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Information specific to this data set:

Obligation condition: reporting of this data element is optional for non-government residential mental health services and specialised mental health services provided by private hospitals that receive state or territory government funding.

Specialised mental health service—hours staffed

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Specialised mental health service—number of hours staffed, average hours NN
<i>METeOR identifier:</i>	288877
<i>Registration status:</i>	NHIG, Standard 08/12/2004
<i>Definition:</i>	The average number of hours per day during which a residential mental health service has appropriately trained staff employed on-site. Training may include formal qualifications and/or on the job training.

Data element concept attributes

<i>Data element concept:</i>	Specialised mental health service—number of hours staffed
<i>Definition:</i>	The average number of hours per day during which a residential mental health service has appropriately trained staff employed on-site. Training may include formal qualifications and/or on the job training.
<i>Context:</i>	Specialised mental health services.
<i>Object class:</i>	Specialised mental health service
<i>Property:</i>	Number of hours staffed

Value domain attributes

Representational attributes

<i>Representation class:</i>	Average				
<i>Data type:</i>	Number				
<i>Format:</i>	NN				
<i>Maximum character length:</i>	2				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99</td><td>Unknown/not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	99	Unknown/not stated/inadequately described
Value	Meaning				
99	Unknown/not stated/inadequately described				
<i>Unit of measure:</i>	Hour (h)				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Whole numbers of hours staffed (no decimals or fractions). Valid numbers are 1 to 24. The hours staffed provides a measure of service intensity for the reporting and analysis of staff, financial and activity data. For residential mental health services, this refers to the number of hours per day during which appropriately trained staff (either with formal qualifications and/or on the job training) are employed on site, as their normal place of employment, within the service unit. It excludes periods where the service unit is only staffed by a resident sleepover staff member or any period where staff are present but not employed on site at the
-----------------------	--

service unit.

Excludes ambulatory and admitted patient services.

Round to nearest whole hour.

Where the number of hours staffed varies by day, average the number of hours staffed over a week, including the weekend.

Relational attributes

Implementation in Data Set Specifications:

Mental health establishments NMDS 2005-2006 NHIG,
Superseded 07/12/2005

Implementation start date: 01/07/2005

Mental health establishments NMDS 2005-2006 NHIG,
Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Mental health establishments NMDS 2006-2007 NHIG,
Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Mental health establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised mental health service—supported public housing places

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Specialised mental health service – number of supported public housing places, total N[N(5)]
<i>METeOR identifier:</i>	288945
<i>Registration status:</i>	NHIG, Standard 08/12/2004
<i>Definition:</i>	The total number of public housing places supported by specialised mental health services available at 30 June, targeted to people affected by mental illness or psychiatric disability. These are places provided by the public housing authority under a formal partnership agreement with the relevant State or Territory health authority. Such agreements commit the State or Territory health authority to assist people within their homes by providing ongoing clinical and disability support, including outreach services.

Data element concept attributes

<i>Data element concept:</i>	Specialised mental health service – number of supported public housing places
<i>Definition:</i>	The total number of public housing places supported by specialised mental health services available at 30 June, targeted to people affected by mental illness or psychiatric disability. These are places provided by the public housing authority under a formal partnership agreement with the relevant State or Territory health authority. Such agreements commit the State or Territory health authority to assist people within their homes by providing ongoing clinical and disability support, including outreach services.
<i>Context:</i>	Specialised mental health services.
<i>Object class:</i>	Specialised mental health service
<i>Property:</i>	Supported public housing places

Collection and usage attributes

<i>Guide for use:</i>	Excludes residential mental health services defined as specialised mental health services.
-----------------------	--

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[N(5)]
<i>Maximum character length:</i>	6

Source and reference attributes

<i>Steward:</i>	Australian Institute of Health and Welfare
-----------------	--

Data element attributes

Relational attributes

Implementation in Data Set Specifications:

Mental health establishments NMDS 2005-2006 NHIG,
Superseded 07/12/2005

Implementation start date: 01/07/2005

Mental health establishments NMDS 2005-2006 NHIG,
Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Mental health establishments NMDS 2006-2007 NHIG,
Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Mental health establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (transplantation unit), yes/no code N
<i>METeOR identifier:</i>	270443
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to organ retrieval, transplantation and ongoing care of the transplant recipient, is provided within an establishment. <ul style="list-style-type: none">• bone marrow• renal• heart, including heart-lung• liver• pancreas.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
-------------------------------------	--

Specialised service indicators—acquired immune deficiency syndrome unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (acquired immune deficiency syndrome unit), yes/no code N
<i>METeOR identifier:</i>	270448
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the treatment of Acquired Immune Deficiency Syndrome (AIDS) patients is provided within an establishment as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—acute renal dialysis unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (acute renal dialysis unit), yes/no code N
<i>METeOR identifier:</i>	270435
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to dialysis of renal failure patients requiring acute care is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—acute spinal cord injury unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (acute spinal cord injury unit), yes/no code N
<i>METeOR identifier:</i>	270432
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the initial treatment and subsequent ongoing management and rehabilitation of patients with acute spinal cord injury, largely conforming to Australian Health Minister's Advisory Council guidelines for service provision, is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set</i>	Public hospital establishments NMDS NHIG, Superseded

Specifications:

21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—alcohol and drug unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (alcohol and drug unit), yes/no code N
<i>METeOR identifier:</i>	270431
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a facility/service dedicated to the treatment of alcohol and drug dependence is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—bone marrow transplantation unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (bone marrow transplantation unit), yes/no code N
<i>METeOR identifier:</i>	308862
<i>Registration status:</i>	NHIG, Standard 07/09/2005
<i>Definition:</i>	Whether or not a specialised facility for bone marrow transplantation is provided within the establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—burns unit (level III)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (burns unit (level III)), yes/no code N
<i>METeOR identifier:</i>	270438
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the initial treatment and subsequent rehabilitation of the severely injured burns patient (usually >10 per cent of patients body surface affected) is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—cardiac surgery unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (cardiac surgery unit), yes/no code N
<i>METeOR identifier:</i>	270434
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to operative and peri-operative care of patients with cardiac disease is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—clinical genetics unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (clinical genetics unit), yes/no code N
<i>METeOR identifier:</i>	270444
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to diagnostic and counselling services for clients who are affected by, at risk of, or anxious about genetic disorders, is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—comprehensive epilepsy centre

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (comprehensive epilepsy centre), yes/no code N
<i>METeOR identifier:</i>	270442
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to seizure characterisation, evaluation of therapeutic regimes, pre-surgical evaluation and epilepsy surgery for patients with refractory epilepsy, is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—coronary care unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (coronary care unit), yes/no code N
<i>METeOR identifier:</i>	270433
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to acute care services for patients with cardiac diseases is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—diabetes unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (diabetes unit), yes/no code N
<i>METeOR identifier:</i>	270449
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the treatment of diabetics is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—domiciliary care service

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (domiciliary care service), yes/no code N
<i>METeOR identifier:</i>	270430
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a facility/service dedicated to the provision of nursing or other professional paramedical care or treatment and non-qualified domestic assistance to patients in their own homes or in residential institutions not part of the establishment is provided by the establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—geriatric assessment unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (geriatric assessment unit), yes/no code N
<i>METeOR identifier:</i>	270429
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not facilities dedicated to the Commonwealth-approved assessment of the level of dependency of (usually) aged individuals either for purposes of initial admission to a long-stay institution or for purposes of reassessment of dependency levels of existing long-stay institution residents, is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—heart, lung transplantation unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (heart, lung transplantation unit), yes/no code N
<i>METeOR identifier:</i>	308866
<i>Registration status:</i>	NHIG, Standard 07/09/2005
<i>Definition:</i>	Whether or not a specialised facility for heart including heart lung transplantation is provided within the establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—hospice care unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (hospice care unit), yes/no code N
<i>METeOR identifier:</i>	270427
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a facility dedicated to the provision of palliative care to terminally ill patients is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—in-vitro fertilisation unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (in-vitro fertilisation unit), yes/no code N
<i>METeOR identifier:</i>	270441
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the investigation of infertility provision of in-vitro fertilisation services is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—**infectious diseases unit**

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (infectious diseases unit), yes/no code N
<i>METeOR identifier:</i>	270447
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the treatment of infectious diseases is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—intensive care unit (level III)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (intensive care unit (level III)), yes/no code N
<i>METeOR identifier:</i>	270426
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the care of paediatric and adult patients requiring intensive care and sophisticated technological support services is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—liver transplantation unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (liver transplantation unit), yes/no code N
<i>METeOR identifier:</i>	308868
<i>Registration status:</i>	NHIG, Standard 07/09/2005
<i>Definition:</i>	Whether or not a specialised facility for liver transplantation is provided within the establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—maintenance renal dialysis centre

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (maintenance renal dialysis centre), yes/no code N
<i>METeOR identifier:</i>	270437
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	An indicator of a specialised facility dedicated to maintenance dialysis of renal failure patients, as represented by a code. It may be a separate facility (possibly located on hospital grounds) or known as a satellite centre or a hospital-based facility but is not a facility solely providing training services.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—major plastic/reconstructive surgery unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (major plastic/reconstructive surgery unit), yes/no code N
<i>METeOR identifier:</i>	270439
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to general purpose plastic and specialised reconstructive surgery, including maxillofacial, microsurgery and hand surgery, is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—neonatal intensive care unit (level III)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (neonatal intensive care unit (level III)), yes/no code N
<i>METeOR identifier:</i>	270436
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the care of neonates requiring care and sophisticated technological support, is provided within an establishment, as represented by a code. Patients usually require intensive cardiorespiratory monitoring, sustained assistance ventilation, long-term oxygen administration and parenteral nutrition.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set</i>	Public hospital establishments NMDS NHIG, Superseded

Specifications:

21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—neuro surgical unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (neuro surgical unit), yes/no code N
<i>METeOR identifier:</i>	270446
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the surgical treatment of neurological conditions is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—nursing home care unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (nursing home care unit), yes/no code N
<i>METeOR identifier:</i>	270428
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a facility dedicated to the provision of nursing home care is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—obstetric/maternity

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (obstetric/maternity), yes/no code N
<i>METeOR identifier:</i>	270150
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the care of obstetric/maternity patients is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—oncology unit, cancer treatment

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (oncology unit) (cancer treatment), yes/no code N
<i>METeOR identifier:</i>	270440
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to multidisciplinary investigation, management, rehabilitation and support services for cancer patients, is provided within an establishment, as represented by a code. Treatment services include surgery, chemotherapy and radiation.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—pancreas transplantation unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (pancreas transplantation unit), yes/no code N
<i>METeOR identifier:</i>	308870
<i>Registration status:</i>	NHIG, Standard 07/09/2005
<i>Definition:</i>	Whether or not a specialised facility for pancreas transplantation is provided within the establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—psychiatric unit/ward

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (psychiatric unit/ward), yes/no code N
<i>METeOR identifier:</i>	270425
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised unit/ward dedicated to the treatment and care of admitted patients with psychiatric, mental, or behavioural disorders, is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—rehabilitation unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (rehabilitation unit), yes/no code N
<i>METeOR identifier:</i>	270450
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not dedicated units within recognised hospitals which provide post-acute rehabilitation and are designed as such by the State health authorities (see metadata item Type of episode of care) are provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—renal transplantation unit

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (renal transplantation unit), yes/no code N
<i>METeOR identifier:</i>	308864
<i>Registration status:</i>	NHIG, Standard 07/09/2005
<i>Definition:</i>	Whether or not a specialised facility for renal transplantation is provided within the establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—sleep centre

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (sleep centre), yes/no code N
<i>METeOR identifier:</i>	270445
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility linked to a sleep laboratory dedicated to the investigation and management of sleep disorders is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006

Public hospital establishments NMDS NHIG, Superseded
23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialised service indicators—specialist paediatric

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment—specialised service indicator (specialist paediatric), yes/no code N
<i>METeOR identifier:</i>	270424
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether or not a specialised facility dedicated to the care of children aged 14 or less is provided within an establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment—specialised service indicator
<i>Definition:</i>	Specialised services provided in establishments.
<i>Context:</i>	Health services: Essential to provide a broad picture of the availability of key specialised services by State and region and to assist with planning if services are over supplied in one region relative to another.
<i>Object class:</i>	Establishment
<i>Property:</i>	Specialised service indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Source and reference attributes

<i>Origin:</i>	National Health Data Committee
----------------	--------------------------------

Relational attributes

<i>Related metadata references:</i>	Supersedes Specialised service indicators, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Public hospital establishments NMDS NHIG, Superseded 21/03/2006 <i>Implementation start date:</i> 01/07/2005 <i>Implementation end date:</i> 30/06/2006 Public hospital establishments NMDS NHIG, Superseded

23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG,
Standard 23/10/2006

Implementation start date: 01/07/2007

Specialist private sector rehabilitation care indicator

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of care – specialist private sector rehabilitation care indicator, code N
<i>METeOR identifier:</i>	270397
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether the rehabilitation care that a patient receives from a private hospital meets the criteria for 'Specialist private sector rehabilitation care', as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Episode of care – specialist private sector rehabilitation care indicator
<i>Definition:</i>	An indicator of whether the rehabilitation care that a patient receives from a private hospital meets the criteria for 'Specialist private sector rehabilitation care' (as determined by the Commonwealth Department of Health and Ageing).
<i>Context:</i>	Admitted and non-admitted patients receiving rehabilitation care from a private hospital.
<i>Object class:</i>	Episode of care
<i>Property:</i>	Specialist private sector rehabilitation care indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No
Value	Meaning						
1	Yes						
2	No						

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>This metadata item is a qualifier of the three 'Rehabilitation' care types for admitted patients in private hospitals. When an admitted patient in a private hospital is receiving rehabilitation care (as defined in Hospital service – care type, code N[N].N), this metadata item should be recorded to denote whether or not that care meets the criteria for 'specialist rehabilitation'.</p> <p>These are the criteria determined by The Commonwealth Department of Health and Ageing in respect of patients treated in the private sector, specialist rehabilitation is:</p> <ul style="list-style-type: none">• Provided by a specialist rehabilitation unit (a separate physical space and a specialist rehabilitation team providing admitted
-----------------------	--

- patient and/or ambulatory care) meeting guidelines issued by the Commonwealth Department of Health and Ageing, and
- provided by a multi-disciplinary team which is under the clinical management of a consultant in rehabilitation medicine or equivalent, and
 - provided for a person with limited functioning (impairments, activity limitations and participation restrictions) and for whom there is a reasonable expectation of functional gain, and
 - for whom the primary treatment goal is improvement in functioning status which is evidenced in the medical record by:
 - an individualised and documented initial and periodic assessment of functional ability, or
 - an individualised multi-disciplinary rehabilitation plan which includes agreed rehabilitation goals and indicative timeframes.

Comments:

This metadata item has been developed by the Private Rehabilitation Working Group, and agreed by the private rehabilitation hospital sector, the private health insurance sector and the Commonwealth Department of Health and Ageing. Whilst most patients will be treated by a consultant in rehabilitation medicine (a Fellow of the Australasian Faculty of Rehabilitation Medicine) there are circumstances in which the treating doctor will not be a Fellow of the Faculty. These include, but are not limited to, care provided in geographic areas where there is a shortage of Fellows of the Australasian Faculty of Rehabilitation Medicine.

Source and reference attributes

Submitting organisation:

Private Rehabilitation Working Group
Commonwealth Department of Health and Ageing

Staging basis of cancer

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Cancer staging – staging basis of cancer, code A
<i>METeOR identifier:</i>	296981
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The timing and evidence for T, N and M cancer stage values, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Cancer staging – staging basis of cancer
<i>Definition:</i>	The timing and evidence for T, N and M cancer stage values.
<i>Context:</i>	For survival analysis adjusted by stage at diagnosis and distribution of cancer cases by type and stage.
<i>Object class:</i>	Cancer staging
<i>Property:</i>	Staging basis of cancer

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	String						
<i>Format:</i>	A						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>P</td><td>Pathological</td></tr><tr><td>C</td><td>Clinical</td></tr></tbody></table>	Value	Meaning	P	Pathological	C	Clinical
Value	Meaning						
P	Pathological						
C	Clinical						

Collection and usage attributes

<i>Guide for use:</i>	<p>CODE P Pathological Pathological stage is based on histological evidence acquired before treatment, supplemented or modified by additional evidence acquired from surgery and from pathological examination.</p> <p>CODE C Clinical Clinical stage is based on evidence obtained prior to treatment from physical examination, imaging, endoscopy, biopsy, surgical exploration or other relevant examinations.</p> <p>Refer to the latest edition of the UICC reference manual TNM Classification of Malignant Tumours for coding rules.</p>
-----------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	Australian Institute of Health and Welfare
---------------------------------	--

Data element attributes

Collection and usage attributes

Collection methods:

From information provided by the treating doctor and recorded on the patient's medical record.

Relational attributes

Implementation in Data Set Specifications:

Cancer (clinical) DSS NHIG, Standard 07/12/2005

Cancer (clinical) DSS NHIG, Superseded 07/12/2005

Staging scheme source

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Cancer staging – cancer staging scheme source, code N
<i>METeOR identifier:</i>	296988
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The reference which describes in detail the methods of staging and the definitions for the classification system used in determining the extent of cancer at the time of diagnosis, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Cancer staging – cancer staging scheme source
<i>Definition:</i>	The reference which describes in detail the methods of staging and the definitions for the classification system used in determining the extent of cancer at the time of diagnosis.
<i>Context:</i>	For survival analysis adjusted by stage at diagnosis and distribution of cancer cases by type and stage.
<i>Object class:</i>	Cancer staging
<i>Property:</i>	Cancer staging scheme source

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code														
<i>Data type:</i>	Number														
<i>Format:</i>	N														
<i>Maximum character length:</i>	1														
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>TNM Classification of Malignant Tumours (UICC)</td></tr><tr><td>2</td><td>Durie & Salmon for multiple myeloma staging</td></tr><tr><td>3</td><td>FAB for leukaemia classification</td></tr><tr><td>4</td><td>Australian Clinico-Pathological Staging (ACPS) System</td></tr><tr><td>8</td><td>Other</td></tr><tr><td>9</td><td>Unknown</td></tr></tbody></table>	Value	Meaning	1	TNM Classification of Malignant Tumours (UICC)	2	Durie & Salmon for multiple myeloma staging	3	FAB for leukaemia classification	4	Australian Clinico-Pathological Staging (ACPS) System	8	Other	9	Unknown
Value	Meaning														
1	TNM Classification of Malignant Tumours (UICC)														
2	Durie & Salmon for multiple myeloma staging														
3	FAB for leukaemia classification														
4	Australian Clinico-Pathological Staging (ACPS) System														
8	Other														
9	Unknown														
<i>Supplementary values:</i>															

Source and reference attributes

<i>Reference documents:</i>	Durie BGM, Salmon SE. <i>A clinical staging system for multiple myeloma correlation of measured myeloma cell mass with presenting clinical features, response to treatment and survival.</i> Cancer 36:842-54 (1975).
	Bennett JM, Catovsky D, Daniel MT, Flandrin G, Galton DA, Gralnick HR, Sultan C. <i>Proposed revised criteria for the classification of acute myeloid leukemia: a report of the French-American-British Cooperative Group.</i> Ann Intern Med 103(4): 620-

625 (1985).

Cheson BD, Cassileth PA, Head DR, Schiffer CA, Bennett JM, Bloomfield CD, Brunning R, Gale RP, Grever MR, Keating MJ, et al. *Report of the National Cancer Institute-sponsored workshop on definitions of diagnosis and response in acute myeloid leukemia*. *J Clin Oncol* 8(5): 813-819 (1990).

Davis NC, Newland RC. *The reporting of colorectal cancer: the Australian Clinicopathological Staging system*. *Aust NZ J Surg* 52:395-397 (1982).

Public Health Division NSW *Clinical Cancer Data Collection for Outcomes and Quality. Data Dictionary Version 1* Sydney NSW Health Dept (2001).

NHMRC *Guidelines for the prevention, early detection and management of colorectal cancer (CRC)* (1999).

Data element attributes

Collection and usage attributes

Guide for use:

It is recommended that the TNM Manual of the UICC be used whenever it is applicable. The classifications published in the American Joint Committee on Cancer (AJCC) Cancer Staging Manual are identical to the TNM classifications of the UICC. TNM is not applicable to all tumour sites. Staging is of limited use in acute leukaemias, although a staging system is used for chronic lymphocytic leukaemia. Separate staging systems exist for lymphomas and myeloma. The *NHMRC Guidelines for the prevention, early detection and management of colorectal cancer (CRC)* support the use of the Australian Clinico-Pathological Staging (ACPS) System. A table of correspondences between ACPS and TNM classifications is available. The current edition of each staging scheme should be used.

Source and reference attributes

Origin:

International Union Against Cancer (UICC).
FAB (French-American-British) Group.
NSW Health Department.
National Health & Medical Research Council.
Clinical Oncological Society of Australia.
Australian Cancer Network.

Relational attributes

Related metadata references:

Supersedes Staging scheme source, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
See also Cancer staging – cancer staging scheme source edition number, code N[N] NHIG, Standard 04/06/2004

Implementation in Data Set Specifications:

Cancer (clinical) DSS NHIG, Standard 07/12/2005
Cancer (clinical) DSS NHIG, Superseded 07/12/2005

Staging scheme source edition number

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Cancer staging – cancer staging scheme source edition number, code N[N]
<i>METeOR identifier:</i>	297011
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The edition of the reference used for the purposes of staging the cancer, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Cancer staging – cancer staging scheme source edition number
<i>Definition:</i>	The edition of the reference used for the purposes of staging the cancer.
<i>Context:</i>	For survival analysis adjusted by stage at diagnosis and distribution of cancer cases by type and stage.
<i>Object class:</i>	Cancer staging
<i>Property:</i>	Cancer staging scheme source edition number

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N[N]						
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>88</td><td>Not applicable (Cases that do not have a recommended staging scheme)</td></tr><tr><td>99</td><td>Unknown edition</td></tr></tbody></table>	Value	Meaning	88	Not applicable (Cases that do not have a recommended staging scheme)	99	Unknown edition
Value	Meaning						
88	Not applicable (Cases that do not have a recommended staging scheme)						
99	Unknown edition						

Collection and usage attributes

<i>Guide for use:</i>	Record the edition number (i.e. 1 - 87).
-----------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	Australian Institute of Health and Welfare
---------------------------------	--

Data element attributes

Source and reference attributes

<i>Origin:</i>	Commission on Cancer, Standards of the Commission on Cancer Registry Operations and Data Standards (ROADS) Volume II (1998).
----------------	--

Relational attributes

<i>Related metadata references:</i>	Supersedes Staging scheme source edition number, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set</i>	Cancer (clinical) DSS NHIG, Standard 07/12/2005

Specifications:

Cancer (clinical) DSS NHIG, Superseded 07/12/2005

State/Territory of birth

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Birth event – state/territory of birth, code N
<i>METeOR identifier:</i>	270151
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The state/territory in which the baby was delivered, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Birth event – state/territory of birth
<i>Definition:</i>	The state/territory in which the baby was delivered.
<i>Object class:</i>	Birth event
<i>Property:</i>	State/territory of birth

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																				
<i>Data type:</i>	Number																				
<i>Format:</i>	N																				
<i>Maximum character length:</i>	1																				
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>New South Wales</td></tr><tr><td>2</td><td>Victoria</td></tr><tr><td>3</td><td>Queensland</td></tr><tr><td>4</td><td>South Australia</td></tr><tr><td>5</td><td>Western Australia</td></tr><tr><td>6</td><td>Tasmania</td></tr><tr><td>7</td><td>Northern Territory</td></tr><tr><td>8</td><td>Australian Capital Territory</td></tr><tr><td>9</td><td>Other territories (Cocos (Keeling) Islands, Christmas Island and Jervis Bay Territory)</td></tr></tbody></table>	Value	Meaning	1	New South Wales	2	Victoria	3	Queensland	4	South Australia	5	Western Australia	6	Tasmania	7	Northern Territory	8	Australian Capital Territory	9	Other territories (Cocos (Keeling) Islands, Christmas Island and Jervis Bay Territory)
Value	Meaning																				
1	New South Wales																				
2	Victoria																				
3	Queensland																				
4	South Australia																				
5	Western Australia																				
6	Tasmania																				
7	Northern Territory																				
8	Australian Capital Territory																				
9	Other territories (Cocos (Keeling) Islands, Christmas Island and Jervis Bay Territory)																				

Collection and usage attributes

<i>Guide for use:</i>	The order presented here is the standard for the Australian Bureau of Statistics (ABS). Other organisations (including the Australian Institute of Health and Welfare) publish data in state order based on population (that is, Western Australia before South Australia and Australian Capital Territory before Northern Territory).
-----------------------	--

Source and reference attributes

<i>Reference documents:</i>	Australian Bureau of Statistics 2005. Australian Standard Geographical Classification (ASGC). Cat. no. 1216.0. Canberra: ABS. Viewed on 30/09/2005
-----------------------------	--

Data element attributes

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Related metadata references: Supersedes State/Territory of birth, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Health care client identification NHIG, Superseded 04/05/2005
Implementation start date: 01/01/2003

Health care client identification DSS NHIG, Standard 04/05/2005

NCSIMG, Standard 03/10/2006

Perinatal NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Perinatal NMDS NHIG, Superseded 06/09/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Perinatal NMDS 2007-2008 NHIG, Standard 06/09/2006

Implementation start date: 01/07/2007

Status of the baby

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Birth—birth status, code N
<i>METeOR identifier:</i>	269949
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The status of the baby at birth as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Birth—birth status
<i>Definition:</i>	Status of the baby at birth.
<i>Context:</i>	Perinatal statistics
<i>Object class:</i>	Birth
<i>Property:</i>	Birth status

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code								
<i>Data type:</i>	Number								
<i>Format:</i>	N								
<i>Maximum character length:</i>	1								
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Live birth</td></tr><tr><td>2</td><td>Stillbirth (fetal death)</td></tr><tr><td>9</td><td>Not stated</td></tr></tbody></table>	Value	Meaning	1	Live birth	2	Stillbirth (fetal death)	9	Not stated
Value	Meaning								
1	Live birth								
2	Stillbirth (fetal death)								
9	Not stated								
<i>Supplementary values:</i>									

Collection and usage attributes

<i>Guide for use:</i>	<p>Live birth is the complete expulsion or extraction from its mother of a product of conception, irrespective of the duration of the pregnancy which, after such separation, breathes or shows any other evidence of life, such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles, whether or not the umbilical cord has been cut or the placenta is attached; each product of such a birth is considered liveborn (WHO, 1992 definition).</p> <p>Stillbirth is a fetal death prior to the complete expulsion or extraction from its mother of a product of conception of 20 or more completed weeks of gestation or of 400 g or more birthweight; the death is indicated by the fact that after such separation the fetus does not breathe or show any other evidence of life, such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles. (This is the same as the WHO definition of fetal death, except that there are no limits of gestational age or birthweight for the WHO definition.)</p>
-----------------------	--

Source and reference attributes

Reference documents:

International Classification of Diseases and Related Health Problems, 10th Revision, Vol 1, WHO 1992.

Data element attributes

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Related metadata references: Supersedes Status of the baby, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Perinatal NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Perinatal NMDS NHIG, Superseded 06/09/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

Essential to analyse outcome of pregnancy.

Perinatal NMDS 2007-2008 NHIG, Standard 06/09/2006

Implementation start date: 01/07/2007

Information specific to this data set:

Essential to analyse outcome of pregnancy.

Street name (person)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person (address)—street name, text [A(30)]
<i>METeOR identifier:</i>	270019
<i>Registration status:</i>	NHIG, Standard 01/03/2005 NCSIMG, Standard 30/09/2005
<i>Definition:</i>	The concatenation of a person's street type and street suffix resulting in a name that identifies a public thoroughfare and differentiates it from others in the same suburb/town/locality, as represented by text.

Data element concept attributes

<i>Data element concept:</i>	Person (address)—street name
<i>Definition:</i>	The concatenation of a person's street type and street suffix resulting in a name that identifies a public thoroughfare and differentiates it from others in the same suburb/town/locality.
<i>Object class:</i>	Person
<i>Property:</i>	Street name

Value domain attributes

Representational attributes

<i>Representation class:</i>	Text
<i>Data type:</i>	String
<i>Format:</i>	[A(30)]
<i>Maximum character length:</i>	30

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	To be used in conjunction with street type. To be used in conjunction with street suffix.
<i>Comments:</i>	Where suburb/town/locality, state/territory and Postcode - Australian are insufficient to assign a Statistical Local Area (SLA) code from the Australian Standard Geographical Classification (Australian Bureau of Statistics, Cat. No. 1216.0) , the Street name metadata item in conjunction with street type, house/property identifier and street suffix should also be used.

Source and reference attributes

<i>Origin:</i>	Health Data Standards Committee Australia Post Address Presentation Standard
----------------	---

Relational attributes

<i>Related metadata references:</i>	Supersedes Street name, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005 Is used in the formation of Person (address)—address line, text
-------------------------------------	---

Implementation in Data Set Specifications:

[X(180)] NHIG, Standard 04/05/2005, NCSIMG, Standard 30/09/2005

Is used in the formation of Person (address) – health address line, text [X(180)] NHIG, Superseded 04/05/2005

Health care client identification DSS NHIG, Standard 04/05/2005

NCSIMG, Standard 03/10/2006

Health care provider identification DSS NHIG, Standard 04/05/2005

Street name (service provider organisation)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Service provider organisation (address) – street name, text [A(30)]
<i>METeOR identifier:</i>	290218
<i>Registration status:</i>	NHIG, Standard 04/05/2005 NCSIMG, Standard 30/09/2005
<i>Definition:</i>	The concatenation of an organisation's street type and street suffix resulting in a name that identifies a public thoroughfare and differentiates it from others in the same suburb/town/locality, as represented by text.

Data element concept attributes

<i>Data element concept:</i>	Service provider organisation (address) – street name
<i>Definition:</i>	The concatenation of an organisation's street type and street suffix resulting in a name that identifies a public thoroughfare and differentiates it from others in the same suburb/town/locality.
<i>Object class:</i>	Service provider organisation
<i>Property:</i>	Street name

Value domain attributes

Representational attributes

<i>Representation class:</i>	Text
<i>Data type:</i>	String
<i>Format:</i>	[A(30)]
<i>Maximum character length:</i>	30

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	To be used in conjunction with street type. To be used in conjunction with street suffix.
<i>Comments:</i>	Where suburb/town/locality, state/territory and Postcode - Australian are insufficient to assign a Statistical Local Area (SLA) code from the Australian Standard Geographical Classification (Australian Bureau of Statistics, Cat. No. 1216.0) , the Street name metadata item in conjunction with street type, house/property identifier and street suffix should also be used.

Source and reference attributes

<i>Origin:</i>	Health Data Standards Committee Australia Post Address Presentation Standard
----------------	---

Relational attributes

<i>Related metadata references:</i>	Is used in the formation of Service provider organisation
-------------------------------------	---

*Implementation in Data Set
Specifications:*

(address)—address line, text [X(180)] NHIG, Standard
04/05/2005, NCSIMG, Standard 30/09/2005

Health care provider identification DSS NHIG, Standard
04/05/2005

Street suffix code (person)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person (address)—street suffix, code A[A]
<i>METeOR identifier:</i>	270022
<i>Registration status:</i>	NHIG, Standard 01/03/2005 NCSIMG, Standard 30/09/2005
<i>Definition:</i>	The abbreviated suffix that identifies the type of street where a person resides, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person (address)—street suffix
<i>Definition:</i>	The abbreviated suffix that identifies the type of street where a person resides.
<i>Object class:</i>	Person
<i>Property:</i>	Street suffix

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																										
<i>Data type:</i>	String																										
<i>Format:</i>	A[A]																										
<i>Maximum character length:</i>	2																										
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>CN</td><td>Central</td></tr><tr><td>E</td><td>East</td></tr><tr><td>EX</td><td>Extension</td></tr><tr><td>LR</td><td>Lower</td></tr><tr><td>N</td><td>North</td></tr><tr><td>NE</td><td>North East</td></tr><tr><td>NW</td><td>North West</td></tr><tr><td>S</td><td>South</td></tr><tr><td>SE</td><td>South East</td></tr><tr><td>SW</td><td>South West</td></tr><tr><td>UP</td><td>Upper</td></tr><tr><td>W</td><td>West</td></tr></tbody></table>	Value	Meaning	CN	Central	E	East	EX	Extension	LR	Lower	N	North	NE	North East	NW	North West	S	South	SE	South East	SW	South West	UP	Upper	W	West
Value	Meaning																										
CN	Central																										
E	East																										
EX	Extension																										
LR	Lower																										
N	North																										
NE	North East																										
NW	North West																										
S	South																										
SE	South East																										
SW	South West																										
UP	Upper																										
W	West																										

Data element attributes

Collection and usage attributes

<i>Collection methods:</i>	To be used in conjunction with street name. To be used in conjunction with street type. For example: Browns Rd W
----------------------------	--

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare
Origin: AS4590 Interchange of client information, Australia Post
Address Presentation Standard

Relational attributes

Related metadata references: Supersedes Street suffix code, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
Is used in the formation of Person (address) – address line, text [X(180)] NHIG, Standard 04/05/2005, NCSIMG, Standard 30/09/2005
Is used in the formation of Person (address) – health address line, text [X(180)] NHIG, Superseded 04/05/2005

Implementation in Data Set Specifications: Health care client identification DSS NHIG, Standard 04/05/2005
NCSIMG, Standard 03/10/2006
Health care provider identification DSS NHIG, Standard 04/05/2005

Street suffix code (service provider organisation)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Service provider organisation (address) – street suffix, code A[A]
<i>METeOR identifier:</i>	290170
<i>Registration status:</i>	NHIG, Standard 04/05/2005 NCSIMG, Standard 30/09/2005
<i>Definition:</i>	The abbreviated suffix that identifies the type of street where an organisation is located, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Service provider organisation (address) – street suffix
<i>Definition:</i>	The abbreviated suffix that identifies the type of street where an organisation is located.
<i>Object class:</i>	Service provider organisation
<i>Property:</i>	Street suffix

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																										
<i>Data type:</i>	String																										
<i>Format:</i>	A[A]																										
<i>Maximum character length:</i>	2																										
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>CN</td><td>Central</td></tr><tr><td>E</td><td>East</td></tr><tr><td>EX</td><td>Extension</td></tr><tr><td>LR</td><td>Lower</td></tr><tr><td>N</td><td>North</td></tr><tr><td>NE</td><td>North East</td></tr><tr><td>NW</td><td>North West</td></tr><tr><td>S</td><td>South</td></tr><tr><td>SE</td><td>South East</td></tr><tr><td>SW</td><td>South West</td></tr><tr><td>UP</td><td>Upper</td></tr><tr><td>W</td><td>West</td></tr></tbody></table>	Value	Meaning	CN	Central	E	East	EX	Extension	LR	Lower	N	North	NE	North East	NW	North West	S	South	SE	South East	SW	South West	UP	Upper	W	West
Value	Meaning																										
CN	Central																										
E	East																										
EX	Extension																										
LR	Lower																										
N	North																										
NE	North East																										
NW	North West																										
S	South																										
SE	South East																										
SW	South West																										
UP	Upper																										
W	West																										

Data element attributes

Collection and usage attributes

<i>Collection methods:</i>	To be used in conjunction with street name. To be used in conjunction with street type. For example:
----------------------------	---

Browns Rd W

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare
Origin: AS4590 Interchange of client information, Australia Post
Address Presentation Standard

Relational attributes

Related metadata references: Is used in the formation of Service provider organisation (address) – address line, text [X(180)] NHIG, Standard 04/05/2005, NCSIMG, Standard 30/09/2005
Implementation in Data Set Specifications: Health care provider identification DSS NHIG, Standard 04/05/2005

Street type code (person)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person (address)—street type, code A[AAA]
<i>METeOR identifier:</i>	270020
<i>Registration status:</i>	NHIG, Standard 01/03/2005 NCSIMG, Standard 30/09/2005
<i>Definition:</i>	The type of public thoroughfare where a person resides, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person (address)—street type
<i>Definition:</i>	The type of public thoroughfare where a person resides.
<i>Object class:</i>	Person
<i>Property:</i>	Street type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code
<i>Data type:</i>	String
<i>Format:</i>	A[AAA]
<i>Maximum character length:</i>	4

Collection and usage attributes

Guide for use: The following is a list of commonly used abbreviations from AS 4590:

Street type	Abbreviation
Alley	Ally
Arcade	Arc
Avenue	Ave
Boulevard	Bvd
Bypass	Bypa
Circuit	Cct
Close	Cl
Corner	Crn
Court	Ct
Crescent	Cres
Cul-de-sac	Cds
Drive	Dr
Esplanade	Esp
Green	Grn
Grove	Gr
Highway	Hwy
Junction	Jnc
Lane	Lane
Link	Link

Mews	Mews
Parade	Pde
Place	Pl
Ridge	Rdge
Road	Rd
Square	Sq
Street	St
Terrace	Tce

Data element attributes

Source and reference attributes

<i>Submitting organisation:</i>	Standards Australia
<i>Origin:</i>	Health Data Standards Committee AS4590 Interchange of client information, Australia Post Address Presentation Standard

Relational attributes

<i>Related metadata references:</i>	Supersedes Street type code, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005 Is used in the formation of Person (address) – address line, text [X(180)] NHIG, Standard 04/05/2005, NCSIMG, Standard 30/09/2005 Is used in the formation of Person (address) – health address line, text [X(180)] NHIG, Superseded 04/05/2005
<i>Implementation in Data Set Specifications:</i>	Health care client identification DSS NHIG, Standard 04/05/2005 NCSIMG, Standard 03/10/2006 Health care provider identification DSS NHIG, Standard 04/05/2005

Street type code (service provider organisation)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Service provider organisation (address) – street type, code A[AAA]
<i>METeOR identifier:</i>	290193
<i>Registration status:</i>	NHIG, Standard 04/05/2005 NCSIMG, Standard 30/09/2005
<i>Definition:</i>	The type of public thoroughfare where an organisation is located, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Service provider organisation (address) – street type
<i>Definition:</i>	The type of public thoroughfare where an organisation is located.
<i>Object class:</i>	Service provider organisation
<i>Property:</i>	Street type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code
<i>Data type:</i>	String
<i>Format:</i>	A[AAA]
<i>Maximum character length:</i>	4

Collection and usage attributes

Guide for use: The following is a list of commonly used abbreviations from AS 4590:

Street type	Abbreviation
Alley	Ally
Arcade	Arc
Avenue	Ave
Boulevard	Bvd
Bypass	Bypa
Circuit	Cct
Close	Cl
Corner	Crn
Court	Ct
Crescent	Cres
Cul-de-sac	Cds
Drive	Dr
Esplanade	Esp
Green	Grn
Grove	Gr
Highway	Hwy
Junction	Jnc

Lane	Lane
Link	Link
Mews	Mews
Parade	Pde
Place	Pl
Ridge	Rdge
Road	Rd
Square	Sq
Street	St
Terrace	Tce

Data element attributes

Collection and usage attributes

Collection methods: To be collected in conjunction with street name. To be collected in conjunction with street suffix.

Source and reference attributes

Origin: AS4590 Interchange of client information, Australia Post Address Presentation Standard

Relational attributes

Related metadata references: Is used in the formation of Service provider organisation (address) – address line, text [X(180)] NHIG, Standard 04/05/2005, NCSIMG, Standard 30/09/2005

Implementation in Data Set Specifications: Health care provider identification DSS NHIG, Standard 04/05/2005

Suburb/town/locality name (person)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person (address)—suburb/town/locality name, text [A(50)]
<i>METeOR identifier:</i>	287326
<i>Registration status:</i>	NHIG, Standard 04/05/2005 NCSIMG, Standard 25/08/2005
<i>Definition:</i>	The full name of the locality contained within the specific address of a person, as represented by text.

Data element concept attributes

<i>Data element concept:</i>	Person (address)—suburb/town/locality name
<i>Definition:</i>	The full name of the locality contained within the specific address of a person.
<i>Context:</i>	In conjunction with the postcode, the data element Suburb/town/locality name is included as an alternative means of reporting information about the geographic location of the residence of a client, or an agency/establishment or where an event occurred. The preferred standard for reporting this information is by using a statistical local area (SLA) in conjunction with a State/Territory code. However, as some agencies may have difficulty allocating SLA codes to the residential locations of their clients without more computerised assistance than is currently available to them, agencies may be given the option of reporting this information by using Australian postcode plus Suburb/town/locality name. Suburb/town/locality name may also be a component of a postal address.
<i>Object class:</i>	Person
<i>Property:</i>	Suburb/town/locality name

Value domain attributes

Representational attributes

<i>Representation class:</i>	Text
<i>Data type:</i>	String
<i>Format:</i>	[A(50)]
<i>Maximum character length:</i>	50

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	The suburb/town/locality name may be a town, city, suburb or commonly used location name such as a large agricultural property or Aboriginal community. This metadata item may be used to describe the location of person. It can be a component of a street or postal address. The Australian Bureau of Statistics has suggested that a
-----------------------	--

maximum field length of 50 characters should be sufficient to record the vast majority of locality names.

Collection methods:

Enter 'Unknown' when the locality name or geographic area for a person or event is not known. Enter 'No fixed address' when a person has no fixed address or is **homeless**.

Source and reference attributes

Origin:

National Health Data Committee
National Community Services Data Committee

Reference documents:

AS5017 Health Care Client Identification, 2002, Sydney:
Standards Australia
AS4846 Health Care Provider Identification, 2004, Sydney:
Standards Australia
Australia Post 2005. Australia Postcode File.
Viewed 12 April, www.auspost.com.au/postcodes

Relational attributes

Related metadata references:

Supersedes Person (address) – suburb/town/locality name, text [A(50)] NHIG, Superseded 04/05/2005, NCSIMG, Superseded 25/08/2005
Is used in the formation of Person (address) – postal delivery point identifier, {N(8)} NHIG, Standard 04/05/2005, NCSIMG, Standard 25/08/2005
Is used in the formation of Dwelling – geographic location, remoteness structure code (ASGC 2004) N[N] NHDAMG, Retired 10/02/2006

Implementation in Data Set Specifications:

Computer Assisted Telephone Interview demographic module DSS NHIG, Standard 04/05/2005

Information specific to this data set:

For data collection using Computer Assisted Telephone Interviewing (CATI), the suggested question is:
What is your suburb, town or community?
(Single response)
Enter town/suburb/community.

Health care client identification DSS NHIG, Standard 04/05/2005
NCSIMG, Standard 03/10/2006

Information specific to this data set:

This data should be verified against the Australia Post Postcode File (see www.auspost.com.au/postcodes). Alternatively, contact State or Territory Health Authorities for Postcode files.

Health care provider identification DSS NHIG, Standard 04/05/2005

Information specific to this data set:

This data should be verified against the Australia Post Postcode File (see www.auspost.com.au/postcodes). Alternatively, contact State or Territory Health Authorities for Postcode files.

Suburb/town/locality name (service provider organisation)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Service provider organisation (address) – suburb/town/locality name, text [A(50)]
<i>METeOR identifier:</i>	290059
<i>Registration status:</i>	NHIG, Standard 04/05/2005 NCSIMG, Standard 31/08/2005
<i>Definition:</i>	The full name of the general locality containing the specific address of an organisation, as represented by text.

Data element concept attributes

<i>Data element concept:</i>	Service provider organisation (address) – suburb/town/locality name
<i>Definition:</i>	The full name of the general locality containing the specific address of an organisation.
<i>Context:</i>	In conjunction with the data element Postcode-Australian, the data element Suburb/town/locality name is included as an alternative means of reporting information about the geographic location of an agency/establishment. The preferred standard for reporting this information is by using a statistical local area (SLA) in conjunction with a State/Territory code. Suburb/town/locality name may also be a component of a postal address.
<i>Object class:</i>	Service provider organisation
<i>Property:</i>	Suburb/town/locality name

Value domain attributes

Representational attributes

<i>Representation class:</i>	Text
<i>Data type:</i>	String
<i>Format:</i>	[A(50)]
<i>Maximum character length:</i>	50

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>The suburb/town/locality name, may be a town, city, suburb or commonly used location name such as a large agricultural property or Aboriginal community.</p> <p>The Australian Bureau of Statistics has suggested that a maximum field length of 50 characters should be sufficient to record the vast majority of locality names.</p> <p>This metadata item may be used to describe the location of an organisation. It can be a component of a street or postal address.</p>
-----------------------	---

Collection methods: Enter 'Unknown' when the locality name or geographic area for an organisation is not known.

Source and reference attributes

Origin: National Health Data Committee
National Community Services Data Committee
Australia Post 2005. Australia Postcode File. Viewed 12 April
www.auspost.com.au/postcodes

Reference documents: AS5017 Health Care Client Identification, 2002, Sydney:
Standards Australia
AS4846 Health Care Provider Identification, 2004, Sydney:
Standards Australia

Relational attributes

Related metadata references: Is used in the formation of Service provider organisation (address) – postal delivery point identifier, {N(8)} NHIG, Standard 04/05/2005, NCSIMG, Standard 31/08/2005

Implementation in Data Set Specifications: Health care provider identification DSS NHIG, Standard 04/05/2005

Information specific to this data set:

This data should be verified against the Australia Post Postcode File (see www.auspost.com.au/postcodes). Alternatively, contact State or Territory Health Authorities for Postcode files.

Surgical specialty

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Elective surgery waiting list episode – surgical specialty (of scheduled doctor), code NN
<i>METeOR identifier:</i>	270146
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The area of clinical expertise held by the doctor who will perform the elective surgery, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Elective surgery waiting list episode – surgical specialty (of scheduled doctor)
<i>Definition:</i>	The area of clinical expertise held by the doctor who will perform the elective surgery.
<i>Context:</i>	Elective surgery: Many hospitals manage their waiting lists on a specialty basis. Current data shows that the total ready-for-care times waited and numbers of long wait patients vary significantly between specialties. Furthermore, the hospital capacity to handle the demand for elective surgery varies with specialty.
<i>Object class:</i>	Elective surgery waiting list episode
<i>Property:</i>	Surgical specialty

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																								
<i>Data type:</i>	String																								
<i>Format:</i>	NN																								
<i>Maximum character length:</i>	2																								
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>Cardio-thoracic surgery</td></tr><tr><td>02</td><td>Ear, nose and throat surgery</td></tr><tr><td>03</td><td>General surgery</td></tr><tr><td>04</td><td>Gynaecology</td></tr><tr><td>05</td><td>Neurosurgery</td></tr><tr><td>06</td><td>Ophthalmology</td></tr><tr><td>07</td><td>Orthopaedic surgery</td></tr><tr><td>08</td><td>Plastic surgery</td></tr><tr><td>09</td><td>Urology</td></tr><tr><td>10</td><td>Vascular surgery</td></tr><tr><td>11</td><td>Other</td></tr></tbody></table>	Value	Meaning	01	Cardio-thoracic surgery	02	Ear, nose and throat surgery	03	General surgery	04	Gynaecology	05	Neurosurgery	06	Ophthalmology	07	Orthopaedic surgery	08	Plastic surgery	09	Urology	10	Vascular surgery	11	Other
Value	Meaning																								
01	Cardio-thoracic surgery																								
02	Ear, nose and throat surgery																								
03	General surgery																								
04	Gynaecology																								
05	Neurosurgery																								
06	Ophthalmology																								
07	Orthopaedic surgery																								
08	Plastic surgery																								
09	Urology																								
10	Vascular surgery																								
11	Other																								

Collection and usage attributes

Comments:

The above classifications are consistent with the Recommended Medical Specialties and Qualifications agreed by the National Specialist Qualification Advisory Committee of Australia, September 1993. Vascular surgery is a subspecialty of general surgery. The Royal Australian College of Surgeons has a training program for vascular surgeons. The specialties listed above refer to the surgical component of these specialties - ear, nose and throat surgery refers to the surgical component of the specialty otolaryngology; gynaecology refers to the gynaecological surgical component of obstetrics and gynaecology; ophthalmology refers to the surgical component of the specialty (patients awaiting argon laser phototherapy are not included).

Data element attributes

Source and reference attributes

Submitting organisation: Hospital Access Program Waiting Lists Working Group
Waiting Times Working Group

Origin: National Health Data Committee

Relational attributes

Related metadata references: Supersedes Surgical specialty, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Elective surgery waiting times (census data) NMDS NHIG, Standard 07/12/2005

Implementation start date: 30/09/2006

Elective surgery waiting times (census data) NMDS NHIG, Superseded 07/12/2005

Implementation start date: 30/09/2002

Implementation end date: 30/06/2006

Elective surgery waiting times (removals data) NMDS NHIG, Standard 07/12/2005

Implementation start date: 01/07/2006

Elective surgery waiting times (removals data) NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2002

Implementation end date: 30/06/2006

Surgical treatment procedure for cancer

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Cancer treatment – surgical procedure for cancer, procedure code (ACHI 5th edn) NNNNN-NN
<i>METeOR identifier:</i>	333816
<i>Registration status:</i>	NHIG, Standard 07/12/2005
<i>Definition:</i>	The surgical procedure used in the primary treatment of the cancer, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Cancer treatment – surgical procedure for cancer
<i>Definition:</i>	The surgical procedure used in the primary treatment of the cancer.
<i>Object class:</i>	Cancer treatment
<i>Property:</i>	Surgical procedure for cancer

Value domain attributes

Representational attributes

<i>Classification scheme:</i>	Australian Classification of Health Interventions (ACHI) 5th edition
<i>Representation class:</i>	Code
<i>Data type:</i>	Number
<i>Format:</i>	NNNNN-NN
<i>Maximum character length:</i>	7

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Each surgical treatment procedure used in the initial treatment of the cancer should be recorded. Surgical procedures performed for palliative purposes only should not be included. For surgical procedures involved in the administration of another modality (eg., implantation of infusion pump, isolated limb perfusion/infusion, intra-operative radiotherapy) record both the surgery and the other modality. Any systemic treatment which can be coded as a procedure through ACHI should be so coded (eg., stem cell or bone marrow infusion).
-----------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	National Cancer Control Initiative
<i>Origin:</i>	National Centre for Classification in Health New South Wales Department of Health, Public Health Division
<i>Reference documents:</i>	NSW Department of Health NSW Clinical Cancer Data

Collection for Outcomes and Quality. Data Dictionary Version 1 (2001).

Relational attributes

Related metadata references:

Supersedes Cancer treatment – surgical procedure for cancer, procedure code (ICD-10-AM 4th edn) NNNNN-NN NHIG, Superseded 07/12/2005

Implementation in Data Set Specifications:

Cancer (clinical) DSS NHIG, Standard 07/12/2005

Information specific to this data set:

This item is collected for determining outcome by treatment type.

Systemic therapy agent name

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Cancer treatment – systemic therapy agent name (primary cancer), antineoplastic drug code (Self-Instructional Manual for Tumour Registrars Book 8 3rd edn) X[X(39)]
<i>METeOR identifier:</i>	288446
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The chemotherapeutic agent or anti-cancer drug used for treatment of the primary cancer, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Cancer treatment – systemic therapy agent name (primary cancer)
<i>Definition:</i>	The standard chemotherapeutic agent or anti-cancer drug used for treatment of the primary cancer.
<i>Object class:</i>	Cancer treatment
<i>Property:</i>	Systemic therapy agent name

Value domain attributes

Representational attributes

<i>Classification scheme:</i>	Self-Instructional Manual for Tumour Registrars Book 8 Antineoplastic Drugs, 3rd edition
<i>Representation class:</i>	Code
<i>Data type:</i>	String
<i>Format:</i>	X[X(39)]
<i>Maximum character length:</i>	40

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>The purpose of collecting specific treatment information is to account for all treatment types, which may assist in evaluation of effectiveness of different treatment patterns. The actual agents used will sometimes be of interest.</p> <p>Systemic therapy often involves treatment with a combination of agents. These may be known by acronyms but since details of drugs and acronyms may vary it is recommended that each agent be recorded separately.</p> <p>Oral chemotherapy normally given on an outpatient basis should also be included.</p> <p>New codes and names will need to be added as new agents become available for clinical use.</p> <p>Hormone therapy agents and immunotherapy agents should be recorded under this data element.</p>
<i>Collection methods:</i>	The full name of the agent(s) should be recorded if the coding manual is not available.

Comments: Collecting dates for systemic therapy will allow evaluation of treatments delivered and of time intervals from diagnosis to treatment, from treatment to recurrence and from treatment to death.

Source and reference attributes

Origin: National Cancer Institute Surveillance, Epidemiology and End Results (SEER) Program

Reference documents: Surveillance, Epidemiology and End Results (SEER) Program Self-instructional manual for tumour registrars: Book 8 - Antineoplastic drugs 3rd Edition National Cancer Institute.

Relational attributes

Related metadata references: Supersedes Systemic therapy agent name, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Cancer (clinical) DSS NHIG, Standard 07/12/2005

Information specific to this data set:

This item is collected for the analysis of outcome by treatment type.

Cancer (clinical) DSS NHIG, Superseded 07/12/2005

Information specific to this data set:

This item is collected for the analysis of outcome by treatment type.

Teaching status

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment – teaching status (university affiliation), code N
<i>METeOR identifier:</i>	270148
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	An indicator to identify the non-direct patient care activity of teaching for a particular establishment, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Establishment – teaching status (university affiliation)
<i>Definition:</i>	An indicator (yes / no) to identify the non-direct patient care activity of teaching for a particular establishment. This is where teaching (associated with a university) is a major program activity of the establishment. It is primarily intended to relate to teaching hospitals affiliated with universities providing undergraduate medical education as advised by the relevant state health authority.
<i>Context:</i>	Health services: The non-direct care activity of teaching can involve the consumption of considerable resources. In comparisons of cost in relation to establishment output, it is important to be aware of particular establishments which are devoting substantial resources to activities not relating to output as measured in terms of either in-patient bed days or outpatient occasions of service. Teaching can be one of the variables in any regression analysis undertaken.
<i>Object class:</i>	Establishment
<i>Property:</i>	Teaching status

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code								
<i>Data type:</i>	Number								
<i>Format:</i>	N								
<i>Maximum character length:</i>	1								
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr><tr><td>9</td><td>Unknown</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No	9	Unknown
Value	Meaning								
1	Yes								
2	No								
9	Unknown								
<i>Supplementary values:</i>									

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	In this context, teaching relates to teaching hospitals affiliated
-----------------------	--

with universities providing undergraduate medical education as advised by the relevant state health authority.

Comments:

The initial intention based on the Taskforce on National Hospital Statistics approach had been to have non-direct care activity indicators for all of the following non-direct patient care activities:

- teaching
- research
- group or community contacts
- public health activities
- mobile centre and/or part-time service.

However, the Resources Working Party decided to delete 2, 3, 4 and 5 and place the emphasis on teaching where teaching (associated with a university) was a major program activity of the hospital. The working party took the view that it was extremely difficult to identify research activities in health institutions because many staff consider that they do research as part of their usual duties. The research indicator was thus deleted and the teaching indicator was agreed to relate to teaching hospitals affiliated with universities providing undergraduate medical education, as advised by the relevant state health authority. If a teaching hospital is identified by a Yes/no indicator then it is not necessary to worry about research (based on the assumption that if you have teaching, you have research).

Source and reference attributes

Origin: National Health Data Committee

Relational attributes

Related metadata references: Supersedes Teaching status, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Public hospital establishments NMDS NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Public hospital establishments NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Public hospital establishments NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Telephone number

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – telephone number, text [X(40)]
<i>METeOR identifier:</i>	270266
<i>Registration status:</i>	NHIG, Standard 01/03/2005 NCSIMG, Standard 01/03/2005
<i>Definition:</i>	The person's contact telephone number, as represented by text.

Data element concept attributes

<i>Data element concept:</i>	Person – telephone number
<i>Definition:</i>	Person contact telephone number.
<i>Object class:</i>	Person
<i>Property:</i>	Telephone number

Value domain attributes

Representational attributes

<i>Representation class:</i>	Text
<i>Data type:</i>	String
<i>Format:</i>	[X(40)]
<i>Maximum character length:</i>	40

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	More than one phone number may be recorded as required. Each phone number should have an appropriate telephone number type code assigned. Record the full phone number (including any prefixes) with no punctuation (hyphens or brackets).
<i>Collection methods:</i>	Prefix plus telephone number: Record the prefix plus telephone number. The default should be the local prefix with an ability to overtype with a different prefix. For example, 08 8226 6000 or 0417 123456. Punctuation: Do not record punctuation. For example, (08) 8226 6000 or 08-8226 6000 would not be correct. Unknown: Leave the field blank.
<i>Comments:</i>	Concerned with the use of person identification data. For organisations that create, use or maintain records on people. Organisations should use this standard, where appropriate, for collecting data when registering people. The positive and unique identification of people is a critical event in service

delivery, with direct implications for the safety and quality of care delivered by health and community services.

Source and reference attributes

Submitting organisation: Standards Australia
Origin: National Health Data Committee
National Community Services Data Committee
Standards Australia 2002. Australian Standard AS5017-2002
Health Care Client Identification. Sydney: Standards Australia

Relational attributes

Related metadata references: Supersedes Telephone number, version 2, DE, Int. NCSDD & NHDD, NCSIMG & NHIMG, Superseded 01/03/2005
Implementation in Data Set Specifications: Health care client identification NHIG, Superseded 04/05/2005
Implementation start date: 01/01/2003
Information specific to this data set:
Numbers and spaces only.

Telephone number type

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person (telephone) – telephone number type, code A
<i>METeOR identifier:</i>	270299
<i>Registration status:</i>	NHIG, Standard 01/03/2005 NCSIMG, Standard 01/03/2005
<i>Definition:</i>	The type of telephone number recorded for a person, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person (telephone) – telephone number type
<i>Definition:</i>	The type of telephone number recorded for a person.
<i>Object class:</i>	Person
<i>Property:</i>	Telephone number type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code														
<i>Data type:</i>	String														
<i>Format:</i>	A														
<i>Maximum character length:</i>	1														
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>B</td><td>Business or work</td></tr><tr><td>H</td><td>Home</td></tr><tr><td>M</td><td>Personal mobile</td></tr><tr><td>N</td><td>Contact number (not own)</td></tr><tr><td>O</td><td>Business or work mobile</td></tr><tr><td>T</td><td>Temporary</td></tr></tbody></table>	Value	Meaning	B	Business or work	H	Home	M	Personal mobile	N	Contact number (not own)	O	Business or work mobile	T	Temporary
Value	Meaning														
B	Business or work														
H	Home														
M	Personal mobile														
N	Contact number (not own)														
O	Business or work mobile														
T	Temporary														

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Where more than one telephone number has been recorded, then each telephone number should have the appropriate telephone number type code assigned.
<i>Comments:</i>	Concerned with the use of person identification data. For organisations that create, use or maintain records on people. Organisations should use this standard, where appropriate, for collecting data when registering people. The positive and unique identification of people is a critical event in service delivery, with direct implications for the safety and quality of care delivered by health and community services.

Source and reference attributes

Submitting organisation: Standards Australia
Origin: National Health Data Committee
National Community Services Data Committee
Reference documents: Standards Australia 2002. Australian Standard AS5017-2002
Health Care Client Identification. Sydney: Standards Australia

Relational attributes

Related metadata references: Supersedes Telephone number type, version 2, DE, Int. NCSDD & NHDD, NCSIMG & NHIMG, Superseded 01/03/2005
Implementation in Data Set Specifications: Health care client identification NHIG, Superseded 04/05/2005
Implementation start date: 01/01/2003

Time creatine kinase MB isoenzyme measured

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – creatine kinase myocardial band isoenzyme measured time, hhmm
<i>METeOR identifier:</i>	285179
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The time at which the person's creatine kinase myocardial band (CK-MB) isoenzyme was measured.

Data element concept attributes

<i>Data element concept:</i>	Person – creatine kinase myocardial band isoenzyme measured time
<i>Definition:</i>	The time at which the person's creatine kinase myocardial band (CK-MB) isoenzyme was measured.
<i>Context:</i>	Health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Creatine kinase myocardial band isoenzyme measured time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Time
<i>Data type:</i>	Date/Time
<i>Format:</i>	hhmm
<i>Maximum character length:</i>	4

Source and reference attributes

<i>Reference documents:</i>	ISO 8601:2000 : Data elements and interchange formats - Information interchange - Representation of dates and times
-----------------------------	---

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record the time in 24-hour clock format.
-----------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	Acute coronary syndrome data working group
<i>Steward:</i>	The National Heart Foundation of Australia and The Cardiac Society of Australia and New Zealand

Relational attributes

<i>Related metadata references:</i>	Supersedes Time creatine kinase MB isoenzyme (CK-MB) measured, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded
07/12/2005

Time of first angioplasty balloon inflation or stenting

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – first angioplasty balloon inflation or stenting time, hhmm
<i>METeOR identifier:</i>	285191
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The time of the first angioplasty balloon inflation or stent placement.

Data element concept attributes

<i>Data element concept:</i>	Person – first angioplasty balloon inflation or stenting time
<i>Definition:</i>	The time of the first angioplasty balloon inflation or stent placement.
<i>Context:</i>	Health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	First angioplasty balloon inflation or stenting time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Time
<i>Data type:</i>	Date/Time
<i>Format:</i>	hhmm
<i>Maximum character length:</i>	4

Source and reference attributes

<i>Reference documents:</i>	ISO 8601:2000 : Data elements and interchange formats - Information interchange - Representation of dates and times
-----------------------------	---

Data element attributes

Source and reference attributes

<i>Submitting organisation:</i>	Acute coronary syndrome data working group
<i>Steward:</i>	The National Heart Foundation of Australia and The Cardiac Society of Australia and New Zealand

Relational attributes

<i>Related metadata references:</i>	Supersedes Time of first angioplasty balloon inflation or stenting, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005
	<i>Implementation start date:</i> 07/12/2005
	<i>Information specific to this data set:</i>

For Acute coronary syndrome (ACS) reporting, refers to

coronary arteries.

Acute coronary syndrome (clinical) DSS NHIG, Superseded
07/12/2005

Information specific to this data set:

For Acute coronary syndrome (ACS) reporting, refers to
coronary arteries.

Time of intravenous fibrinolytic therapy

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – intravenous fibrinolytic therapy time, hhmm
<i>METeOR identifier:</i>	285201
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The time intravenous (IV) fibrinolytic therapy was first administered to a person.

Data element concept attributes

<i>Data element concept:</i>	Person – intravenous fibrinolytic therapy time
<i>Definition:</i>	The time intravenous (IV) fibrinolytic therapy was first administered.
<i>Context:</i>	Health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Intravenous fibrinolytic therapy time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Time
<i>Data type:</i>	Date/Time
<i>Format:</i>	hhmm
<i>Maximum character length:</i>	4

Source and reference attributes

<i>Reference documents:</i>	ISO 8601:2000 : Data elements and interchange formats - Information interchange - Representation of dates and times
-----------------------------	---

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	If initiated by a bolus dose whether in a pre-hospital setting, emergency department or inpatient unit/ward, the time the initial bolus was administered should be reported.
-----------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	Acute coronary syndrome data working group
<i>Steward:</i>	The National Heart Foundation of Australia and The Cardiac Society of Australia and New Zealand

Relational attributes

<i>Related metadata references:</i>	Supersedes Time of intravenous fibrinolytic therapy, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005 <i>Implementation start date:</i> 07/12/2005

Information specific to this data set:

For Acute coronary syndrome (ACS) reporting, refers to coronary arteries.

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Information specific to this data set:

For Acute coronary syndrome (ACS) reporting, refers to coronary arteries.

Time of triage

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Non-admitted patient emergency department service episode – triage time, hhmm
<i>METeOR identifier:</i>	313817
<i>Registration status:</i>	NHIG, Standard 07/12/2005
<i>Definition:</i>	The time at which the patient is triaged .
<i>Context:</i>	Emergency Department care.

Data element concept attributes

<i>Data element concept:</i>	Non-admitted patient emergency department service episode – triage time
<i>Definition:</i>	The time at which the patient is triaged.
<i>Context:</i>	Emergency Department care.
<i>Object class:</i>	Non-admitted patient emergency department service episode
<i>Property:</i>	Triage time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Time
<i>Data type:</i>	Date/Time
<i>Format:</i>	hhmm
<i>Maximum character length:</i>	4

Source and reference attributes

<i>Reference documents:</i>	ISO 8601:2000 : Data elements and interchange formats - Information interchange - Representation of dates and times
-----------------------------	---

Data element attributes

Collection and usage attributes

<i>Collection methods:</i>	Collected in conjunction with non-admitted patient emergency department service episode – triage date.
----------------------------	---

Source and reference attributes

<i>Submitting organisation:</i>	Australian Government Department of Health and Ageing
---------------------------------	---

Relational attributes

<i>Related metadata references:</i>	Supersedes Triage – triage time, hhmm NHIG, Superseded 07/12/2005
<i>Implementation in Data Set Specifications:</i>	Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005 <i>Implementation start date:</i> 07/12/2005 Non-admitted patient emergency department care NMDS

NHIG, Standard 24/03/2006

Implementation start date: 01/07/2006

Non-admitted patient emergency department care NMDS

NHIG, Superseded 24/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Time patient presents

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Health service event – presentation time, hhmm
<i>METeOR identifier:</i>	270080
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The time at which the patient presents for the delivery of a service.

Data element concept attributes

<i>Data element concept:</i>	Health service event – presentation time
<i>Definition:</i>	The time at which the patient presents for the delivery of a service.
<i>Context:</i>	Admitted patient care. Community health care. Hospital non-admitted patient care: Required to identify commencement of a visit and for calculation of waiting times.
<i>Object class:</i>	Health service event
<i>Property:</i>	Presentation time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Time
<i>Data type:</i>	Date/Time
<i>Format:</i>	hhmm
<i>Maximum character length:</i>	4

Source and reference attributes

<i>Reference documents:</i>	ISO 8601:2000 : Data elements and interchange formats - Information interchange - Representation of dates and times
-----------------------------	---

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>For community health care, outreach services and services provided via telephone or telehealth, this may be the time at which the service provider presents to the patient or the telephone/telehealth session commences.</p> <p>The time of patient presentation at the emergency department is the earliest occasion of being registered clerically or triaged.</p> <p>The time that the patient presents is not necessarily:</p> <ul style="list-style-type: none">• the listing time for care (see listing date for care for an analogous concept), nor• the time at which care is scheduled to be provided, nor• the time at which commencement of care actually occurs
-----------------------	--

(for admitted patients see admission time, for hospital non-admitted patient care and community health care see service commencement time).

Source and reference attributes

Submitting organisation: National Institution Based Ambulatory Model Reference Group
Origin: National Health Data Committee

Relational attributes

Related metadata references: Supersedes Time patient presents, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005
Is used in the formation of Non-admitted patient emergency department service episode – waiting time (to service delivery), total minutes NNNNN NHIG, Standard 01/03/2005
Is used in the formation of Non-admitted patient emergency department service episode – service episode length, total minutes NNNNN NHIG, Standard 01/03/2005
Is used in the formation of Non-admitted patient emergency department service episode – waiting time (to hospital admission), total hours and minutes NNNN NHIG, Standard 01/03/2005

Implementation in Data Set Specifications: Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005
Implementation start date: 07/12/2005
Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005
Non-admitted patient emergency department care NMDS NHIG, Standard 24/03/2006
Implementation start date: 01/07/2006
Non-admitted patient emergency department care NMDS NHIG, Superseded 07/12/2005
Non-admitted patient emergency department care NMDS NHIG, Superseded 24/03/2006
Implementation start date: 01/07/2005
Implementation end date: 30/06/2006

Time troponin measured

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – troponin level measured time, hhmm
<i>METeOR identifier:</i>	285211
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The time at which the troponin (T or I) was measured.

Data element concept attributes

<i>Data element concept:</i>	Person – troponin level measured time
<i>Definition:</i>	The time at which the troponin (T or I) was measured.
<i>Context:</i>	Health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Troponin level measured time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Time
<i>Data type:</i>	Date/Time
<i>Format:</i>	hhmm
<i>Maximum character length:</i>	4

Source and reference attributes

<i>Reference documents:</i>	ISO 8601:2000 : Data elements and interchange formats - Information interchange - Representation of dates and times
-----------------------------	---

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	This metadata item pertains to the measuring of troponin at any time point during this current event.
-----------------------	---

Source and reference attributes

<i>Submitting organisation:</i>	Acute coronary syndrome data working group
<i>Steward:</i>	The National Heart Foundation of Australia and The Cardiac Society of Australia and New Zealand

Relational attributes

<i>Related metadata references:</i>	Supersedes Time troponin measured, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005 <i>Implementation start date:</i> 07/12/2005 Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Tobacco smoking status

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – tobacco smoking status, code N
<i>METeOR identifier:</i>	270311
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	A person's current and past smoking behaviour, as represented by a code.
<i>Context:</i>	Public health and health care

Data element concept attributes

<i>Data element concept:</i>	Person – tobacco smoking status
<i>Definition:</i>	A person's current and past smoking behaviour.
<i>Context:</i>	Public health and health care
<i>Object class:</i>	Person
<i>Property:</i>	Tobacco smoking status

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code												
<i>Data type:</i>	Number												
<i>Format:</i>	N												
<i>Maximum character length:</i>	1												
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Daily smoker</td></tr><tr><td>2</td><td>Weekly smoker</td></tr><tr><td>3</td><td>Irregular smoker</td></tr><tr><td>4</td><td>Ex-smoker</td></tr><tr><td>5</td><td>Never smoked</td></tr></tbody></table>	Value	Meaning	1	Daily smoker	2	Weekly smoker	3	Irregular smoker	4	Ex-smoker	5	Never smoked
Value	Meaning												
1	Daily smoker												
2	Weekly smoker												
3	Irregular smoker												
4	Ex-smoker												
5	Never smoked												

Collection and usage attributes

<i>Guide for use:</i>	CODE 1 Daily smoker A person who smokes daily
	CODE 2 Weekly smoker A person who smokes at least weekly but not daily
	CODE 3 Irregular smoker A person who smokes less than weekly
	CODE 4 Ex-smoker A person who does not smoke at all now, but has smoked at least 100 cigarettes or a similar amount of other tobacco products in his/her lifetime.
	CODE 5 Never-smoker A person who does not smoke now and has smoked fewer than 100 cigarettes or similar amount of other tobacco products in his/her lifetime.

Source and reference attributes

Reference documents: Standard Questions on the Use of Tobacco Among Adults (1998)

Data element attributes

Collection and usage attributes

Collection methods: The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults - interviewer administered (Questions 1 and 4) and self-administered (Questions 1 and 1a) versions. The questionnaires are designed to cover persons aged 18 years and over.

Comments: There are two other ways of categorising this information:

- Regular and irregular smokers where a regular smoker includes someone who is a daily smoker or a weekly smoker. 'Regular' smokers is the preferred category to be reported in prevalence estimates.
- Daily and occasional smokers where an occasional smoker includes someone who is a weekly or irregular smoker. The category of 'occasional' smoker can be used when the aim of the study is to draw contrast between daily smokers and other smokers.

Where this information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

Smoker type is used to define subpopulations of adults (age 18+ years) based on their smoking behaviour.

Smoking has long been known as a health risk factor. Population studies indicate a relationship between smoking and increased mortality/morbidity.

This data element can be used to estimate smoking prevalence. Other uses are:

- To evaluate health promotion and disease prevention programs (assessment of interventions)
- To monitor health risk factors and progress towards National Health Goals and Targets

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references: Supersedes Tobacco smoking status, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Information specific to this data set:

Smoker type is used to define sub-populations of adults (age 18+ years) based on their smoking behaviour.

Smoking has long been known as a health risk factor. Population studies indicate a relationship between smoking and increased mortality/morbidity. This metadata item can be used to estimate smoking prevalence.

Other uses are:

- To evaluate health promotion and disease prevention programs (assessment of interventions)
- To monitor health risk factors and progress towards National Health Goals and Targets

Acute coronary syndrome (clinical) DSS NHIG, Superseded
07/12/2005

Information specific to this data set:

Smoker type is used to define sub-populations of adults (age 18+ years) based on their smoking behaviour.

Smoking has long been known as a health risk factor. Population studies indicate a relationship between smoking and increased mortality/morbidity. This metadata item can be used to estimate smoking prevalence.

Other uses are:

- To evaluate health promotion and disease prevention programs (assessment of interventions)
- To monitor health risk factors and progress towards National Health Goals and Targets

Cardiovascular disease (clinical) DSS NHIG, Superseded
15/02/2006

Cardiovascular disease (clinical) DSS NHIG, Standard
15/02/2006

Tobacco smoking status (diabetes mellitus)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – regular tobacco smoking indicator (last 3 months), code N
<i>METeOR identifier:</i>	302467
<i>Registration status:</i>	NHIG, Standard 21/09/2005
<i>Definition:</i>	Whether an individual has been a regular smoker (daily or weekly) of any tobacco material over the previous 3 months, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – regular tobacco smoking indicator
<i>Definition:</i>	Whether an individual has been a regular smoker (daily or weekly) of any tobacco material.
<i>Context:</i>	Public health, health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Regular tobacco smoking indicator

Collection and usage attributes

<i>Comments:</i>	Smoking is the act of drawing into the mouth and puffing out the smoke of tobacco contained in a cigarette, cigar or pipe. Tobacco smoke contains a number of harmful substances including poisons, various irritant and carcinogenic compounds.
------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	Australian Institute of Health and Welfare
---------------------------------	--

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code								
<i>Data type:</i>	Number								
<i>Format:</i>	N								
<i>Maximum character length:</i>	1								
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Yes</td></tr><tr><td>2</td><td>No</td></tr><tr><td>9</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	1	Yes	2	No	9	Not stated/inadequately described
Value	Meaning								
1	Yes								
2	No								
9	Not stated/inadequately described								
<i>Supplementary values:</i>	9 Not stated/inadequately described								

Collection and usage attributes

<i>Guide for use:</i>	CODE 9 Not stated/inadequately described This code is not for use in primary data collections.
-----------------------	---

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	CODE 1 Yes: Record if the person has smoked daily or weekly over the previous 3 months. CODE 2 No: Record if the person has not smoked daily or weekly over the previous 3 months or has been an irregular smoker.
<i>Collection methods:</i>	Ask the individual if he/she has regularly smoked (daily or weekly) any tobacco material over the past 3 months.

Source and reference attributes

<i>Submitting organisation:</i>	National diabetes data working group
<i>Origin:</i>	National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary.

Relational attributes

<i>Related metadata references:</i>	Supersedes Person – tobacco smoking status (previous three months), code N NHIG, Superseded 21/09/2005
<i>Implementation in Data Set Specifications:</i>	Diabetes (clinical) DSS NHIG, Standard 21/09/2005 <i>Information specific to this data set:</i> For people with diabetes smoking is one of the most powerful treatable risk factors. Associated with hypertension, diabetes and hypercholesterolemia, smoking is a definite health hazard for coronary heart disease.

Tobacco smoking—consumption/quantity (cigarettes)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – number of cigarettes smoked (per day), total N[N]
<i>METeOR identifier:</i>	270332
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The total number of cigarettes (manufactured or roll-your-own) smoked per day by a person.
<i>Context:</i>	Public health and health care

Data element concept attributes

<i>Data element concept:</i>	Person – number of cigarettes smoked
<i>Definition:</i>	The number of cigarettes smoked by a person.
<i>Object class:</i>	Person
<i>Property:</i>	Number of cigarettes smoked

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	Number				
<i>Format:</i>	N[N]				
<i>Maximum character length:</i>	2				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	99	Not stated/inadequately described
Value	Meaning				
99	Not stated/inadequately described				
<i>Unit of measure:</i>	Cigarette				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>This metadata item is relevant only for persons who currently smoke cigarettes daily or at least weekly. Daily consumption should be reported, rather than weekly consumption. Weekly consumption is converted to daily consumption by dividing by 7 and rounding to the nearest whole number.</p> <p>Quantities greater than 98 (extremely rare) should be recorded as 98.</p>
<i>Collection methods:</i>	<p>The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults (1998) - interviewer administered (Questions 3a and 3b) and self-administered (Questions 2a and 2b) versions.</p> <p>The questions cover persons aged 18 and over.</p>
<i>Comments:</i>	<p>The number of cigarettes smoked is an important measure of the magnitude of the tobacco problem for an individual.</p> <p>Research shows that of Australians who smoke, the overwhelming majority smoke cigarettes (manufactured or roll-your-own) rather than other tobacco products.</p>

From a public health point of view, consumption level is relevant only for regular smokers (those who smoke daily or at least weekly).

Data on quantity smoked can be used to:

- evaluate health promotion and disease prevention programs (assessment of interventions)
- monitor health risk factors and progress towards National Health Goals and Targets
- ascertain determinants and consequences of smoking
- assess a person's exposure to tobacco smoke.

Where this information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - consumption/ quantity (cigarettes), version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications:

Cardiovascular disease (clinical) DSS NHIG, Superseded 15/02/2006

Information specific to this data set:

The number of cigarettes smoked is an important measure of the magnitude of the tobacco problem for an individual. Research shows that of Australians who smoke, the overwhelming majority smoke cigarettes (manufactured or roll-your-own) rather than other tobacco products. From a public health point of view, consumption level is relevant only for regular smokers (those who smoke daily or at least weekly).

Data on quantity smoked can be used to:

- evaluate health promotion and disease prevention programs (assessment of interventions)
- monitor health risk factors and progress towards National Health Goals and Targets
- ascertain determinants and consequences of smoking
- assess a person's exposure to tobacco smoke.

Cardiovascular disease (clinical) DSS NHIG, Standard 15/02/2006

Information specific to this data set:

The number of cigarettes smoked is an important measure of the magnitude of the tobacco problem for an individual. Research shows that of Australians who smoke, the overwhelming majority smoke cigarettes (manufactured or

roll-your-own) rather than other tobacco products. From a public health point of view, consumption level is relevant only for regular smokers (those who smoke daily or at least weekly).

Data on quantity smoked can be used to:

- evaluate health promotion and disease prevention programs (assessment of interventions)
- monitor health risk factors and progress towards National Health Goals and Targets
- ascertain determinants and consequences of smoking
- assess a person's exposure to tobacco smoke.

Tobacco smoking—duration (daily smoking)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person—tobacco smoking duration (daily smoking), total years N[N]
<i>METeOR identifier:</i>	270330
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The total duration in years, of daily smoking for a person who is now a daily smoker or has been a daily smoker in the past.
<i>Context:</i>	Public health and health care

Data element concept attributes

<i>Data element concept:</i>	Person—tobacco smoking duration
<i>Definition:</i>	Duration of smoking for a person who is now a smoker or has been a smoker in the past.
<i>Object class:</i>	Person
<i>Property:</i>	Tobacco smoking duration

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	Number				
<i>Format:</i>	N[N]				
<i>Maximum character length:</i>	2				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	99	Not stated/inadequately described
Value	Meaning				
99	Not stated/inadequately described				
<i>Unit of measure:</i>	Year				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	In order to estimate duration of smoking the person's date of birth or current age should also be collected. If a person reports that they smoke daily now then duration is the difference between the start-age and the person's current age. If a person reports that they smoked daily in the past but do not smoke daily now then duration is the difference between the quit age and the start age. Record duration of less than one year as 0.
<i>Collection methods:</i>	The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults (1998) - interviewer administered (Question 1, 5, 6, 7) and self-administered (Question 1, 3, 3a, 4) versions. The questions cover persons aged 18 years and over.
<i>Comments:</i>	Duration of daily smoking is an indicator of exposure to increased risk to health. In this data element, duration is measured as the years elapsed from the time the person first

started smoking daily and when they most recently quit smoking daily (or the present for those persons who still smoke daily). There may have been intervening periods when the person did not smoke daily. However, as the negative health effects of smoking accumulate over time, the information on duration of daily smoking, as measured in this data element, remains useful, despite any intervening periods of non-daily smoking.

Where this information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - duration (daily smoking), version 1, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Is formed using Person – tobacco smoking start age (daily smoking), total years NN NHIG, Standard 01/03/2005

Is formed using Person – tobacco smoking quit age (daily smoking), total years NN NHIG, Standard 01/03/2005

Tobacco smoking—ever daily use

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – tobacco smoking daily use status, code N
<i>METeOR identifier:</i>	270329
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	An indicator of whether a person has ever smoked tobacco in any form on a daily basis in their lifetime, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – tobacco smoking daily use status
<i>Definition:</i>	Whether a person has ever smoked tobacco in any form daily in his or her lifetime.
<i>Context:</i>	Public health and health care
<i>Object class:</i>	Person
<i>Property:</i>	Tobacco smoking daily use status

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Ever-daily</td></tr><tr><td>2</td><td>Never-daily</td></tr></tbody></table>	Value	Meaning	1	Ever-daily	2	Never-daily
Value	Meaning						
1	Ever-daily						
2	Never-daily						

Collection and usage attributes

<i>Guide for use:</i>	<p>CODE 1 Ever-daily</p> <p>If a person reports that they now smoke cigarettes, cigars, pipes or any other tobacco products daily OR if they report that in the past they have been a daily smoker, they are coded to 1 (ever-daily).</p> <p>CODE 2 Never-daily</p> <p>If a person reports that they have never smoked cigarettes, cigars, pipes or any other tobacco products daily AND they have never in the past been a daily smoker then they are coded to 2 (never-daily).</p>
-----------------------	--

Data element attributes

Collection and usage attributes

<i>Collection methods:</i>	The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults (1998) - interviewer administered (Question 1 and 5) and self-
----------------------------	--

Comments:

administered (Question 1 and 3) versions. The questions cover persons aged 18 years and over.

Whether a person has ever smoked on a daily basis can be used to assess an individual's health risk from smoking and to monitor population trends in smoking behaviour.

It can also be used to:

- evaluate health promotion and disease prevention programs (assessment of interventions);
- monitor health risk factors;
- ascertain determinants and consequences of smoking.

Where the information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - ever daily use, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Tobacco smoking—frequency

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – tobacco smoking frequency, code N
<i>METeOR identifier:</i>	270328
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	How often a person now smokes a tobacco product, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – tobacco smoking frequency
<i>Definition:</i>	How often a person now smokes a tobacco product.
<i>Context:</i>	Public health and health care
<i>Object class:</i>	Person
<i>Property:</i>	Tobacco smoking frequency

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code										
<i>Data type:</i>	Number										
<i>Format:</i>	N										
<i>Maximum character length:</i>	1										
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Smokes daily</td></tr><tr><td>2</td><td>Smokes at least weekly, but not daily</td></tr><tr><td>3</td><td>Smokes less often than weekly</td></tr><tr><td>4</td><td>Does not smoke at all</td></tr></tbody></table>	Value	Meaning	1	Smokes daily	2	Smokes at least weekly, but not daily	3	Smokes less often than weekly	4	Does not smoke at all
Value	Meaning										
1	Smokes daily										
2	Smokes at least weekly, but not daily										
3	Smokes less often than weekly										
4	Does not smoke at all										

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	To record multiple use data, repeat the data field as many times as necessary, viz: product1, product2 etc. In most instances, data on both product and frequency are needed. In such situations, repeat both fields as many times as necessary, viz: product1, frequency1, product2, frequency2 etc.
<i>Collection methods:</i>	The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults (1998) - interviewer administered (Question 1) and self-administered (Question 1) versions. The questions relate to smoking of manufactured cigarettes, roll-your-own cigarettes, cigars, pipes and other tobacco products and are designed to cover persons aged 18 years and over.
<i>Comments:</i>	The frequency of smoking helps to assess a person's exposure to tobacco smoke which is a known risk factor for cardiovascular disease and cancer. From a public health point of view, the level

of consumption of tobacco as measured by frequency of smoking tobacco products is only relevant for regular smokers (persons who smoke daily or at least weekly).

Where this information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - frequency, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Tobacco smoking—product

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – tobacco product smoked, code N
<i>METeOR identifier:</i>	270327
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The type of tobacco product smoked by a person, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – tobacco product smoked
<i>Definition:</i>	The type of tobacco product smoked by a person.
<i>Context:</i>	Public health and health care
<i>Object class:</i>	Person
<i>Property:</i>	Tobacco product smoked

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code														
<i>Data type:</i>	Number														
<i>Format:</i>	N														
<i>Maximum character length:</i>	1														
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Cigarettes - manufactured</td></tr><tr><td>2</td><td>Cigarettes - roll-your-own</td></tr><tr><td>3</td><td>Cigars</td></tr><tr><td>4</td><td>Pipes</td></tr><tr><td>5</td><td>Other tobacco product</td></tr><tr><td>6</td><td>None</td></tr></tbody></table>	Value	Meaning	1	Cigarettes - manufactured	2	Cigarettes - roll-your-own	3	Cigars	4	Pipes	5	Other tobacco product	6	None
Value	Meaning														
1	Cigarettes - manufactured														
2	Cigarettes - roll-your-own														
3	Cigars														
4	Pipes														
5	Other tobacco product														
6	None														

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	To record multiple use data, repeat the data field as many times as necessary, viz: product1, product2 etc. In most instances, data on both product and frequency are needed. In such situations, repeat both fields as many times as necessary, viz: product1, frequency1, product2, frequency2 etc.
<i>Collection methods:</i>	The recommended standard for collecting information about smoking the above tobacco products is the Standard Questions on the Use of Tobacco Among Adults (1998) - interviewer or self-administered versions.
<i>Comments:</i>	Tobacco smoking is a known risk factor for cardiovascular disease and cancer. The type of tobacco product smoked by a person in conjunction with information about the frequency of

smoking assists with establishing a profile of smoking behaviour at the individual or population level and with monitoring shifts from cigarette smoking to other types of tobacco products and vice versa.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - product, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Tobacco smoking—quit age (daily smoking)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person—tobacco smoking quit age (daily smoking), total years NN
<i>METeOR identifier:</i>	270323
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The age in years at which a person who has smoked daily in the past and is no longer a daily smoker most recently stopped smoking daily.
<i>Context:</i>	Public health and health care

Data element concept attributes

<i>Data element concept:</i>	Person—tobacco smoking quit age
<i>Definition:</i>	The age at which a person stopped smoking.
<i>Object class:</i>	Person
<i>Property:</i>	Tobacco smoking quit age

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	String				
<i>Format:</i>	NN				
<i>Maximum character length:</i>	2				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	99	Not stated/inadequately described
Value	Meaning				
99	Not stated/inadequately described				
<i>Unit of measure:</i>	Year				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>In order to estimate quit-age, the person's date of birth or current age should also be collected. Quit-age may be directly reported, or derived from the date the person quit smoking or the length of time since quitting, once the person's date of birth (or current age) is known.</p> <p>Quit-age is relevant only to persons who have been daily smokers in the past and are not current daily smokers.</p>
<i>Collection methods:</i>	<p>The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults - interviewer administered (Question 6) and self-administered (Question 3a) versions. The questions cover persons aged 18 years and over.</p> <p>The relevant question in each version of the questionnaires refers to when the person finally stopped smoking daily, whereas the definition for this metadata item refers to when the</p>

person most recently stopped smoking daily. However, in order to provide information on when the person most recently stopped smoking daily, the most appropriate question to ask at the time of collecting the information is when the person finally stopped smoking daily.

Comments:

Quit-age and start-age provide information on the duration of daily smoking and exposure to increased risk to health.

Where the information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - quit age (daily smoking), version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Is used in the formation of Person – time since quitting tobacco smoking (daily smoking), code NN NHIG, Standard 01/03/2005

Is used in the formation of Person – tobacco smoking duration (daily smoking), total years N[N] NHIG, Standard 01/03/2005

Tobacco smoking—start age (daily smoking)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person—tobacco smoking start age (daily smoking), total years NN
<i>METeOR identifier:</i>	270324
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The age in years at which a person who has ever been a daily smoker, first started to smoke daily.
<i>Context:</i>	Public health and health care

Data element concept attributes

<i>Data element concept:</i>	Person—tobacco smoking start age
<i>Definition:</i>	The age at which a person first started to smoke.
<i>Object class:</i>	Person
<i>Property:</i>	Tobacco smoking start age

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	String				
<i>Format:</i>	NN				
<i>Maximum character length:</i>	2				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	99	Not stated/inadequately described
Value	Meaning				
99	Not stated/inadequately described				
<i>Unit of measure:</i>	Year				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record age in completed years. This information is relevant only if a person currently smokes daily or has smoked daily in the past.
<i>Collection methods:</i>	The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults (1998) - interviewer administered (Question 7) and self-administered (Question 4) versions. The questions cover persons aged 18 years and over.
<i>Comments:</i>	Start-age may be used to derive duration of smoking, which is a much stronger predictor of the risks associated with smoking than is the total amount of tobacco smoked over time. Where the information is collected by survey and the sample permits, population estimates should be presented by sex and age groups. The recommended age groups are: It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is

also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - start age (daily smoking), version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Is used in the formation of Person – tobacco smoking duration (daily smoking), total years N[N] NHIG, Standard 01/03/2005

Tobacco smoking—time since quitting (daily smoking)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – time since quitting tobacco smoking (daily smoking), code NN
<i>METeOR identifier:</i>	270356
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The time since a person most recently quit daily smoking, as represented by a code.
<i>Context:</i>	Public health and health care

Data element concept attributes

<i>Data element concept:</i>	Person – time since quitting tobacco smoking
<i>Definition:</i>	Time since a person most recently quit smoking.
<i>Object class:</i>	Person
<i>Property:</i>	Time since quitting tobacco smoking

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																																						
<i>Data type:</i>	String																																						
<i>Format:</i>	NN																																						
<i>Maximum character length:</i>	2																																						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>12 months (1 year)</td></tr><tr><td>02</td><td>2 years etc. to 78</td></tr><tr><td>79</td><td>79+ years</td></tr><tr><td>80</td><td>Less than 1 month</td></tr><tr><td>81</td><td>1 month</td></tr><tr><td>82</td><td>2 months</td></tr><tr><td>83</td><td>3 months</td></tr><tr><td>84</td><td>4 months</td></tr><tr><td>85</td><td>5 months</td></tr><tr><td>86</td><td>6 months</td></tr><tr><td>87</td><td>7 months</td></tr><tr><td>88</td><td>8 months</td></tr><tr><td>89</td><td>9 months</td></tr><tr><td>90</td><td>10 months</td></tr><tr><td>91</td><td>11 months</td></tr><tr><td>92</td><td>months, not specified</td></tr><tr><td>93</td><td>years, not specified</td></tr><tr><td>99</td><td>not stated</td></tr></tbody></table>	Value	Meaning	01	12 months (1 year)	02	2 years etc. to 78	79	79+ years	80	Less than 1 month	81	1 month	82	2 months	83	3 months	84	4 months	85	5 months	86	6 months	87	7 months	88	8 months	89	9 months	90	10 months	91	11 months	92	months, not specified	93	years, not specified	99	not stated
Value	Meaning																																						
01	12 months (1 year)																																						
02	2 years etc. to 78																																						
79	79+ years																																						
80	Less than 1 month																																						
81	1 month																																						
82	2 months																																						
83	3 months																																						
84	4 months																																						
85	5 months																																						
86	6 months																																						
87	7 months																																						
88	8 months																																						
89	9 months																																						
90	10 months																																						
91	11 months																																						
92	months, not specified																																						
93	years, not specified																																						
99	not stated																																						
<i>Supplementary values:</i>	99 not stated																																						

Data element attributes

Collection and usage attributes

Guide for use:

In order to estimate time since quitting for all respondents, the person's date of birth or current age should also be collected.

For optimal flexibility of use, the time since quitting is coded as months or years. However, people may report the time that they quit smoking in various ways (e.g. age, a date, or a number of days or weeks ago). When the information is reported in weeks and is less than 4, or in days and is less than 28, then code 80.

When the person reports the time since quitting as weeks ago, convert into months by dividing by 4 (rounded down to the nearest month).

If days reported are between 28 and 59, then code 81.

Where the information is about age only, time since quitting (daily use) is the difference between quit-age and age at survey.

Collection methods:

The recommended standard for collecting this information is the Standard Questions on the Use of Tobacco Among Adults (1998) - interviewer administered (Question 6) and self-administered (Question 3) versions.

Comments:

Time since quitting daily smoking may give an indication of improvement in the health risk profile of a person. It is also useful in evaluating health promotion campaigns.

Where this information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected. It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Relational attributes

Related metadata references:

Supersedes Tobacco smoking - time since quitting (daily smoking), version 1, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Is formed using Person – tobacco smoking quit age (daily smoking), total years NN NHIG, Standard 01/03/2005

Total contract patient days

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Admitted patient hospital stay – number of patient days (of contracted care), total N[NN]
<i>METeOR identifier:</i>	270301
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Sum of the number of contract patient days for all periods within the hospital stay.

Data element concept attributes

<i>Data element concept:</i>	Admitted patient hospital stay – number of patient days (of contracted care)
<i>Definition:</i>	Number of contract patient days (contracted care completion date minus contracted care commencement date) for all periods within the hospital stay.
<i>Context:</i>	Admitted patient care.
<i>Object class:</i>	Admitted patient hospital stay
<i>Property:</i>	Number of patient days

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NN]
<i>Maximum character length:</i>	3
<i>Unit of measure:</i>	Day

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>Count number of days.</p> <p>A day is measured from midnight to 2359 hours.</p> <p>Contract patient days are included in the total count of patient days. If necessary, contract patient days can be distinguished from other patient days by using the following rules:</p> <ul style="list-style-type: none">• The day the contract commences is counted as a contract patient day.• If the patient is on contract from midnight to 2359 count as a contract patient day.• The day a contract is completed is not counted as a contract patient day.• If the patient is admitted and commences a contract on the same day, this is not counted as a contract patient day.• If a contract is completed and the patient is separated on the same day, the day should not be counted as a contract
-----------------------	--

or other patient day.

Relational attributes

Related metadata references:

Supersedes Total contract patient days, version 1, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Total hours worked by a medical practitioner

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Medical practitioner – hours worked, total NNN
<i>METeOR identifier:</i>	270136
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The total hours worked in a week in a job by a medical practitioner, including any on-call hours actually worked (includes patient care and administration).
<i>Context:</i>	Health labour force

Data element concept attributes

<i>Data element concept:</i>	Medical practitioner – hours worked
<i>Definition:</i>	The number of hours worked by a medical practitioner.
<i>Object class:</i>	Medical practitioner
<i>Property:</i>	Hours worked

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	String				
<i>Format:</i>	NNN				
<i>Maximum character length:</i>	3				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>999</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	999	Not stated/inadequately described
Value	Meaning				
999	Not stated/inadequately described				
<i>Unit of measure:</i>	Hour (h)				

Collection and usage attributes

<i>Guide for use:</i>	Total hours expressed as 000, 001 etc.
-----------------------	--

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	This metadata item relates to each position (job) held by a medical practitioner, not the aggregate of hours worked in all.
<i>Collection methods:</i>	There are inherent problems in asking for information on number of hours usually worked per week, for example, reaching a satisfactory definition and communicating this definition to the respondents in a self-administered survey. Whether hours worked are collected for main job only, or main job and one or more additional jobs, it is important that a total for all jobs is included.
<i>Comments:</i>	It is often argued that health professionals contribute a considerable amount of time to voluntary professional work and that this component needs to be identified. This should be considered as an additional item, and kept segregated from

data on paid hours worked.

Source and reference attributes

Submitting organisation: National Health Labour Force Data Working Group

Relational attributes

Related metadata references: Supersedes Total hours worked by a medical practitioner, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Health labour force NMDS NHIG, Standard 01/03/2005

Implementation start date: 01/07/2006

Information specific to this data set:

Value must be less than 169 (except for 999).

Used in relation to issues of economic activity, productivity, wage rates, working conditions etc. Used to develop capacity measures relating to total time available. Assists in analysis of human resource requirements and labour force modelling. Used to determine full-time and part-time work status and to compute full-time equivalents (FTE) (see entry for FTE). Often the definition for full-time or FTE differs (35, 37.5 and 40 hours) and knowing total hours and numbers of individuals allows for variances in FTE.

Total leave days

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of admitted patient care – number of leave days, total N[NN]
<i>METeOR identifier:</i>	270251
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Sum of the length of leave (date returned from leave minus date went on leave) for all periods within the hospital stay.

Data element concept attributes

<i>Data element concept:</i>	Episode of admitted patient care – number of leave days
<i>Definition:</i>	Sum of the length of leave (date returned from leave minus date went on leave) for all periods within the hospital stay.
<i>Context:</i>	Recording of leave days allows for exclusion of these from the calculation of patient days. This is important for analysis of costs per patient and for planning. The maximum limit allowed for leave affects admission and separation rates, particularly for long-stay patients who may have several leave periods.
<i>Object class:</i>	Episode of admitted patient care
<i>Property:</i>	Number of leave days

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NN]
<i>Maximum character length:</i>	3
<i>Unit of measure:</i>	Day

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>A day is measured from midnight to midnight.</p> <p>The following rules apply in the calculation of leave days for both overnight and same-day patients:</p> <ul style="list-style-type: none">• The day the patient goes on leave is counted as a leave day.• The day the patient is on leave is counted as a leave day.• The day the patient returns from leave is counted as a patient day.• If the patient is admitted and goes on leave on the same day, this is counted as a patient day, not a leave day.• If the patient returns from leave and then goes on leave again on the same day, this is counted as a leave day.• If the patient returns from leave and is separated on the same day, the day should not be counted as either a patient
-----------------------	---

day or a leave day.

Comments:

It should be noted that for private patients in public and private hospitals, s.3 (12) of the Health Insurance Act 1973 (Cwlth) currently applies a different leave day count, Commonwealth Department of Human Services and Health HBF Circular 354 (31 March 1994). This metadata item was modified in July 1996 to exclude the previous differentiation between the psychiatric and other patients.

Source and reference attributes

Origin:

National Health Data Committee

Relational attributes

Related metadata references:

Supersedes Total leave days, version 3, DE, NHDD, NHIMG, Superseded 01/03/2005

Is used in the formation of Episode of admitted patient care – length of stay (excluding leave days), total N[NN] NHIG, Standard 01/03/2005

Is used in the formation of Episode of care – number of psychiatric care days, total N[NNNN] NHIG, Standard 01/03/2005

Is used in the formation of Episode of admitted patient care – major diagnostic category, code (AR-DRG v5.1) NN NHIG, Standard 01/03/2005

Is used in the formation of Episode of admitted patient care – diagnosis related group, code (AR-DRG v5.1) ANNA NHIG, Standard 01/03/2005

Implementation in Data Set Specifications:

Admitted patient care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies:

(Episode of admitted patient care – separation date, DDMMYYYY minus Episode of admitted patient care – admission date, DDMMYYYY) minus Admitted patient hospital stay – number of leave days, total N[NN] must be ≥ 0 days.

Admitted patient care NMDS 2006-2007 NHIG, Superseded 29/11/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies:

(Episode of admitted patient care – separation date, DDMMYYYY minus Episode of admitted patient care – admission date, DDMMYYYY) minus Admitted patient hospital stay – number of leave days, total N[NN] must be ≥ 0 days.

Admitted patient care NMDS 2007-2008 NHIG, Standard

29/11/2006

Implementation start date: 01/07/2006

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies:

(Episode of admitted patient care – separation date, DDMMYYYY minus Episode of admitted patient care – admission date, DDMMYYYY) minus Admitted patient hospital stay – number of leave days, total N[NN] must be ≥ 0 days.

Admitted patient mental health care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies:

(Episode of admitted patient care – separation date, DDMMYYYY minus Episode of admitted patient care – admission date, DDMMYYYY) minus Admitted patient hospital stay – number of leave days, total N[NN] must be ≥ 0 days.

Admitted patient mental health care NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies:

(Episode of admitted patient care – separation date, DDMMYYYY minus Episode of admitted patient care – admission date, DDMMYYYY) minus Admitted patient hospital stay – number of leave days, total N[NN] must be ≥ 0 days.

Admitted patient mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies:

(Episode of admitted patient care – separation date, DDMMYYYY minus Episode of admitted patient care – admission date, DDMMYYYY) minus Admitted patient hospital stay – number of leave days, total N[NN] must be ≥ 0 days.

Total number of group sessions

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment – number of group sessions, total N[NNNNN]
<i>METeOR identifier:</i>	292789
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The total number of groups of patients/clients receiving services. Each group is to count once, irrespective of size or the number of staff providing services.

Data element concept attributes

<i>Data element concept:</i>	Establishment – number of group sessions
<i>Definition:</i>	The number of groups of patients/clients receiving services. Each group is to count once, irrespective of size or the number of staff providing services.
<i>Context:</i>	The resources required to provide services to groups of patients are different from those required to provide services to an equivalent number of individuals. Hence services to groups of non-admitted patients or outreach clients should be counted separately from services to individuals.
<i>Object class:</i>	Establishment
<i>Property:</i>	Number of group sessions

Source and reference attributes

<i>Submitting organisation:</i>	National minimum data set working parties.
---------------------------------	--

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NNNNN]
<i>Maximum character length:</i>	6
<i>Unit of measure:</i>	Group session

Data element attributes

Collection and usage attributes

<i>Collection methods:</i>	At present, occasions of service to groups are counted in an inconsistent manner. The numbers of occasions of service should be collected for both individual and group sessions for public psychiatric hospitals and alcohol and drug hospitals.
<i>Comments:</i>	This data element is derived from data elements that are not currently specified in the National Health Data Dictionary, but which are recorded in various ways by hospitals and/or outpatient departments. Examples include identifiers of individual consultations/visits, diagnostic tests, etc.

Source and reference attributes

Submitting organisation: National minimum data set working parties.

Relational attributes

Related metadata references: Supersedes Group sessions, version 1, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Outpatient care NMDS NHIG, Standard 04/05/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Total occasions of service

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Establishment – number of occasions of service, total N[NNNNNN]
<i>METeOR identifier:</i>	291066
<i>Registration status:</i>	NHIG, Standard 04/05/2005
<i>Definition:</i>	The total number of occasions of examination, consultation, treatment or other service provided to a patient.

Data element concept attributes

<i>Data element concept:</i>	Establishment – number of occasions of service
<i>Definition:</i>	The number of occasions of examination, consultation, treatment or other service provided to a patient.
<i>Context:</i>	Occasions of service are required as a measure of non-admitted patient service provision.
<i>Object class:</i>	Establishment
<i>Property:</i>	Number of occasions of service

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NNNNNN]
<i>Maximum character length:</i>	7
<i>Unit of measure:</i>	Occasion of service

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	May apply to imaging, pathology and/or pharmacy services, according to relevant National Minimum Data Set (NMDS) specifications. Each diagnostic test or simultaneous set of related diagnostic tests for the one patient referred to a hospital pathology department consists of one occasion of service.
<i>Collection methods:</i>	This data element must be collected in conjunction with outpatient clinic type. Data compiled from service or clinic activity records. Outpatient Care NMDS the following guides for use also apply: <ul style="list-style-type: none">• Patients participating in group sessions each receive a count of one occasion of service, and these counts are assigned separately.• Outreach clinic services should be counted at the facility where the patient medical record is held.• Where a patient receives multidisciplinary care, within one booked appointment, one occasion of service shall be

- recorded, regardless of the number of providers involved.
- Where patients have more than one booked appointment, each appointment will be counted as one occasion of service. (Example: three booked appointments on a single day will be counted as three OOS).
- The occasion of service count should be attributed to the clinic type associated with the booked appointment.
- Not counted are telephone calls and telemedicine contacts (as listed in the data element Non-admitted patient service event – service mode, hospital code N{.N})

Comments: Replaces Occasions of Service version 1, generalising its applicability to encompass the Outpatient Care NMDS, commencing 1 July 2005.

Source and reference attributes

Submitting organisation: National minimum data set working parties, including Outpatient NMDS Working Group 2004.

Relational attributes

Related metadata references: See also Establishment – outpatient clinic type, code N[N] NHIG, Standard 04/05/2005

Implementation in Data Set Specifications: Outpatient care NMDS NHIG, Standard 04/05/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Total psychiatric care days

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of care – number of psychiatric care days, total N[NNNN]
<i>METeOR identifier:</i>	270300
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The sum of the number of days or part days of stay that the person received care as an admitted patient or resident within a designated psychiatric unit, minus the sum of leave days occurring during the stay within the designated unit.

Data element concept attributes

<i>Data element concept:</i>	Episode of care – number of psychiatric care days
<i>Definition:</i>	The sum of the number of days or part days of stay that the person received care as an admitted patient or resident within a designated psychiatric unit, minus the sum of leave days occurring during the stay within the designated unit.
<i>Context:</i>	Admitted patient and residential mental health care: This metadata item is required to identify the characteristics of patients treated in specialist psychiatric units located within acute care hospitals or 24-hour staffed community-based residential services and to analyse the activities of these units and services. Community mental health care: This metadata item is required to identify the characteristics of patients treated in specialist psychiatric 24-hour staffed community-based residential services and to analyse the activities of these units. The metadata item is necessary to describe and evaluate the progress of mainstreaming of mental health services.
<i>Object class:</i>	Episode of care
<i>Property:</i>	Number of psychiatric care days

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NNNN]
<i>Maximum character length:</i>	5
<i>Unit of measure:</i>	Day

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Designated psychiatric units are staffed by health professionals with specialist mental health qualifications or training and have
-----------------------	--

as their principal function the treatment and care of patients affected by mental disorder. The unit may or may not be recognised under relevant State and Territory legislation to treat patients on an involuntary basis. Patients are admitted patients in the acute and psychiatric hospitals and residents in community based residences.

Public acute care hospitals:

Designated psychiatric units in public acute care hospitals are normally recognised by the State/Territory health authority in the funding arrangements applying to those hospitals.

Private acute care hospitals:

Designated psychiatric units in private acute care hospitals normally require license or approval by the State/Territory health authority in order to receive benefits from health funds for the provision of psychiatric care.

Psychiatric hospitals:

Total psychiatric care days in stand-alone psychiatric hospitals are calculated by counting those days the patient received specialist psychiatric care. Leave days and days on which the patient was receiving other care (e.g. specialised intellectual ability or drug and alcohol care) should be excluded.

Psychiatric hospitals are establishments devoted primarily to the treatment and care of admitted patients with psychiatric, mental or behavioural disorders. Private hospitals formerly approved by the Commonwealth Department of Health under the Health Insurance Act 1973 (Commonwealth) (now licensed/approved by each State/Territory health authority), catering primarily for patients with psychiatric or behavioural disorders are included in this category.

Community-based residential services:

Designated psychiatric units refers to 24-hour staffed community-based residential units established in community settings that provide specialised treatment, rehabilitation or care for people affected by a mental illness or psychiatric disability. Special psychiatric units for the elderly are covered by this category, including psychogeriatric hostels or psychogeriatric nursing homes. Note that residences occupied by admitted patients located on hospital grounds, whether on the campus of a general or stand-alone psychiatric hospital, should be counted in the category of admitted patient services and not as community-based residential services.

Counting of patient days and leave days in designated psychiatric units should follow the standard definitions applying to these items.

For each period of care in a designated psychiatric unit, total days is calculated by subtracting the date on which care commenced within the unit from the date on which the specialist unit care was completed, less any leave days that occurred during the period.

Total psychiatric care days in 24-hour community-based residential care are calculated by counting those days the patient received specialist psychiatric care. Leave days and days on which the patient was receiving other care (e.g. specialised intellectual ability or drug and alcohol care) should be excluded.

Admitted patients in acute care:

Commencement of care within a designated psychiatric unit may be the same as the date the patient was admitted to the hospital, or occur subsequently, following transfer of the patient from another hospital ward. Where commencement of psychiatric care occurs by transfer from another ward, a new episode of care may be recorded, depending on whether the care type has changed (see metadata item Care type). Completion of care within a designated psychiatric unit may be the same as the date the patient was discharged from the hospital, or occur prior to this on transfer of the patient to another hospital ward. Where completion of psychiatric care is followed by transfer to another hospital ward, a new episode of care may be recorded, depending on whether the care type has changed (see metadata item Care type). Total psychiatric care days may cover one or more periods in a designated psychiatric unit within the overall hospital stay.

Collection methods:

Accurate counting of total days in psychiatric care requires periods in designated psychiatric units to be identified in the person-level data collected by state or territory health authorities. Several mechanisms exist for this data field to be implemented:

- Ideally, the new data field should be collected locally by hospitals and added to the unit record data provided to the relevant state/territory health authority.
- Acute care hospitals in most states and territories include details of the wards in which the patient was accommodated in the unit record data provided to the health authority. Local knowledge should be used to identify designated psychiatric units within each hospital's ward codes, to allow total psychiatric care days to be calculated for each episode of care.
- Acute care hospitals and 24-hour staffed community-based residential services should be identified separately at the level of the establishment.

Comments:

This metadata item was originally designed to monitor trends in the delivery of psychiatric admitted patient care in acute care hospitals. It has been modified to enable collection of data in the community-based residential care sector. The metadata item is intended to improve understanding in this area and contribute to the ongoing evaluation of changes occurring in mental health services.

Source and reference attributes

Submitting organisation:

National Mental Health Information Strategy Committee

Reference documents:

Health Insurance Act 1973 (Commonwealth)

Relational attributes

Related metadata references:

Supersedes Total psychiatric care days, version 2, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Is formed using Establishment – establishment type, sector and services provided code AN.N{.N} NHIG, Standard 01/03/2005

Is formed using Hospital service – care type, code N[N].N NHIG, Standard 01/03/2005

Is formed using Episode of admitted patient care – number of leave days, total N[NN] NHIG, Standard 01/03/2005

Implementation in Data Set Specifications:

Is formed using Episode of admitted patient care – admission date, DDMMYYYY NHIG, Standard 01/03/2005

Is formed using Episode of admitted patient care – separation date, DDMMYYYY NHIG, Standard 01/03/2005

Admitted patient care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

Total days in psychiatric care must be: \geq zero; and \leq length of stay.

Admitted patient care NMDS 2007-2008 NHIG, Standard 29/11/2006

Implementation start date: 01/07/2007

Information specific to this data set:

Total days in psychiatric care must be: \geq zero; and \leq length of stay.

Admitted patient mental health care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

Total days in psychiatric care must be \geq zero;

Total days in psychiatric care must be \leq length of stay.

Admitted patient mental health care NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Information specific to this data set:

Total days in psychiatric care must be \geq zero;

Total days in psychiatric care must be \leq length of stay.

Admitted patient mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Information specific to this data set:

Total days in psychiatric care must be \geq zero;

Total days in psychiatric care must be \leq length of stay.

Treatment delivery setting for alcohol and other drugs

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of treatment for alcohol and other drugs – service delivery setting, code N
<i>METeOR identifier:</i>	270068
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The main physical setting in which the type of treatment that is the principal focus of a client's alcohol and other drug treatment episode is actually delivered irrespective of whether or not this is the same as the usual location of the service provider, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Episode of treatment for alcohol and other drugs – service delivery setting
<i>Definition:</i>	The main physical setting in which the type of treatment that is the principal focus of a client's alcohol and other drug treatment episode is actually delivered, irrespective of whether or not this is the same as the usual location of the service provider.
<i>Context:</i>	Alcohol and other drug treatment services. Required to identify the settings in which treatment is occurring, allowing for trends in treatment patterns to be monitored.
<i>Object class:</i>	Episode of treatment for alcohol and other drugs
<i>Property:</i>	Service delivery setting

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code												
<i>Data type:</i>	Number												
<i>Format:</i>	N												
<i>Maximum character length:</i>	1												
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Non-residential treatment facility</td></tr><tr><td>2</td><td>Residential treatment facility</td></tr><tr><td>3</td><td>Home</td></tr><tr><td>4</td><td>Outreach setting</td></tr><tr><td>8</td><td>Other</td></tr></tbody></table>	Value	Meaning	1	Non-residential treatment facility	2	Residential treatment facility	3	Home	4	Outreach setting	8	Other
Value	Meaning												
1	Non-residential treatment facility												
2	Residential treatment facility												
3	Home												
4	Outreach setting												
8	Other												

Collection and usage attributes

<i>Guide for use:</i>	Only one code to be selected at the end of the alcohol and other drug treatment episode. Agencies should report the setting in which most of the main type of treatment was received by the client during the treatment episode. CODE 1 Non-residential treatment facility This code refers to any non-residential centre that provides
-----------------------	---

alcohol and other drug treatment services, including hospital outpatient services and community health centres.

CODE 2 Residential treatment facility

This code refers to community-based settings in which clients reside either temporarily or long-term in a facility that is not their home or usual place of residence to receive alcohol and other drug treatment. This does not include ambulatory situations, but does include therapeutic community settings.

CODE 3 Home

This code refers to the client's own home or usual place of residence.

CODE 4 Outreach setting

This code refers to an outreach environment, excluding a client's home or usual place of residence, where treatment is provided. An outreach environment may be any public or private location that is not covered by Codes 1-3.

Mobile/outreach alcohol and other drug treatment service providers would usually provide treatment within this setting.

Data element attributes

Source and reference attributes

Submitting organisation: Intergovernmental Committee on Drugs National Minimum Data Set Working Group

Relational attributes

Related metadata references: Supersedes Treatment delivery setting for alcohol and other drugs, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Alcohol and other drug treatment services NMDS NHIG, Superseded 21/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Alcohol and other drug treatment services NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Alcohol and other drug treatment services NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Triglyceride level (measured)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – triglyceride level (measured), total millimoles per litre N[N].N
<i>METeOR identifier:</i>	270229
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	A person's triglyceride level measured in millimoles per litre.

Data element concept attributes

<i>Data element concept:</i>	Person – triglyceride level
<i>Definition:</i>	A person's triglyceride level.
<i>Context:</i>	Public health, health care and clinical setting.
<i>Object class:</i>	Person
<i>Property:</i>	Triglyceride level

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	Number				
<i>Format:</i>	N[N].N				
<i>Maximum character length:</i>	3				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>99.9</td><td>Not stated/inadequately described.</td></tr></tbody></table>	Value	Meaning	99.9	Not stated/inadequately described.
Value	Meaning				
99.9	Not stated/inadequately described.				
<i>Unit of measure:</i>	Millimole per litre (mmol/L)				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record the absolute result of the total triglyceride measurement.
<i>Collection methods:</i>	<p>Measurement of lipid levels should be carried out by laboratories, or practices, which have been accredited to perform these tests by the National Association of Testing Authorities.</p> <ul style="list-style-type: none">To be collected as a single venous blood sample, preferably following a 12-hour fast where only water and medications have been consumed. <p>Note that to calculate the low-density lipoprotein - cholesterol (LDL-C) from the Friedwald Equation (Friedwald et al, 1972):</p> <ul style="list-style-type: none">a fasting level of plasma triglyceride and knowledge of the levels of plasma total cholesterol and high-density lipoprotein - cholesterol (HDL-C) is required,the Friedwald equation becomes unreliable when the plasma triglyceride exceeds 4.5 mmol/L, andthat while levels are reliable for the first 24 hours after the

onset of acute coronary syndromes, they may be unreliable for the subsequent 6 weeks after an event.

Source and reference attributes

Submitting organisation: Cardiovascular Data Working Group

Relational attributes

Related metadata references: Supersedes Triglycerides - measured, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
Is used in the formation of Person – low-density lipoprotein cholesterol level (calculated), total millimoles per litre N[N].N NHIG, Standard 01/03/2005

Implementation in Data Set Specifications: Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Cardiovascular disease (clinical) DSS NHIG, Superseded 15/02/2006

Information specific to this data set:

A relationship between triglyceride and High-density Lipoprotein Cholesterol (HDL-C) and chronic heart disease (CHD) event rates has been shown. This view is supported by the observation that the remnants of triglyceride-rich lipoproteins are the particles that occur in dysbetalipoproteinaemia, a condition associated with a very high risk of premature atherosclerotic vascular disease. There have been two comprehensive reviews of the relationship between plasma triglyceride and CHD (see Criqui et al. 1993 and Austin et al. 1991). Criqui concludes that triglyceride is not an independent predictor of CHD and is probably not causally related to the disease, while Austin provides a compelling case for a causal role of (at least) some triglyceride rich lipoproteins. Conclusions drawn from population studies of the relationship between plasma triglyceride and the risk of CHD include the following:

- an elevated concentration of plasma triglyceride (> 2.0 mmol/L) is predictive of CHD when associated with either an increased concentration of LDL-C or a decreased concentration of HDL-C.
- the relationship between CHD risk and plasma triglyceride is not continuous, with evidence that the risk is greatest in people with triglyceride levels between 2 and 6 mmol/L (Lipid Management Guidelines - 2001, MJA 2001; 175: S57-S88. National Heart Foundation of Australia and the Cardiac Society of Australia and New Zealand).

It is likely that the positive relationship between plasma triglyceride and CHD, as observed in many population studies, is because an elevated level of plasma triglyceride in some people is a reflection of an accumulation of the atherogenic remnants of chylomicrons and very Low-density Lipoprotein (LDL). These particles are rich in both

triglyceride and cholesterol and appear to be at least as atherogenic as LDL.

Cardiovascular disease (clinical) DSS NHIG, Standard 15/02/2006

Information specific to this data set:

A relationship between triglyceride and High-density Lipoprotein Cholesterol (HDL-C) and chronic heart disease (CHD) event rates has been shown. This view is supported by the observation that the remnants of triglyceride-rich lipoproteins are the particles that occur in dysbetalipoproteinaemia, a condition associated with a very high risk of premature atherosclerotic vascular disease. There have been two comprehensive reviews of the relationship between plasma triglyceride and CHD (see Criqui et al. 1993 and Austin et al. 1991). Criqui concludes that triglyceride is not an independent predictor of CHD and is probably not causally related to the disease, while Austin provides a compelling case for a causal role of (at least) some triglyceride rich lipoproteins. Conclusions drawn from population studies of the relationship between plasma triglyceride and the risk of CHD include the following:

- an elevated concentration of plasma triglyceride (> 2.0 mmol/L) is predictive of CHD when associated with either an increased concentration of LDL-C or a decreased concentration of HDL-C.
- the relationship between CHD risk and plasma triglyceride is not continuous, with evidence that the risk is greatest in people with triglyceride levels between 2 and 6 mmol/L (Lipid Management Guidelines - 2001, MJA 2001; 175: S57-S88. National Heart Foundation of Australia and the Cardiac Society of Australia and New Zealand).

It is likely that the positive relationship between plasma triglyceride and CHD, as observed in many population studies, is because an elevated level of plasma triglyceride in some people is a reflection of an accumulation of the atherogenic remnants of chylomicrons and very Low-density Lipoprotein (LDL). These particles are rich in both triglyceride and cholesterol and appear to be at least as atherogenic as LDL.

Diabetes (clinical) DSS NHIG, Superseded 21/09/2005

Information specific to this data set:

Following Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus, the targets for lipids management are:

- to reduce total cholesterol to less than 5.5 mmol/L
- to reduce triglyceride level to less than 2.0 mmol/L
- to increase HDL-C to more than or equal to 1.0 mmol/L.

Alterations in fat transport, often resulting in hypertriglyceridaemia, are well-recognised concomitants of diabetes mellitus.

Elevated plasma triglyceride levels are present in about one third of diabetic patients. It seems that triglycerides are related to the critical role of insulin in the production and removal from plasma of triglyceride-rich lipoproteins. Lifestyle modifications, including weight loss and reduction of excess alcohol intake, are particularly effective for reducing triglyceride and increasing HDL-C.

Diabetes (clinical) DSS NHIG, Standard 21/09/2005

Information specific to this data set:

Following Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus, the targets for lipids management are:

- to reduce total cholesterol to less than 5.5 mmol/L
- to reduce triglyceride level to less than 2.0 mmol/L
- to increase HDL-C to more than or equal to 1.0 mmol/L.

Alterations in fat transport, often resulting in hypertriglyceridaemia, are well-recognised concomitants of diabetes mellitus.

Elevated plasma triglyceride levels are present in about one third of diabetic patients. It seems that triglycerides are related to the critical role of insulin in the production and removal from plasma of triglyceride-rich lipoproteins. Lifestyle modifications, including weight loss and reduction of excess alcohol intake, are particularly effective for reducing triglyceride and increasing HDL-C.

Troponin assay type

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – troponin assay type, code N
<i>METeOR identifier:</i>	285225
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	The type of troponin assay (I or T) used to assess the person's troponin levels, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – troponin assay type
<i>Definition:</i>	Identifies the type of troponin assay (I or T) used to assess the person's troponin levels.
<i>Context:</i>	Health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Troponin assay type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code										
<i>Data type:</i>	Number										
<i>Format:</i>	N										
<i>Maximum character length:</i>	1										
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Cardiac troponin T (cTnT)</td></tr><tr><td>2</td><td>Cardiac troponin I (cTnI)</td></tr><tr><td>3</td><td>Not taken</td></tr><tr><td>9</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	1	Cardiac troponin T (cTnT)	2	Cardiac troponin I (cTnI)	3	Not taken	9	Not stated/inadequately described
Value	Meaning										
1	Cardiac troponin T (cTnT)										
2	Cardiac troponin I (cTnI)										
3	Not taken										
9	Not stated/inadequately described										
<i>Supplementary values:</i>											

Source and reference attributes

<i>Submitting organisation:</i>	Australian Institute of Health and Welfare
---------------------------------	--

Data element attributes

Source and reference attributes

<i>Submitting organisation:</i>	Acute coronary syndrome data working group.
<i>Steward:</i>	The National Heart Foundation of Australia and The Cardiac Society of Australia and New Zealand

Relational attributes

<i>Related metadata references:</i>	Supersedes Troponin assay type, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Information specific to this data set:

For Acute coronary syndrome (ACS) reporting, record the type of troponin assay (I or T) used to assess troponin levels during this presentation.

Acute coronary syndrome (clinical) DSS NHIG, Superseded
07/12/2005

Information specific to this data set:

For Acute coronary syndrome (ACS) reporting, record the type of troponin assay (I or T) used to assess troponin levels during this presentation.

Troponin assay—upper limit of normal range (micrograms per litre)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Laboratory standard – upper limit of normal range for troponin assay, total micrograms per litre N[NNN]
<i>METeOR identifier:</i>	285326
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	Laboratory standard for the value of `troponin T' or `troponin I' measured in micrograms per litre that is the upper boundary of the normal reference range.

Data element concept attributes

<i>Data element concept:</i>	Laboratory standard – upper limit of normal range of troponin assay
<i>Definition:</i>	Laboratory standard for the value of `troponin T' or `troponin I' that is the upper boundary of the normal reference range.
<i>Context:</i>	Health care and clinical settings.
<i>Object class:</i>	Laboratory standard
<i>Property:</i>	Upper limit of normal range of troponin assay

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NNN]
<i>Maximum character length:</i>	4
<i>Supplementary values:</i>	Value Meaning 9999 Not stated/inadequately described
<i>Unit of measure:</i>	Microgram per litre (µg/L)

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record the upper limit of normal (usually the ninety-ninth percentile of a normal population) for the individual laboratory.
-----------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	Acute coronary syndrome data working group
<i>Steward:</i>	The National Heart Foundation of Australia and The Cardiac Society of Australia and New Zealand

Relational attributes

<i>Related metadata references:</i>	Supersedes Troponin assay - upper limit of normal range, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
-------------------------------------	--

Implementation in Data Set Specifications:

Acute coronary syndrome (clinical) DSS NHIG, Standard
07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded
07/12/2005

Troponin level (measured)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – troponin level (measured), total micrograms per litre NN.NN
<i>METeOR identifier:</i>	285253
<i>Registration status:</i>	NHIG, Standard 04/06/2004
<i>Definition:</i>	A person's troponin measured in micrograms per litre.

Data element concept attributes

<i>Data element concept:</i>	Person – troponin level
<i>Definition:</i>	A person's measured troponin.
<i>Context:</i>	Health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Troponin level

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total						
<i>Data type:</i>	Number						
<i>Format:</i>	NN.NN						
<i>Maximum character length:</i>	4						
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>88.88</td><td>Not measured</td></tr><tr><td>99.99</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	88.88	Not measured	99.99	Not stated/inadequately described
Value	Meaning						
88.88	Not measured						
99.99	Not stated/inadequately described						
<i>Unit of measure:</i>	Microgram per litre (µg/L)						

Collection and usage attributes

<i>Guide for use:</i>	CODE 88.88 Not measured This code is used if test for troponin (T or I) was not done.
-----------------------	--

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Measured in different assays dependant upon laboratory methodology. When only one troponin level is recorded, this should be the peak level during the admission.
-----------------------	--

Source and reference attributes

<i>Submitting organisation:</i>	Acute coronary syndrome data working group
<i>Steward:</i>	The National Heart Foundation of Australia and The Cardiac Society of Australia and New Zealand

Relational attributes

Related metadata references:

Supersedes Troponin measured, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

See also Laboratory standard – upper limit of normal range for troponin assay, total micrograms per litre N[NNN] NHIG, Standard 04/06/2004

Implementation in Data Set Specifications:

Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Information specific to this data set:

For Acute coronary syndrome (ACS) reporting, can be used to determine diagnostic strata.

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Information specific to this data set:

For Acute coronary syndrome (ACS) reporting, can be used to determine diagnostic strata.

Tumour size at diagnosis (solid tumours)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person with cancer – solid tumour size (at diagnosis), total millimetres NNN
<i>METeOR identifier:</i>	270184
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The largest dimension of a solid tumour, measured in millimetres.

Data element concept attributes

<i>Data element concept:</i>	Person with cancer – solid tumour size
<i>Definition:</i>	The largest dimension of a solid tumour.
<i>Object class:</i>	Person with cancer
<i>Property:</i>	Solid tumour size

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	String
<i>Format:</i>	NNN
<i>Maximum character length:</i>	3
<i>Supplementary values:</i>	Value Meaning 999 Unknown
<i>Unit of measure:</i>	Millimetre (mm)

Collection and usage attributes

<i>Guide for use:</i>	Size in millimetres with valid values 001 to 997.
-----------------------	---

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	The reporting standard for the size of solid tumours is: Breast cancer or other solid neoplasms - the largest tumour dimension, measured to a precision of 1mm.
-----------------------	--

Relational attributes

<i>Related metadata references:</i>	Supersedes Tumour size at diagnosis - solid tumours, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Cancer (clinical) DSS NHIG, Standard 07/12/2005 <i>Information specific to this data set:</i> This is used to measure the diameter of the largest dimension of breast cancers and other solid neoplasms for patient management, population cancer statistics and research.

Cancer (clinical) DSS NHIG, Superseded 07/12/2005

Information specific to this data set:

This is used to measure the diameter of the largest dimension of breast cancers and other solid neoplasms for patient management, population cancer statistics and research.

Tumour thickness at diagnosis (melanoma)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person with cancer – melanoma thickness (at diagnosis), total millimetres NNN.NN
<i>METeOR identifier:</i>	270185
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The measured thickness of a melanoma in millimetres.

Data element concept attributes

<i>Data element concept:</i>	Person with cancer – melanoma thickness
<i>Definition:</i>	The thickness of a melanoma.
<i>Context:</i>	Patient management, population cancer statistics and research.
<i>Object class:</i>	Person with cancer
<i>Property:</i>	Melanoma thickness

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	String				
<i>Format:</i>	NNN.NN				
<i>Maximum character length:</i>	5				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>999.99</td><td>Unknown</td></tr></tbody></table>	Value	Meaning	999.99	Unknown
Value	Meaning				
999.99	Unknown				
<i>Unit of measure:</i>	Millimetre (mm)				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	The reporting standard for the thickness of melanoma is: Primary cutaneous melanoma - the depth of penetration of tumour cells below the basal layer of the skin; measured to a precision of 0.01mm. Size in millimetres - valid values are: 000.01 to 997.99
-----------------------	---

Relational attributes

<i>Related metadata references:</i>	Supersedes Tumour thickness at diagnosis - melanoma, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Cancer (clinical) DSS NHIG, Standard 07/12/2005 Cancer (clinical) DSS NHIG, Superseded 07/12/2005

Type and sector of employment establishment

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Health professional – establishment type (employment), industry code NN
<i>METeOR identifier:</i>	269954
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The sector of employment and main type of work/speciality area of the health professional, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Health professional – establishment type (employment)
<i>Definition:</i>	<p>For each health profession, type of employment establishment is a self-reporting, condensed industry of employment classification that can be cross-referenced to the Australian and New Zealand Standard Industrial Classification.</p> <p>Sector of employment establishment is government (public) or non-government (private), according to whether or not the employer is a Commonwealth, State or local government agency.</p>
<i>Context:</i>	Health labour force
<i>Object class:</i>	Health professional
<i>Property:</i>	Establishment type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code														
<i>Data type:</i>	String														
<i>Format:</i>	NN														
<i>Maximum character length:</i>	2														
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>Private medical practitioner rooms/surgery (including 24-hour medical clinics)</td></tr><tr><td>02</td><td>Other public non-residential health care facility (e.g. Aboriginal health service, ambulatory centre, outpatient clinic, day surgery centre, medical centre, community health centre)</td></tr><tr><td>03</td><td>Other private non-residential health care (e.g. Aboriginal health service, ambulatory centre, outpatient clinic, day surgery centre, medical centre, community health centre)</td></tr><tr><td>04</td><td>Hospital - acute care (including psychiatric or specialist hospital) hospital (public)</td></tr><tr><td>05</td><td>Hospital - acute care (including psychiatric or specialist hospital) hospital (private)</td></tr><tr><td>06</td><td>Residential health care (e.g. nursing home, hospice, physical disabilities residential centre)</td></tr></tbody></table>	Value	Meaning	01	Private medical practitioner rooms/surgery (including 24-hour medical clinics)	02	Other public non-residential health care facility (e.g. Aboriginal health service, ambulatory centre, outpatient clinic, day surgery centre, medical centre, community health centre)	03	Other private non-residential health care (e.g. Aboriginal health service, ambulatory centre, outpatient clinic, day surgery centre, medical centre, community health centre)	04	Hospital - acute care (including psychiatric or specialist hospital) hospital (public)	05	Hospital - acute care (including psychiatric or specialist hospital) hospital (private)	06	Residential health care (e.g. nursing home, hospice, physical disabilities residential centre)
Value	Meaning														
01	Private medical practitioner rooms/surgery (including 24-hour medical clinics)														
02	Other public non-residential health care facility (e.g. Aboriginal health service, ambulatory centre, outpatient clinic, day surgery centre, medical centre, community health centre)														
03	Other private non-residential health care (e.g. Aboriginal health service, ambulatory centre, outpatient clinic, day surgery centre, medical centre, community health centre)														
04	Hospital - acute care (including psychiatric or specialist hospital) hospital (public)														
05	Hospital - acute care (including psychiatric or specialist hospital) hospital (private)														
06	Residential health care (e.g. nursing home, hospice, physical disabilities residential centre)														

	facility (public)
07	Residential health care (e.g. nursing home, hospice, physical disabilities residential centre) facility (private)
08	Tertiary education institution (public)
09	Tertiary education institution (private)
10	Defence forces
11	Government department or agency (e.g. laboratory, research organisation etc.)
12	Private industry/private enterprise (e.g. insurance, pathology, bank)
13	Other (specified) public
14	Other (specified) private
<i>Supplementary values:</i>	99 Unknown/inadequately described/not stated

Collection and usage attributes

Guide for use:

Establishments are coded into self reporting groupings in the public and private sectors. This can be seen in the code list for medical practitioners.

Minor variations in ordering of sequence and disaggregation of the principal categories will be profession-specific as appropriate; where a more detailed set of codes is used, the essential criterion is that there should not be an overlap of the detailed codes across the Australian and New Zealand Standard Industrial Classification category definitions.

Note:

Public psychiatric hospitals are non-acute care facilities, whereas private psychiatric hospitals are acute care facilities. To minimise the possibility of respondent confusion and mis-reporting, public psychiatric hospitals are included in the grouping for acute care public hospitals.

Source and reference attributes

Origin:

Australian Bureau of Statistics 1993. Australian and New Zealand Standard Industrial Classification (ANZSIC). Cat. No. 1292.0. Canberra: ABS

Data element attributes

Collection and usage attributes

Comments:

Day surgery centres, outpatient clinics and medical centres approved as hospitals under the Health Insurance Act 1973 (Commonwealth) have emerged as a new category for investigation. These will be included in a review of the National Health Labour Force Collection questions and coding frames.

Source and reference attributes

Submitting organisation:

National Health Labour Force Data Working Group

Reference documents:

Australian Bureau of Statistics 1993. Australian and New Zealand Standard Industrial Classification (ANZSIC). Cat. No. 1292.0. Canberra: ABS

Relational attributes

Related metadata references:

Supersedes Type and sector of employment establishment, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications:

Health labour force NMDS NHIG, Standard 01/03/2005

Implementation start date: 01/07/2006

Information specific to this data set:

To analyse distribution of service providers by setting (defined by industry of employer and sector), cross-classified with main type of work and/or specialty area.

Type of accommodation

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – accommodation type (usual), code N[N]
<i>METeOR identifier:</i>	270088
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The type of accommodation setting in which a person usually lives/lived, as represented by a code.
<i>Context:</i>	Admitted patient mental health care: Permits analysis of the usual residential accommodation type of people prior to admission to institutional health care. The setting in which the person usually lives can have a bearing on the types of treatment and support required by the person and the outcomes that result from their treatment.

Data element concept attributes

<i>Data element concept:</i>	Person – accommodation type
<i>Definition:</i>	The setting in which a person lives.
<i>Object class:</i>	Person
<i>Property:</i>	Accommodation type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																				
<i>Data type:</i>	Number																				
<i>Format:</i>	N[N]																				
<i>Maximum character length:</i>	2																				
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Private residence (e.g. house, flat, bedsitter, caravan, boat, independent unit in retirement village), including privately and publicly rented homes</td></tr><tr><td>2</td><td>Psychiatric hospital</td></tr><tr><td>3</td><td>Residential aged care service</td></tr><tr><td>4</td><td>Specialised alcohol/other drug treatment residence</td></tr><tr><td>5</td><td>Specialised mental health community-based residential support service</td></tr><tr><td>6</td><td>Domestic-scale supported living facility (eg. group home for people with disabilities)</td></tr><tr><td>7</td><td>Boarding/rooming house/hostel or hostel type accommodation, not including aged persons' hostel</td></tr><tr><td>8</td><td>Homeless persons' shelter</td></tr><tr><td>9</td><td>Shelter/refuge (not including homeless)</td></tr></tbody></table>	Value	Meaning	1	Private residence (e.g. house, flat, bedsitter, caravan, boat, independent unit in retirement village), including privately and publicly rented homes	2	Psychiatric hospital	3	Residential aged care service	4	Specialised alcohol/other drug treatment residence	5	Specialised mental health community-based residential support service	6	Domestic-scale supported living facility (eg. group home for people with disabilities)	7	Boarding/rooming house/hostel or hostel type accommodation, not including aged persons' hostel	8	Homeless persons' shelter	9	Shelter/refuge (not including homeless)
Value	Meaning																				
1	Private residence (e.g. house, flat, bedsitter, caravan, boat, independent unit in retirement village), including privately and publicly rented homes																				
2	Psychiatric hospital																				
3	Residential aged care service																				
4	Specialised alcohol/other drug treatment residence																				
5	Specialised mental health community-based residential support service																				
6	Domestic-scale supported living facility (eg. group home for people with disabilities)																				
7	Boarding/rooming house/hostel or hostel type accommodation, not including aged persons' hostel																				
8	Homeless persons' shelter																				
9	Shelter/refuge (not including homeless)																				

		persons' shelter)
	10	Other supported accommodation
	11	Prison/remand centre/youth training centre
	12	Public place (homeless)
	13	Other accommodation, not elsewhere classified
<i>Supplementary values:</i>	14	Unknown/unable to determine

Collection and usage attributes

Guide for use:

CODE 3 Residential aged care service

Includes nursing home beds in acute care hospitals.

CODE 4 Specialised alcohol/other drug treatment residence

Includes alcohol/other drug treatment units in psychiatric hospitals.

CODE 5 Specialised mental health community-based residential support service

Specialised mental health community-based residential support services are defined as community-based residential supported accommodation specifically targeted at people with psychiatric disabilities which provides 24-hour support/rehabilitation on a residential basis.

CODE 6 Domestic-scale supported living facility (eg. group home for people with disabilities)

Domestic-scale supported living facilities include group homes for people with disabilities, cluster apartments where a support worker lives on-site, community residential apartments (except mental health), congregate care arrangements. Support is provided by staff on either a live-in or rostered basis, and they may or may not have 24-hour supervision and care.

CODE 10 Other supported accommodation

Includes other supported accommodation facilities such as hostels for people with disabilities and Residential Services/Facilities (Victoria and South Australia only). These facilities provide board and lodging and rostered care workers provide client support services.

Data element attributes

Collection and usage attributes

Guide for use:

'Usual' is defined as the type of accommodation the person has lived in for the most amount of time over the past three months prior to admission to institutional health care or first contact with a community service setting. If a person stays in a particular place of accommodation for four or more days a week over the period, that place of accommodation would be the person's type of usual accommodation. In practice, receiving an answer to questioning about a person's usual accommodation setting may be difficult to achieve. The place the person perceives as their usual accommodation will often prove to be the best approximation of their type of usual accommodation.

Comments:

The changes made to this metadata item are in accordance with the requirements of the National Mental Health Information Strategy Committee and take into consideration corresponding

definitions in other data dictionaries (e.g. Home and Community Care Data Dictionary Version 1 and National Community Services Data Dictionary Version 1).

Relational attributes

Related metadata references:

Supersedes Type of accommodation, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications:

Admitted patient mental health care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient mental health care NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Admitted patient mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Type of augmentation of labour

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Birth event – labour augmentation type, code N
<i>METeOR identifier:</i>	270036
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Methods used to assist progress of labour, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Birth event – labour augmentation type
<i>Definition:</i>	Methods used to assist progress of labour.
<i>Context:</i>	Perinatal care
<i>Object class:</i>	Birth event
<i>Property:</i>	Labour augmentation type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code														
<i>Data type:</i>	Number														
<i>Format:</i>	N														
<i>Maximum character length:</i>	1														
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>0</td><td>None</td></tr><tr><td>1</td><td>Oxytocin</td></tr><tr><td>2</td><td>Prostaglandins</td></tr><tr><td>3</td><td>Artificial rupture of membranes (ARM)</td></tr><tr><td>4</td><td>Other</td></tr><tr><td>5</td><td>Not stated</td></tr></tbody></table>	Value	Meaning	0	None	1	Oxytocin	2	Prostaglandins	3	Artificial rupture of membranes (ARM)	4	Other	5	Not stated
Value	Meaning														
0	None														
1	Oxytocin														
2	Prostaglandins														
3	Artificial rupture of membranes (ARM)														
4	Other														
5	Not stated														
<i>Supplementary values:</i>															

Collection and usage attributes

<i>Comments:</i>	Prostaglandin is listed as a method of augmentation in the data domain. Advice from the Royal Australian and New Zealand College of Obstetricians and Gynaecologists and the manufacturer indicates that vaginal prostaglandin use is not recommended or supported as a method of augmentation of labour as it may significantly increase the risk of uterine hyperstimulation. In spite of this, the method is being used and it is considered important to monitor its use for augmentation.
------------------	--

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	More than one method of augmentation can be recorded, except where 0=none applies.
-----------------------	--

Collection units need to edit carefully the use of prostaglandins as an augmentation method. Results from checking records have shown that either the onset of labour was incorrect or that the augmentation method was incorrectly selected.

Comments:

Type of augmentation determines the progress and duration of labour and may influence the method of delivery and the health status of the baby at birth.

Source and reference attributes

Submitting organisation:

National Perinatal Data Development Committee

Relational attributes

Related metadata references:

Supersedes Type of augmentation of labour, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Type of labour induction

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Birth event – labour induction type, code N
<i>METeOR identifier:</i>	270037
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Method used to induce labour, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Birth event – labour induction type
<i>Definition:</i>	Method used to induce labour.
<i>Context:</i>	Perinatal statistics
<i>Object class:</i>	Birth event
<i>Property:</i>	Labour induction type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code												
<i>Data type:</i>	Number												
<i>Format:</i>	N												
<i>Maximum character length:</i>	1												
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>0</td><td>None</td></tr><tr><td>1</td><td>Oxytocin</td></tr><tr><td>2</td><td>Prostaglandins</td></tr><tr><td>3</td><td>Artificial rupture of membranes (ARM)</td></tr><tr><td>4</td><td>Other</td></tr></tbody></table>	Value	Meaning	0	None	1	Oxytocin	2	Prostaglandins	3	Artificial rupture of membranes (ARM)	4	Other
Value	Meaning												
0	None												
1	Oxytocin												
2	Prostaglandins												
3	Artificial rupture of membranes (ARM)												
4	Other												

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	More than one method of induction can be recorded, except where 0=none applies.
<i>Comments:</i>	Type of induction determines the progress and duration of labour and may influence the method of delivery and the health status of the baby at birth.

Source and reference attributes

<i>Submitting organisation:</i>	National Perinatal Data Development Committee
---------------------------------	---

Relational attributes

<i>Related metadata references:</i>	Supersedes Type of labour induction, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
-------------------------------------	--

Type of usual accommodation

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – accommodation type (prior to admission), code N
<i>METeOR identifier:</i>	270079
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The type of physical accommodation the person lived in prior to admission .
<i>Context:</i>	Admitted patient mental health care: Permits analysis of the prior residential accommodation type of people admitted to residential aged care services or other institutional care.

Data element concept attributes

<i>Data element concept:</i>	Person – accommodation type
<i>Definition:</i>	The setting in which a person lives.
<i>Object class:</i>	Person
<i>Property:</i>	Accommodation type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																
<i>Data type:</i>	Number																
<i>Format:</i>	N																
<i>Maximum character length:</i>	1																
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>House or flat</td></tr><tr><td>2</td><td>Independent unit as part of retirement village or similar</td></tr><tr><td>3</td><td>Hostel or hostel type accommodation</td></tr><tr><td>4</td><td>Psychiatric hospital</td></tr><tr><td>5</td><td>Acute hospital</td></tr><tr><td>6</td><td>Other accommodation</td></tr><tr><td>7</td><td>No usual residence</td></tr></tbody></table>	Value	Meaning	1	House or flat	2	Independent unit as part of retirement village or similar	3	Hostel or hostel type accommodation	4	Psychiatric hospital	5	Acute hospital	6	Other accommodation	7	No usual residence
Value	Meaning																
1	House or flat																
2	Independent unit as part of retirement village or similar																
3	Hostel or hostel type accommodation																
4	Psychiatric hospital																
5	Acute hospital																
6	Other accommodation																
7	No usual residence																

Collection and usage attributes

<i>Collection methods:</i>	The above classifications have been based on Question 16 of Form NH5. The Australian Government Department of Health and Aged Care has introduced a new Aged Care Application and Approval form which replaces the NH5.
----------------------------	--

Data element attributes

Collection and usage attributes

Collection methods:

This metadata item is not available for New South Wales State nursing homes. As this item includes only details of physical accommodation before admission it was decided to have details of the relational basis of accommodation before admission collected as a separate metadata item (see metadata item Admission mode).

Source and reference attributes

Submitting organisation:

National minimum data set working parties

Relational attributes

Related metadata references:

Supersedes Type of usual accommodation, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications:

Admitted patient mental health care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient mental health care NMDS NHIG, Superseded 23/10/2006

Implementation start date: 01/07/2006

Implementation end date: 30/06/2007

Admitted patient mental health care NMDS 2007-2008 NHIG, Standard 23/10/2006

Implementation start date: 01/07/2007

Type of visit to emergency department

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Non-admitted patient emergency department service episode – type of visit to emergency department, code N
<i>METeOR identifier:</i>	270362
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The reason the patient presents to an emergency department, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Non-admitted patient emergency department service episode – type of visit to emergency department
<i>Definition:</i>	The reason the patient presents to an emergency department.
<i>Context:</i>	Hospital non-admitted patient care
<i>Object class:</i>	Non-admitted patient emergency department service episode
<i>Property:</i>	Type of visit to emergency department

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code												
<i>Data type:</i>	Number												
<i>Format:</i>	N												
<i>Maximum character length:</i>	1												
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Emergency presentation: attendance for an actual or suspected condition which is sufficiently serious to require acute unscheduled care.</td></tr><tr><td>2</td><td>Return visit, planned: presentation is planned and is a result of a previous emergency department presentation or return visit.</td></tr><tr><td>3</td><td>Pre-arranged admission: a patient who presents at the emergency department for either clerical, nursing or medical processes to be undertaken, and admission has been pre-arranged by the referring medical officer and a bed allocated.</td></tr><tr><td>4</td><td>Patient in transit: the emergency department is responsible for care and treatment of a patient awaiting transport to another facility.</td></tr><tr><td>5</td><td>Dead on arrival: a patient who is dead on arrival at the emergency department.</td></tr></tbody></table>	Value	Meaning	1	Emergency presentation: attendance for an actual or suspected condition which is sufficiently serious to require acute unscheduled care.	2	Return visit, planned: presentation is planned and is a result of a previous emergency department presentation or return visit.	3	Pre-arranged admission: a patient who presents at the emergency department for either clerical, nursing or medical processes to be undertaken, and admission has been pre-arranged by the referring medical officer and a bed allocated.	4	Patient in transit: the emergency department is responsible for care and treatment of a patient awaiting transport to another facility.	5	Dead on arrival: a patient who is dead on arrival at the emergency department.
Value	Meaning												
1	Emergency presentation: attendance for an actual or suspected condition which is sufficiently serious to require acute unscheduled care.												
2	Return visit, planned: presentation is planned and is a result of a previous emergency department presentation or return visit.												
3	Pre-arranged admission: a patient who presents at the emergency department for either clerical, nursing or medical processes to be undertaken, and admission has been pre-arranged by the referring medical officer and a bed allocated.												
4	Patient in transit: the emergency department is responsible for care and treatment of a patient awaiting transport to another facility.												
5	Dead on arrival: a patient who is dead on arrival at the emergency department.												

Data element attributes

Collection and usage attributes

Comments: Required for analysis of emergency department services.

Source and reference attributes

Submitting organisation: National Institution Based Ambulatory Model Reference Group

Origin: National Health Data Committee

Relational attributes

Related metadata references: Supersedes Type of visit to emergency department, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Non-admitted patient emergency department care NMDS NHIG, Standard 24/03/2006

Implementation start date: 01/07/2006

Non-admitted patient emergency department care NMDS NHIG, Superseded 07/12/2005

Non-admitted patient emergency department care NMDS NHIG, Superseded 24/03/2006

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Urgency of admission

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Episode of admitted patient care – admission urgency status, code N
<i>METeOR identifier:</i>	269986
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether the admission has an urgency status assigned and, if so, whether admission occurred on an emergency basis, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Episode of admitted patient care – admission urgency status
<i>Definition:</i>	<p>Whether the admission has an urgency status assigned and, if so, whether admission occurred on an emergency basis.</p> <p>An emergency admission is an admission of a patient for care or treatment which, in the opinion of the treating clinician, is necessary and admission for which should occur within 24 hours.</p> <p>An elective admission is an admission of a patient for care or treatment which, in the opinion of the treating clinician, is necessary and admission for which can be delayed for at least 24 hours.</p> <p>Admissions for which an urgency status is usually not assigned are:</p> <ul style="list-style-type: none">• admissions for normal delivery (obstetric)• admissions which begin with the birth of the patient, or when it was intended that the birth occur in the hospital, commence shortly after the birth of the patient• statistical admissions• planned readmissions for the patient to receive limited care or treatment for a current condition, for example dialysis or chemotherapy.
<i>Context:</i>	Admitted patient care.
<i>Object class:</i>	Episode of admitted patient care
<i>Property:</i>	Admission urgency status

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Urgency status assigned - emergency</td></tr><tr><td>2</td><td>Urgency status assigned - elective</td></tr></tbody></table>	Value	Meaning	1	Urgency status assigned - emergency	2	Urgency status assigned - elective
Value	Meaning						
1	Urgency status assigned - emergency						
2	Urgency status assigned - elective						

	3	Urgency status not assigned
<i>Supplementary values:</i>	9	Not known/not reported

Data element attributes

Collection and usage attributes

Guide for use:

CODE 1 Urgency status assigned - emergency

Emergency admission:

The following guidelines may be used by health professionals, hospitals and health insurers in determining whether an emergency admission has occurred. These guidelines should not be considered definitive.

An emergency admission occurs if one or more of the following clinical conditions are applicable such that the patient required admission within 24 hours.

Such a patient would be:

- at risk of serious morbidity or mortality and requiring urgent assessment and/or resuscitation; or
- suffering from suspected acute organ or system failure; or
- suffering from an illness or injury where the viability or function of a body part or organ is acutely threatened; or
- suffering from a drug overdose, toxic substance or toxin effect; or
- experiencing severe psychiatric disturbance whereby the health of the patient or other people is at immediate risk; or
- suffering severe pain where the viability or function of a body part or organ is suspected to be acutely threatened; or
- suffering acute significant haemorrhage and requiring urgent assessment and treatment; or
- suffering gynaecological or obstetric complications; or
- suffering an acute condition which represents a significant threat to the patient's physical or psychological wellbeing; or
- suffering a condition which represents a significant threat to public health.

If an admission meets the definition of emergency above, it is categorised as emergency, regardless of whether the admission occurred within 24 hours of such a categorisation being made, or after 24 hours or more.

CODE 2 Urgency status assigned - Elective

Elective admissions:

If an admission meets the definition of elective above, it is categorised as elective, regardless of whether the admission actually occurred after 24 hours or more, or it occurred within 24 hours. The distinguishing characteristic is that the admission could be delayed by at least 24 hours.

Scheduled admissions:

A patient who expects to have an elective admission will often have that admission scheduled in advance. Whether or not the admission has been scheduled does not affect the categorisation of the admission as emergency or elective, which depends only on whether it meets the definitions above. That is, patients both with and without a scheduled admission can be admitted on

either an emergency or elective basis.

Admissions from elective surgery waiting lists:

Patients on waiting lists for elective surgery are assigned a Clinical urgency status which indicates the clinical assessment of the urgency with which a patient requires elective hospital care. On admission, they will also be assigned an urgency of admission category, which may or may not be elective:

- Patients who are removed from elective surgery waiting lists on admission as an elective patient for the procedure for which they were waiting (see code 1 in metadata item Reason for removal from an elective surgery waiting list code N) will be assigned an Admission urgency status code N code of 2. In that case, their clinical urgency category could be regarded as further detail on how urgent their admission was.
- Patients who are removed from elective surgery waiting lists on admission as an emergency patient for the procedure for which they were waiting (see code 2 in metadata item Reason for removal from an elective surgery waiting list code N), will be assigned an Admission urgency status code N code of 1.

CODE 3 Urgency status not assigned

Admissions for which an urgency status is usually not assigned:

An urgency status can be assigned for admissions of the types listed above for which an urgency status is not usually assigned. For example, a patient who is to have an obstetric admission may have one or more of the clinical conditions listed above and be admitted on an emergency basis.

CODE 9 Not known/not reported

This code is used when it is not known whether or not an urgency status has been assigned, or when an urgency status has been assigned but is not known.

Source and reference attributes

Submitting organisation: Emergency definition working party

Origin: National Health Data Committee

Relational attributes

Related metadata references: Supersedes Urgency of admission, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Admitted patient care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Admitted patient care NMDS 2007-2008 NHIG, Standard 29/11/2006

Implementation start date: 01/07/2007

Vascular history

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – vascular condition status (history), code NN
<i>METeOR identifier:</i>	269958
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	Whether the person has had a history of vascular conditions, as represented by a code.
<i>Context:</i>	The vascular history of the patient is important as an element in defining future risk for a cardiovascular event and as a factor in determining best practice management for various cardiovascular risk factor(s). It may be used to map vascular conditions, assist in risk stratification and link to best practice management.

Data element concept attributes

<i>Data element concept:</i>	Person – vascular condition status
<i>Definition:</i>	Describes the vascular condition of the person.
<i>Context:</i>	Public health, health care and clinical settings
<i>Object class:</i>	Person
<i>Property:</i>	Vascular condition status

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																														
<i>Data type:</i>	String																														
<i>Format:</i>	NN																														
<i>Maximum character length:</i>	2																														
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>Myocardial infarction</td></tr><tr><td>02</td><td>Unstable angina pectoris</td></tr><tr><td>03</td><td>Angina</td></tr><tr><td>04</td><td>Heart failure</td></tr><tr><td>05</td><td>Atrial fibrillation</td></tr><tr><td>06</td><td>Other dysrhythmia or conductive disorder</td></tr><tr><td>07</td><td>Rheumatic heart disease</td></tr><tr><td>08</td><td>Non-rheumatic valvular heart disease</td></tr><tr><td>09</td><td>Left ventricular hypertrophy</td></tr><tr><td>10</td><td>Stroke</td></tr><tr><td>11</td><td>Transient ischaemic attack</td></tr><tr><td>12</td><td>Hypertension</td></tr><tr><td>13</td><td>Peripheral vascular disease (includes abdominal aortic aneurism)</td></tr><tr><td>14</td><td>Deep vein thrombosis</td></tr></tbody></table>	Value	Meaning	01	Myocardial infarction	02	Unstable angina pectoris	03	Angina	04	Heart failure	05	Atrial fibrillation	06	Other dysrhythmia or conductive disorder	07	Rheumatic heart disease	08	Non-rheumatic valvular heart disease	09	Left ventricular hypertrophy	10	Stroke	11	Transient ischaemic attack	12	Hypertension	13	Peripheral vascular disease (includes abdominal aortic aneurism)	14	Deep vein thrombosis
Value	Meaning																														
01	Myocardial infarction																														
02	Unstable angina pectoris																														
03	Angina																														
04	Heart failure																														
05	Atrial fibrillation																														
06	Other dysrhythmia or conductive disorder																														
07	Rheumatic heart disease																														
08	Non-rheumatic valvular heart disease																														
09	Left ventricular hypertrophy																														
10	Stroke																														
11	Transient ischaemic attack																														
12	Hypertension																														
13	Peripheral vascular disease (includes abdominal aortic aneurism)																														
14	Deep vein thrombosis																														

	15	Other atherosclerotic disease
	16	Carotid stenosis
	17	Vascular renal disease
	18	Vascular retinopathy (hypertensive)
	19	Vascular retinopathy (diabetic)
	97	Other vascular
	98	No vascular history
<i>Supplementary values:</i>	99	Unknown/not stated /not specified

Collection and usage attributes

Comments: Can be mapped to the current version of ICD-10-AM.

Source and reference attributes

Origin: International Classification of Diseases - Tenth Revision - Australian Modification (3rd Edition 2000), National Centre for Classification in Health, Sydney

Data element attributes

Collection and usage attributes

Guide for use: More than one code can be recorded.

Collection methods: Ideally, vascular history information is derived from and substantiated by clinical documentation.

Source and reference attributes

Submitting organisation: Cardiovascular Data Working Group

Origin: National Centre for Classification in Health
National Data Standards for Injury Surveillance Advisory Group

Relational attributes

Related metadata references: Supersedes Vascular history, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Cardiovascular disease (clinical) DSS NHIG, Superseded 15/02/2006

Cardiovascular disease (clinical) DSS NHIG, Standard 15/02/2006

Vascular procedures

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – vascular procedures (history), code NN
<i>METeOR identifier:</i>	269962
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The vascular procedures the person has undergone, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – vascular procedure
<i>Definition:</i>	Describes the vascular procedures the person has undergone.
<i>Context:</i>	Public health and health care: This metadata item is important for tracking cardiovascular patient management against appropriate practice for cardiovascular presentation(s) and risk factor(s) the person may exhibit.
<i>Object class:</i>	Person
<i>Property:</i>	Vascular procedure

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																																
<i>Data type:</i>	String																																
<i>Format:</i>	NN																																
<i>Maximum character length:</i>	2																																
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>Amputation for arterial vascular insufficiency</td></tr><tr><td>02</td><td>Carotid endarterectomy</td></tr><tr><td>03</td><td>Carotid angioplasty/stenting</td></tr><tr><td>04</td><td>Coronary angioplasty/stenting</td></tr><tr><td>05</td><td>Coronary artery bypass grafting</td></tr><tr><td>06</td><td>Renal artery angioplasty/stenting</td></tr><tr><td>07</td><td>Heart transplant</td></tr><tr><td>08</td><td>Heart valve surgery</td></tr><tr><td>09</td><td>Abdominal aortic aneurism repair/bypass graft/stenting</td></tr><tr><td>10</td><td>Cerebral circulation angioplasty/stenting</td></tr><tr><td>11</td><td>Femoral/popliteal bypass/graft/stenting</td></tr><tr><td>12</td><td>Congenital heart and blood vessel defect surgery</td></tr><tr><td>13</td><td>Permanent pacemaker implantation</td></tr><tr><td>14</td><td>Implantable cardiac defibrillator</td></tr><tr><td>98</td><td>Other</td></tr></tbody></table>	Value	Meaning	01	Amputation for arterial vascular insufficiency	02	Carotid endarterectomy	03	Carotid angioplasty/stenting	04	Coronary angioplasty/stenting	05	Coronary artery bypass grafting	06	Renal artery angioplasty/stenting	07	Heart transplant	08	Heart valve surgery	09	Abdominal aortic aneurism repair/bypass graft/stenting	10	Cerebral circulation angioplasty/stenting	11	Femoral/popliteal bypass/graft/stenting	12	Congenital heart and blood vessel defect surgery	13	Permanent pacemaker implantation	14	Implantable cardiac defibrillator	98	Other
Value	Meaning																																
01	Amputation for arterial vascular insufficiency																																
02	Carotid endarterectomy																																
03	Carotid angioplasty/stenting																																
04	Coronary angioplasty/stenting																																
05	Coronary artery bypass grafting																																
06	Renal artery angioplasty/stenting																																
07	Heart transplant																																
08	Heart valve surgery																																
09	Abdominal aortic aneurism repair/bypass graft/stenting																																
10	Cerebral circulation angioplasty/stenting																																
11	Femoral/popliteal bypass/graft/stenting																																
12	Congenital heart and blood vessel defect surgery																																
13	Permanent pacemaker implantation																																
14	Implantable cardiac defibrillator																																
98	Other																																

Data element attributes

Collection and usage attributes

<i>Collection methods:</i>	Ideally, Vascular procedure information is derived from and substantiated by clinical documentation.
<i>Comments:</i>	In settings where the monitoring of a person's health is ongoing and where a history can change over time (such as general practice), the Service contact – service contact date, DDMMYYYY should be recorded.

Source and reference attributes

<i>Submitting organisation:</i>	Cardiovascular Data Working Group
<i>Origin:</i>	Australian Institute of Health and Welfare (AIHW) 2001. Heart, stroke and vascular diseases - Australian facts 2001. AIHW Cat. No. CVD 13. Canberra: AIHW, National Heart foundation of Australia, National Stroke Foundation of Australia (CVD Series No. 14)

Relational attributes

<i>Related metadata references:</i>	Supersedes Vascular procedures, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Cardiovascular disease (clinical) DSS NHIG, Superseded 15/02/2006 Cardiovascular disease (clinical) DSS NHIG, Standard 15/02/2006

Visual acuity (left eye)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – visual acuity (left eye), code NN
<i>METeOR identifier:</i>	269963
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	A person's left eye visual acuity, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – visual acuity
<i>Definition:</i>	The visual acuity test measures the smallest letters that a person can read on a standardised chart at a distance of 6 metres (20 feet) wearing glasses if needed.
<i>Context:</i>	Public health, health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Visual acuity

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																												
<i>Data type:</i>	String																												
<i>Format:</i>	NN																												
<i>Maximum character length:</i>	2																												
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>6/5</td></tr><tr><td>02</td><td>6/6</td></tr><tr><td>03</td><td>6/9</td></tr><tr><td>04</td><td>6/12</td></tr><tr><td>05</td><td>6/18</td></tr><tr><td>06</td><td>6/24</td></tr><tr><td>07</td><td>6/36</td></tr><tr><td>08</td><td>6/60</td></tr><tr><td>09</td><td>CF (count fingers)</td></tr><tr><td>10</td><td>HM (hand movement)</td></tr><tr><td>11</td><td>PL (perceive light)</td></tr><tr><td>12</td><td>BL (blind)</td></tr><tr><td>13</td><td>6/7.5</td></tr></tbody></table>	Value	Meaning	01	6/5	02	6/6	03	6/9	04	6/12	05	6/18	06	6/24	07	6/36	08	6/60	09	CF (count fingers)	10	HM (hand movement)	11	PL (perceive light)	12	BL (blind)	13	6/7.5
Value	Meaning																												
01	6/5																												
02	6/6																												
03	6/9																												
04	6/12																												
05	6/18																												
06	6/24																												
07	6/36																												
08	6/60																												
09	CF (count fingers)																												
10	HM (hand movement)																												
11	PL (perceive light)																												
12	BL (blind)																												
13	6/7.5																												
<i>Supplementary values:</i>	99 Not stated/inadequately described																												

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record actual result for both right and left eyes:
-----------------------	--

- 1st field: Right eye
- 2nd field: Left eye.

Test wearing distance glasses if prescribed.
Use pinhole if vision less than 6/6.

Collection methods:

One of the most often utilised tests for visual acuity uses the Snellen chart.

- At a distance of 6 metres all subjects should be able to read the 6/6 line with each eye using the proper refractive correction.
- Both eyes are to be opened and then cover one eye with the ocular occluder.
- The observer has to read out the smallest line of letters that he/she can see from the chart.
- This is to be repeated with the other eye.

Eye examination should be performed by an ophthalmologist or a suitably trained clinician:

- within five years of **diagnosis** and then every 1-2 years for patients whose diabetes onset was at age under 30 years
- at diagnosis and then every 1-2 years for patients whose diabetes onset was at age 30 years or more.

Source and reference attributes

Origin:

National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary

Reference documents:

Vision Australia, No 2, 1997/8; University of Melbourne
World Health Organization
US National Library of Medicine
Diabetes Control and Complications Trial: DCCT New England
Journal of Medicine, 329(14), September 30, 1993
Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus

Relational attributes

Related metadata references:

Supersedes Visual acuity, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
See also Person – visual acuity (right eye), code NN NHIG, Standard 01/03/2005

Implementation in Data Set Specifications:

Diabetes (clinical) DSS NHIG, Superseded 21/09/2005

Information specific to this data set:

Record actual result for both right and left eyes (this is a repeating field):

- 1st field: Right eye
- 2nd field: Left eye.

Diabetes (clinical) DSS NHIG, Standard 21/09/2005

Information specific to this data set:

Patients with diabetes have an increased risk of developing several eye complications including retinopathy, cataract and glaucoma that can lead to loss of vision. Regular eye checkups are important for patients suffering from diabetes mellitus. This helps to detect and

treat abnormalities early and to avoid or postpone vision-threatening complications. Assessment by an ophthalmologist is essential:

- at initial examination if the corrected visual acuity is less than 6/6 in either eye
- if at subsequent examinations declining visual acuity is detected
- if any retinal abnormality is detected
- if clear view of retina is not obtained.

Visual acuity (right eye)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – visual acuity (right eye), code NN
<i>METeOR identifier:</i>	270381
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	A person's right eye visual acuity, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Person – visual acuity
<i>Definition:</i>	The visual acuity test measures the smallest letters that a person can read on a standardised chart at a distance of 6 metres (20 feet) wearing glasses if needed.
<i>Context:</i>	Public health, health care and clinical settings.
<i>Object class:</i>	Person
<i>Property:</i>	Visual acuity

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code																												
<i>Data type:</i>	String																												
<i>Format:</i>	NN																												
<i>Maximum character length:</i>	2																												
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>6/5</td></tr><tr><td>02</td><td>6/6</td></tr><tr><td>03</td><td>6/9</td></tr><tr><td>04</td><td>6/12</td></tr><tr><td>05</td><td>6/18</td></tr><tr><td>06</td><td>6/24</td></tr><tr><td>07</td><td>6/36</td></tr><tr><td>08</td><td>6/60</td></tr><tr><td>09</td><td>CF (count fingers)</td></tr><tr><td>10</td><td>HM (hand movement)</td></tr><tr><td>11</td><td>PL (perceive light)</td></tr><tr><td>12</td><td>BL (blind)</td></tr><tr><td>13</td><td>6/7.5</td></tr></tbody></table>	Value	Meaning	01	6/5	02	6/6	03	6/9	04	6/12	05	6/18	06	6/24	07	6/36	08	6/60	09	CF (count fingers)	10	HM (hand movement)	11	PL (perceive light)	12	BL (blind)	13	6/7.5
Value	Meaning																												
01	6/5																												
02	6/6																												
03	6/9																												
04	6/12																												
05	6/18																												
06	6/24																												
07	6/36																												
08	6/60																												
09	CF (count fingers)																												
10	HM (hand movement)																												
11	PL (perceive light)																												
12	BL (blind)																												
13	6/7.5																												
<i>Supplementary values:</i>	99 Not stated/inadequately described																												

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record actual result for both right and left eyes:
-----------------------	--

- 1st field: Right eye
- 2nd field: Left eye.

Test wearing distance glasses if prescribed.

Use pinhole if vision less than 6/6.

Collection methods:

One of the most often utilised tests for visual acuity uses the Snellen chart.

- At a distance of 6 metres all subjects should be able to read the 6/6 line with each eye using the proper refractive correction.
- Both eyes are to be opened and then cover one eye with the ocular occluder.
- The observer has to read out the smallest line of letters that he/she can see from the chart.
- This is to be repeated with the other eye.

Eye examination should be performed by an ophthalmologist or a suitably trained clinician:

- within five years of **diagnosis** and then every 1-2 years for patients whose diabetes onset was at age under 30 years
- at diagnosis and then every 1-2 years for patients whose diabetes onset was at age 30 years or more.

Source and reference attributes

Origin:

National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary

Reference documents:

Vision Australia, No 2, 1997/8; University of Melbourne

World Health Organization

US National Library of Medicine

Diabetes Control and Complications Trial: DCCT New England Journal of Medicine, 329(14), September 30, 1993

Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus

Relational attributes

Related metadata references:

See also Person – visual acuity (left eye), code NN NHIG, Standard 01/03/2005

Supersedes Visual acuity, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005

Implementation in Data Set Specifications:

Diabetes (clinical) DSS NHIG, Superseded 21/09/2005

Information specific to this data set:

Record actual result for both right and left eyes (this is a repeating field):

- 1st field: Right eye
- 2nd field: Left eye.

Diabetes (clinical) DSS NHIG, Standard 21/09/2005

Information specific to this data set:

Patients with diabetes have an increased risk of developing several eye complications including retinopathy, cataract and glaucoma that can lead to loss of vision. Regular eye checkups are important for patients suffering from diabetes mellitus. This helps to detect and

treat abnormalities early and to avoid or postpone vision-threatening complications. Assessment by an ophthalmologist is essential:

- at initial examination if the corrected visual acuity is less than 6/6 in either eye
- if at subsequent examinations declining visual acuity is detected
- if any retinal abnormality is detected
- if clear view of retina is not obtained.

Waist circumference (measured)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – waist circumference (measured), total centimetres NN[N].N
<i>METeOR identifier:</i>	270129
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	A person's waist circumference measured in centimetres.

Data element concept attributes

<i>Data element concept:</i>	Person – waist circumference
<i>Definition:</i>	A person's waist circumference measured half way between the inferior margin of the last rib and the crest of the ilium in the mid-axillary plane.
<i>Context:</i>	Public health, health care and clinical settings: Originally used in the calculation of Waist-to-hip ratio which requires the measurement of hip circumference and waist circumference as a predictor of obesity-related morbidity and mortality. More recently it has been used in its own right as an indicator of risk associated with excess abdominal fat.
<i>Object class:</i>	Person
<i>Property:</i>	Waist circumference

Collection and usage attributes

<i>Comments:</i>	This metadata item applies to persons of all ages. There is evidence that waist circumference alone might be used to identify people at health risk both from being overweight and from having a central fat distribution (Lean et al. 1995; Han et al. 1995; Pouliot et al. 1994; Seidell et al. 1992). It has been suggested that waist circumference as an index of truncal adiposity in adults may have certain advantages over other measurements of adiposity in predicting obesity related diseases. However, among children and adolescents, waist circumference measures should only be used as a measure of variation in an individual. As yet, no age appropriate cut-off points indicative of risk factors have been developed for use among children and adolescents.
------------------	---

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	NN[N].N
<i>Maximum character length:</i>	4
<i>Supplementary values:</i>	Value Meaning 999.9 Not measured
<i>Unit of measure:</i>	Centimetre (cm)

Data element attributes

Collection and usage attributes

Collection methods:

The collection of anthropometric measurements, particularly in those who are overweight or obese or who are concerned about their weight, should be performed with great sensitivity, and without drawing attention to an individual's weight.

The measurement protocol described below is that recommended by the World Health Organization (WHO Expert Committee 1995) which was adapted from Lohman et al. (1988) and the International Society for the Advancement of Kinanthropometry as described by Norton et al. (1996).

In order to ensure consistency in measurement, the following measurement protocol should be used.

Measurement protocol:

The measurement of waist circumference requires a narrow (The subject should remove any belts and heavy outer clothing. Measurement of waist circumference should be taken over at most one layer of light clothing. Ideally the measure is made directly over the skin.

The subject stands comfortably with weight evenly distributed on both feet, and the feet separated about 25-30 cm. The arms should hang loosely at the sides. Posture can affect waist circumference. The measurement is taken midway between the inferior margin of the last rib and the crest of the ilium, in the mid-axillary plane. Each landmark should be palpated and marked, and the midpoint determined with a tape measure and marked.

The circumference is measured with an inelastic tape maintained in a horizontal plane, at the end of normal expiration. The tape is snug, but does not compress underlying soft tissues. The measurer is positioned by the side of the subject to read the tape. To ensure contiguity of the two parts of the tape from which the circumference is to be determined, the cross-handed technique of measurement, as described by Norton et al. (1996), should be used. Ideally an assistant will check the position of the tape on the opposite side of the subject's body.

The measurement is recorded at the end of a normal expiration to the nearest 0.1 cm. Take a repeat measurement and record it to the nearest 0.1 cm. If the two measurements disagree by more than 1 cm, take a third measurement. All raw measurements should be recorded on the data collection form. If practical, it is preferable to enter the raw data into the database as this enables intra-observer and, where relevant, inter-observer errors to be assessed. The subject's measured waist circumference is subsequently calculated as the mean of the two observations, or the mean of the two closest measurements if a third is taken, and recorded on the form. If only a mean value is entered into the database then the data collection forms should be retained.

It may be necessary to round the mean value to the nearest 0.1 cm. If so, rounding should be to the nearest even digit to reduce systematic over-reporting (Armitage & Berry 1994). For example, a mean value of 72.25 cm would be rounded to 72.2

cm, while a mean value of 72.35 cm would be rounded to 72.4 cm.

Validation and quality control measures:

Steel tapes should be checked against a 1 metre engineer's rule every 12 months. If tapes other than steel are used they should be checked daily against a steel rule.

Within- and, if relevant, between-observer variability should be reported. They can be assessed by the same (within-) or different (between-) observers repeating the measurement, on the same subjects, under standard conditions after a short time interval. The standard deviation of replicate measurements (technical error of measurement (Pederson & Gore 1996) between observers should not exceed 2% and be less than 1.5% within observers.

Extreme values at the lower and upper end of the distribution of measured waist circumference should be checked both during data collection and after data entry. Individuals should not be excluded on the basis of true biological difference.

Last-digit preference, and preference or avoidance of certain values, should be analysed in the total sample and (if relevant) by observer, survey site and over time if the survey period is long.

Comments:

This metadata item is recommended for use in population surveys and health care settings.

It is recommended that in population surveys, sociodemographic data including ethnicity should be collected, as well as other risk factors including physiological status (e.g. pregnancy), physical activity, smoking and alcohol consumption. Summary statistics may need to be adjusted for these variables.

National health metadata items currently exist for sex, date of birth, country of birth, Indigenous status and smoking. Metadata items are being developed for physical activity.

Presentation of data:

Means, 95% confidence intervals, medians and centiles should be reported to one decimal place. Where the sample permits, population estimates should be presented by sex and 5-year age groups. However 5-year age groups are not generally suitable for children and adolescents. Estimates based on sample surveys may need to take into account sampling weights.

For consistency with conventional practice, and for current comparability with international data sets, recommended centiles are 5, 10, 15, 25, 50, 75, 85, 90 and 95. To estimate the 5th and 95th centiles, a sample size of at least 200 is recommended for each group for which the centiles are being specified.

For reporting purposes, it may be desirable to present waist circumference in categories. It is recommended that 5-cm groupings are used for this purpose. Waist circumference should not be rounded before categorisation. The following categories may be appropriate for describing the waist circumferences of Australian men, women children and adolescents, although the range will depend on the population.

Waist

35 cm = Waist

40 cm = Waist

... in 5 cm categories
105 cm = Waist
Waist => 110 cm

Source and reference attributes

Submitting organisation: World Health Organization International Society for the Advancement of Kinanthropometry

Relational attributes

Related metadata references: Supersedes Waist circumference - measured, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Is used in the formation of Adult – waist-to-hip ratio, N.NN NHIG, Standard 01/03/2005

Implementation in Data Set Specifications: Cardiovascular disease (clinical) DSS NHIG, Superseded 15/02/2006

Cardiovascular disease (clinical) DSS NHIG, Standard 15/02/2006

Waist circumference risk indicator - adults

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Adult – waist circumference risk indicator, Caucasian adult code N
<i>METeOR identifier:</i>	270205
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The sex specific category of risk of metabolic complications associated with excess abdominal adiposity in adult Caucasians, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Adult – waist circumference risk indicator
<i>Definition:</i>	An indicator of the sex specific category of risk of metabolic complications associated with excess abdominal adiposity in an adult.
<i>Context:</i>	<p>Public health and health care:</p> <p>This metadata item is used as an indicator of risk of metabolic complications associated with overweight and obesity including dyslipidaemia, glucose intolerance and hypertension. On a population basis there is a strong association between abdominal obesity and health risk. Body fat distribution has emerged as an important predictor of obesity-related morbidity and mortality. Abdominal obesity, which is more common in men than women, has, in epidemiological studies, been closely associated with conditions such as coronary heart disease, stroke, non-insulin dependent diabetes mellitus and high blood pressure. Waist circumference as an indicator of risk can be used:</p> <ul style="list-style-type: none">• to indicate the prevalence of abdominal obesity and its sociodemographic distribution (problem identification);• to evaluate health promotion and disease prevention programs (assessment of interventions);• to monitor progress towards national public health policy;• to ascertain determinants and consequences of abdominal obesity; and• in nutrition and physical activity surveillance and long-term planning. <p>Waist circumference is a convenient and simple measurement that is unrelated to height, correlates closely with body mass index (BMI) and waist-to-hip ratio (WHR) and is an approximate index of intra-abdominal fat mass and total body fat. Changes in waist circumference can reflect changes in risk factors for cardiovascular disease and other forms of chronic disease, even though the risks seem to vary in different populations.</p>
<i>Object class:</i>	Adult
<i>Property:</i>	Waist circumference risk indicator

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code	
<i>Data type:</i>	Number	
<i>Format:</i>	N	
<i>Maximum character length:</i>	1	
<i>Permissible values:</i>	Value	Meaning
	1	Not at risk (male waist circumference
	2	Increased (male waist circumference \geq 94 cm, female waist circumference \geq 80 cm)
	3	Substantially increased (male waist circumference \geq 102 cm, female waist circumference \geq 88 cm)
<i>Supplementary values:</i>	9	Not stated/inadequately described

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>This metadata item cannot be determined if waist circumference measured has not been collected (i.e. is coded to 999.9) and/or sex is not stated (i.e. coded to 9).</p> <p>This metadata item applies to persons aged 18 years or older.</p>
<i>Collection methods:</i>	<p>This metadata item should be derived after the data entry of waist circumference measured. It should be stored on the raw data set as a continuous variable and should not be aggregated or rounded.</p>
<i>Comments:</i>	<p>This metadata item is recommended for use in population surveys and health care settings.</p> <p>Recent evidence suggests that waist circumference may provide a more practical correlate of abdominal fat distribution and associated ill health.</p> <p>The identification of risk using waist circumference is population-specific and will depend on levels of obesity and other risk factors for cardiovascular disease and non-insulin dependant diabetes mellitus.</p> <p>Populations differ in the level of risk associated with a particular waist circumference, so that globally applicable cut-off points cannot be developed. For example, complications associated with abdominal fat in black women and those of South Asian descent are markedly higher for a given level of BMI than in Europeans. Also, although women have almost the same absolute risk of coronary heart disease as men at the same WHR, they show increases in relative risk of coronary heart disease at lower waist circumferences than men. Thus, there is a need to develop sex-specific waist circumference cut-off points appropriate for different populations. Hence, the cut-off points used for this metadata item are associated with obesity in Caucasians. This issue is being investigated further.</p> <p>Cut-off points for children and adolescents are also being developed. Research shows that a high childhood BMI and high trunk skin fold values are predictive of abdominal obesity as an adult and waist circumference measures in childhood track well into adulthood.</p>

It is recommended that in population surveys, sociodemographic data including ethnicity should be collected, as well as other risk factors including physiological status (e.g. pregnancy), physical activity, smoking and alcohol consumption. Summary statistics may need to be adjusted for these variables.

National health metadata item currently exist for sex, date of birth, country of birth and Indigenous status and smoking. Metadata items are being developed for physical activity.

Source and reference attributes

Origin:

World Health Organization

Reference documents:

Obesity: Preventing and Managing the Global Epidemic: Report of a World Health Organization (WHO) Expert Committee. Geneva: WHO, 2000 as described by Han TS et al (1995)

Relational attributes

Related metadata references:

Supersedes Waist circumference risk indicator - adults, version 1, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Waist-to-hip ratio

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Adult – waist-to-hip ratio, N.NN
<i>METeOR identifier:</i>	270207
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	A ratio calculated by dividing the waist circumference of an adult person by the hip circumference of that same person.

Data element concept attributes

<i>Data element concept:</i>	Adult – waist-to-hip ratio
<i>Definition:</i>	A ratio calculated by dividing the waist circumference of an adult person by the hip circumference of that same person.
<i>Context:</i>	Public health and health care
<i>Object class:</i>	Adult
<i>Property:</i>	Waist-to-hip ratio

Value domain attributes

Representational attributes

<i>Representation class:</i>	Ratio
<i>Data type:</i>	Number
<i>Format:</i>	N.NN
<i>Maximum character length:</i>	3

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>Formula:</p> <p>WHR = waist circumference (cm) divided by hip circumference (cm).</p> <p>Adult WHR is a continuous variable. Adult WHR cannot be calculated if either component necessary for its calculation (i.e. abdominal circumference or hip circumference) has not been collected (i.e. is coded to 999.9).</p>
<i>Collection methods:</i>	<p>As there are no cut-off points for waist to hip ratio for children and adolescents, it is not necessary to calculate this item for those aged under 18 years.</p> <p>Waist-to-hip ratio (WHR) should be derived after the data entry of waist circumference and hip circumference. It should be stored on the raw data set as a continuous variable and should not be aggregated or rounded.</p>
<i>Comments:</i>	<p>Adult cut-off points for WHR, that may define increased risk of cardiovascular disease and all cause mortality, range from 0.9 to 1.0 for men and 0.8 to 0.9 for women (Croft et al. 1995, Bray 1987, Bjorntorp 1985). These values are based primarily on evidence of increased risk of death in European populations, and may not be appropriate for all age and ethnic groups.</p>

In Australia and New Zealand, the cutoffs of >0.9 for males and >0.8 for females were used in the Australian Bureau of Statistics' 1995 National Nutrition Survey.

This metadata item applies to persons aged 18 years or older as no cut off points have been developed for children and adolescents. It is recommended for use in population surveys and health care settings.

More recently it has emerged that waist circumference alone, or in combination with other metabolic measures, is a better indicator of risk and reduces the errors in WHR measurements. WHR is therefore no longer a commonly used measure.

It is recommended that in population surveys, sociodemographic data including ethnicity should be collected, as well as other risk factors including physiological status (e.g. pregnancy), physical activity, smoking and alcohol consumption. Summary statistics may need to be adjusted for these variables.

Body fat distribution has emerged as an important predictor of obesity-related morbidity and mortality. Abdominal obesity, which is more common in men than women, has, in epidemiological studies, been closely associated with conditions such as coronary heart disease, stroke, non-insulin dependent diabetes mellitus and high blood pressure.

Waist- to-hip ratio (WHR) can be used:

- to indicate the prevalence of abdominal obesity and its sociodemographic distribution (problem identification)
- to evaluate health promotion and disease prevention programs (assessment of interventions)
- to monitor progress towards national public health policy
- to ascertain determinants and consequences of abdominal obesity - in nutrition and physical activity surveillance and long-term planning.

Presentation of data:

Means, 95% confidence intervals, medians and centiles should be reported to one decimal place. Where the sample permits, population estimates should be presented by sex and 5-year age groups. Estimates based on sample surveys may need to take into account sampling weights.

For consistency with conventional practice, and for current comparability with international data sets, recommended centiles are 5, 10, 15, 25, 50, 75, 85, 90 and 95. To estimate the 5th and 95th centiles, a sample size of at least 200 is recommended for each group for which the centiles are being specified.

Source and reference attributes

Origin:

National Health Data Committee
National Centre for Monitoring Cardiovascular Disease
Australian Institute of Health and Welfare

Relational attributes

Related metadata references:

Supersedes Waist-to-hip ratio, version 2, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Is formed using Person – waist circumference (measured), total centimetres NN[N].N NHIG, Standard 01/03/2005

Is formed using Person – hip circumference (measured), total

centimetres NN[N].N NHIG, Standard 01/03/2005

Waiting list category

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Elective care waiting list episode – elective care type, code N
<i>METeOR identifier:</i>	335048
<i>Registration status:</i>	NHIG, Standard 07/12/2005
<i>Definition:</i>	The type of elective hospital care that a patient requires, as represented by a code.

Data element concept attributes

<i>Data element concept:</i>	Elective care waiting list episode – elective care type
<i>Definition:</i>	The type of elective hospital care that a patient requires.
<i>Context:</i>	<p>Admitted patients:</p> <p>Hospitals maintain waiting lists which may include patients awaiting hospital care other than elective surgery - for example, dental surgery and oncology treatments.</p> <p>This metadata item is necessary to distinguish patients awaiting elective surgery from those awaiting other types of elective hospital care. The waiting period for patients awaiting transplant or obstetric procedures is largely independent of system resource factors.</p>
<i>Object class:</i>	Elective care waiting list episode
<i>Property:</i>	Elective care type

Value domain attributes

Representational attributes

<i>Representation class:</i>	Code						
<i>Data type:</i>	Number						
<i>Format:</i>	N						
<i>Maximum character length:</i>	1						
<i>Permissible values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Elective surgery</td></tr><tr><td>2</td><td>Other</td></tr></tbody></table>	Value	Meaning	1	Elective surgery	2	Other
Value	Meaning						
1	Elective surgery						
2	Other						

Collection and usage attributes

<i>Guide for use:</i>	<p>Elective surgery comprises elective care where the procedures required by patients are listed in the surgical operations section of the Medicare benefits schedule, with the exclusion of specific procedures frequently done by non-surgical clinicians.</p> <p>Elective care is care that, in the opinion of the treating clinician, is necessary and admission for which can be delayed for at least twenty-four hours.</p> <p>CODE 2 Other</p> <p>Patients awaiting the following procedures should be classified as Code 2 - other:</p> <ul style="list-style-type: none">organ or tissue transplant procedures
-----------------------	---

- procedures associated with obstetrics (e.g. elective caesarean section, cervical suture)
- cosmetic surgery, i.e. when the procedure will not attract a Medicare rebate
- biopsy of:
 - kidney (needle only)
 - lung (needle only)
 - liver and gall bladder (needle only)
- bronchoscopy (including fibre-optic bronchoscopy)
- peritoneal renal dialysis;
- haemodialysis
- colonoscopy
- endoscopic retrograde cholangio-pancreatography (ERCP)
- endoscopy of:
 - biliary tract
 - oesophagus
 - small intestine
 - stomach
- endovascular interventional procedures
- gastroscopy
- miscellaneous cardiac procedures
- oesophagoscopy
- panendoscopy (except when involving the bladder)
- proctosigmoidoscopy
- sigmoidoscopy
- anoscopy
- urethroscopy and associated procedures
- dental procedures not attracting a Medicare rebate
- other diagnostic and non-surgical procedures.

These procedure terms are also defined by the Australian Classification of Health Interventions (ACHI) codes which are listed under Comments below. This coded list is the recommended, but optional, method for determining whether a patient is classified as requiring elective surgery or other care.

CODE 1 Elective surgery

All other elective surgery should be included in this code.

Comments:

The table of Australian Classification of Health Interventions (ACHI) (5th edition) procedure codes was prepared by the National Centre for Classification in Health. Some codes were excluded from the list on the basis that they are usually performed by non-surgeon clinicians. A more extensive and detailed listing of procedure descriptors is under development. This will replace the list in the Guide for use to facilitate more readily the identification of the exclusions when the list of codes is not used.

ACHI CODES FOR THE EXCLUDED PROCEDURES:

Organ or tissue transplant:

90172-00 [555] 90172-01 [555] 90204-00 [659] 90204-01 [659]
 90205-00 [660] 90205-01 [660] 13700-00 [801] 13706-08 [802]
 13706-00 [802] 13706-06 [802] 13706-07 [802] 13706-09 [802]
 13706-10 [802] 30375-21 [817] 90317-00 [954] 90324-00 [981]
 36503-00 [1058] 36503-01 [1058] 14203-01 [1906]

Procedures associated with obstetrics: 16511-00 [1274]
 Obstetric Blocks [1330] to [1345] and [1347]
 90463-01 [1330] 90488-00 [1330]

Biopsy (needle) of:

- kidney: 36561-00 [1047]
- lung: 38412-00 [550]
- liver and gall bladder: 30409-00 [953] 30412-00 [953] 90319-01 [951] 30094-04 [964]

Bronchoscopy:
 41889-00 [543] 41892-00 [544] 41904-00 [546] 41764-02 [416]
 41895-00 [544] 41764-04 [532] 41892-01 [545] 41901-00 [545]
 41898-00 [543] 41898-01 [544] 41889-01 [543] 41849-00 [520]
 41764-03 [520] 41855-00 [520]

Peritoneal renal dialysis:
 13100-06 [1061] 13100-07 [1061] 13100-08 [1061] 13100-00 [1060]

Endoscopy of biliary tract:
 30484-00 [957] 30484-01 [957] 30484-02 [974] 30494-00 [971]
 30452-00 [971] 30491-00 [958] 30491-01 [958] 30485-00 [963]
 30485-01 [963] 30452-01 [958] 30450-00 [959] 30452-02 [959]
 90349-00 [975]

Endoscopy of oesophagus:
 30473-03 [850] 30473-04 [861] 41822-00 [861] 30478-11 [856]
 41819-00 [862] 30478-10 [852] 30478-13 [861] 41816-00 [850]
 41822-00 [861] 41825-00 [852] 30478-12 [856] 41831-00 [862]
 30478-12 [856] 30490-00 [853] 30479-00 [856]

Panendoscopy:
 30476-03 [874] 32095-00 [891] 30568-00 [893] 30569-00 [894]
 30473-05 [1005] 30473-00 [1005] 30473-02 [1005] 30478-00 [1006]
 30478-14 [1006] 30478-01 [1007] 30478-02 [1007] 30478-03 [1007]
 30478-15 [1007] 30478-16 [1007] 30478-17 [1007] 30478-20 [1007]
 30478-21 [1007] 30473-01 [1008] 30478-04 [1008] 30473-06 [1008]
 30478-18 [1008]

Endoscopy of large intestine, rectum and anus:
 32075-00 [904] 32090-00 [905] 32084-00 [905] 30479-02 [908]
 90308-00 [908] 32075-01 [910] 32078-00 [910] 32081-00 [910]
 32090-01 [911] 32093-00 [911] 32084-01 [911] 32087-00 [911]
 30479-01 [931] 90315-00 [933]

Miscellaneous cardiac:
 38603-00 [642] 38600-00 [642] 38256-00 [647] 38256-01 [647]
 38256-02 [647] 38278-00 [648] 38278-01 [648] 38284-00 [648]
 90202-00 [649] 38470-00 [649] 38473-00 [649] 38281-01 [650]
 38281-02 [650] 38281-03 [650] 38281-04 [650] 38281-05 [650]
 38281-06 [650] 38281-07 [651] 38281-07 [651] 38281-08 [651]
 38281-09 [651] 38281-10 [651] 38281-00 [652] 38278-02 [654]
 38456-07 [654] 90203-00 [654] 38284-01 [654] 90219-00 [663]
 38281-11 [655] 38281-12 [655] 38212-00 [665] 38209-00 [665]
 38200-00 [667] 38203-00 [667] 38206-00 [667] 35324-00 [740]
 35315-00 [758] 35315-01 [758] 90214-00 [648] 90202-01 [649]
 38281-13 [652] 38278-03 [654] 38284-02 [654] 90214-01 [654]
 90214-02 [654] 90203-01 [654] 38456-08 [654] 38281-14 [655]
 38212-01 [665] 38213-00 [665]

Endovascular interventional:
 35304-01 [670] 35305-00 [670] 35304-00 [670] 35305-01 [670]
 35310-00 [671] 35310-01 [671] 35310-03 [671] 35310-04 [671]
 35310-02 [671] 35310-05 [671] 34524-00 [694] 13303-00 [694]
 34521-01 [694] 32500-01 [722] 32500-00 [722] 13300-01 [738]
 13300-02 [738] 13319-00 [738] 13300-00 [738] 13815-00 [738]

13815-01 [738] 34521-02 [738] 34530-04 [738] 90220-00 [738]

Urethroscopy:

36800-00 [1090] 36800-01 [1090] 37011-00 [1093] 37008-01 [1093]
37008-00 [1093] 37315-00 [1112] 37318-01 [1116] 36815-01 [1116]
37854-00 [1116] 35527-00 [1116] 37318-04 [1117] 36800-03 [1090]
37318-02 [1116] 37318-03 [1116]

Dental:

Blocks [450] to [490]

97022-00 [451] 97025-00 [451] 97113-00 [453] 97121-01 [454]
97123-01 [454] 97165-01 [455] 97221-00 [456] 97222-00 [456]
97231-00 [456] 97232-00 [456] 97233-00 [456] 97234-00 [456]
97384-00 [461] 97386-01 [461] 97415-00 [462] 97417-00 [462]
97431-00 [463] 97433-00 [463] 97434-00 [463] 97437-00 [463]
97445-00 [464] 97455-00 [464] 97511-01 [465] 97512-01 [465]
97513-01 [465] 97514-02 [465] 97515-02 [465] 97541-01 [465]
97542-01 [465] 97543-01 [465] 97544-00 [465] 97545-00 [465]
97521-01 [466] 97522-01 [466] 97523-01 [466] 97524-00 [466]
97525-00 [466] 97531-00 [466] 97532-00 [466] 97533-00 [466]
97534-00 [466] 97535-00 [466] 97551-01 [466] 97552-01 [466]
97553-01 [466] 97554-01 [466] 97555-01 [466] 97572-01 [469]
97574-01 [469] 97575-00 [469] 97578-00 [469] 97582-01 [469]
97583-01 [469] 97631-00 [470] 97632-00 [471] 97649-00 [471]
97671-00 [473] 97672-00 [473] 97673-00 [473] 97732-00 [474]
97733-00 [474] 97739-00 [474] 97741-00 [475] 97762-00 [476]
97765-00 [476] 97768-00 [476] 97825-00 [479]

Other diagnostic and non-surgical:

90347-01 [983] 30406-00 [983] 90347-02 [983] 30408-00 [983]

Blocks [1820] to [1939], [1940] to [2016]

11003-00 [1825] 11018-00 [1826] 11018-01 [1826] 11614-00 [1851]
11602-00 [1852] 11604-00 [1852] 11605-00 [1852] 11610-00 [1852]
11611-00 [1852] 11612-00 [1852] 11709-00 [1853] 11917-00 [1860]
11919-00 [1860] 96207-00 [1921] 96207-01 [1921] 96207-02 [1921]
96207-03 [1921] 96207-04 [1921] 96207-05 [1921] 96207-06 [1921]
96207-07 [1921] 96207-08 [1921] 96207-09 [1921] 96208-00 [1921]
96208-01 [1921] 96208-02 [1921] 96208-03 [1921] 96208-04 [1921]
96208-05 [1921] 96208-06 [1921] 96208-07 [1921] 96208-08 [1921]
96208-09 [1921] 55084-00 [1943] 55600-00 [1943] 60000-01 [1992]
60048-00 [1996] 60060-00 [1997] 60060-01 [1997] 61442-00 [2012]
96196-00 [1920] 96196-01 [1920] 96196-02 [1920] 96196-03 [1920]
96196-04 [1920] 96196-05 [1920] 96196-06 [1920] 96196-07 [1920]
96196-08 [1920] 96196-09 [1920] 96197-00 [1920] 96197-01 [1920]
96197-02 [1920] 96197-03 [1920] 96197-04 [1920] 96197-05 [1920]
96197-06 [1920] 96197-07 [1920] 96197-08 [1920] 96197-09 [1920]
96198-00 [1920] 96198-01 [1920] 96198-02 [1920] 96198-03 [1920]
96198-04 [1920] 96198-05 [1920] 96198-06 [1920] 96198-07 [1920]
96198-08 [1920] 96198-09 [1920] 96199-00 [1920] 96199-01 [1920]
96199-02 [1920] 96199-03 [1920] 96199-04 [1920] 96199-05 [1920]
96199-06 [1920] 96199-07 [1920] 96199-08 [1920] 96199-09 [1920]
96200-00 [1920] 96200-01 [1920] 96200-02 [1920] 96200-03 [1920]
96200-04 [1920] 96200-05 [1920] 96200-06 [1920] 96200-07 [1920]
96200-08 [1920] 96200-09 [1920] 96201-00 [1920] 96201-01 [1920]
96201-02 [1920] 96201-03 [1920] 96201-04 [1920] 96201-05 [1920]
96201-06 [1920] 96201-07 [1920] 96201-08 [1920] 96201-09 [1920]
96202-00 [1920] 96202-01 [1920] 96202-02 [1920] 96202-03 [1920]
96202-04 [1920] 96202-05 [1920] 96202-06 [1920] 96202-07 [1920]
96202-08 [1920] 96202-09 [1920] 96203-00 [1920] 96203-01 [1920]
96203-02 [1920] 96203-03 [1920] 96203-04 [1920] 96203-05 [1920]

96203-06 [1920] 96203-07 [1920] 96203-08 [1920] 96203-09 [1920]
96204-00 [1920] 96204-01 [1920] 96204-02 [1920] 96204-03 [1920]
96204-04 [1920] 96204-05 [1920] 96204-06 [1920] 96204-07 [1920]
96204-08 [1920] 96204-09 [1920] 96205-00 [1920] 96205-01 [1920]
96205-02 [1920] 96205-03 [1920] 96205-04 [1920] 96205-05 [1920]
96205-06 [1920] 96205-07 [1920] 96205-08 [1920] 96205-09 [1920]
96206-00 [1920] 96206-01 [1920] 96206-02 [1920] 96206-03 [1920]
96206-04 [1920] 96206-05 [1920] 96206-06 [1920] 96206-07 [1920]
96206-08 [1920] 96206-09 [1920]

Source and reference attributes

Reference documents:

National Centre for Classification in Health (NCCH) 2006. The Australian Classification of Health Interventions (ACHI) – Fifth Edition - Tabular list of interventions and Alphabetic index of interventions. Sydney: NCCH, Faculty of Health Sciences, The University of Sydney.

Data element attributes

Source and reference attributes

Submitting organisation:

Hospital Access Program Waiting Lists Working Group
Waiting Times Working Group

Origin:

National Health Data Committee

Relational attributes

Related metadata references:

Supersedes Elective care waiting list episode – elective care type, code N NHIG, Superseded 07/12/2005

Waiting time at a census date

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Elective surgery waiting list episode – waiting time (at a census date), total days N[NNN]
<i>METeOR identifier:</i>	269961
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The time elapsed (in days) for a patient on the elective surgery waiting list from the date they were added to the waiting list to a designated census date.
<i>Context:</i>	Elective surgery

Data element concept attributes

<i>Data element concept:</i>	Elective surgery waiting list episode – waiting time
<i>Definition:</i>	The time elapsed for a patient waiting for elective surgery from the date they were added to the waiting list for the procedure.
<i>Object class:</i>	Elective surgery waiting list episode
<i>Property:</i>	Waiting time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NNN]
<i>Maximum character length:</i>	4
<i>Unit of measure:</i>	Day

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>The number of days is calculated by subtracting the Elective care waiting list episode – listing date for care, DDMMYYYY from the Hospital census (of elective surgery waitlist patients) – census date, DDMMYYYY, minus any days when the patient was 'not ready for care', and also minus any days the patient was waiting with a less urgent clinical urgency category than their clinical urgency category at the census date.</p> <p>Days when the patient was not ready for care is calculated by subtracting the date(s) the person was recorded as 'not ready for care' from the date(s) the person was subsequently recorded as again being 'ready for care'.</p> <p>If, at any time since being added to the waiting list for the elective surgical procedure, the patient has had a less urgent clinical urgency category than the category at the census date, then the number of days waited at the less urgent Elective surgery waiting list episode – clinical urgency, code N category should be subtracted from the total number of days waited.</p>
-----------------------	--

In cases where there has been only one category reassignment (i.e. to the more urgent category attached to the patient at census date) the number of days at the less urgent clinical urgency category should be calculated by subtracting the Elective care waiting list episode – listing date for care, DDMMYYYY from the Elective care waiting list episode – category reassignment date, DDMMYYYY. If the patient's clinical urgency was reclassified more than once, days spent in each period of less urgent clinical urgency than the one applying at the census date should be calculated by subtracting one Elective care waiting list episode – category reassignment date, DDMMYYYY from the subsequent Elective care waiting list episode – category reassignment date, DDMMYYYY, and then adding the days together.

When a patient is admitted from an elective surgery waiting list but the surgery is cancelled and the patient remains on or is placed back on the waiting list within the same hospital, the time waited on the list should continue. Therefore at the census date the patient's waiting time includes the number of days waited on an elective surgery waiting list, both before and after any cancelled surgery admission. The time waited before the cancelled surgery should be counted as part of the total time waited by the patient.

Comments:

Elective surgery waiting times data collections include measures of waiting times at removal and at designated census dates. This metadata item is used to measure waiting times at a designated census date whereas the metadata item Elective surgery waiting list episode – waiting time (at removal), total days N[NNN] measures waiting times at removal.

The calculation of waiting times for patients who are transferred from an elective surgery waiting list managed by one public acute hospital to another will be investigated in the future. In this case, the amount of time waited on previous lists should follow the patient to the next. Therefore at the census date, their waiting time includes the total number of days on all lists (less days not ready for care and days in lower urgency categories).

This is a critical elective surgery waiting times metadata item. It is used to determine whether patients are overdue, or had extended waits at a census date. It is used to assist doctors and patients in making decisions about hospital referral, to assist in the planning and management of hospitals and in health care related research.

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare
Origin: National Health Data Committee

Relational attributes

Related metadata references: Is formed using Elective surgery waiting list episode – patient listing status, readiness for care code N NHIG, Standard 01/03/2005
 Is formed using Hospital census (of elective surgery waitlist patients) – census date, DDMMYYYY NHIG, Standard 01/03/2005
 Is formed using Elective care waiting list episode – listing date

for care, DDMMYYYY NHIG, Standard 01/03/2005
Is formed using Elective care waiting list episode – category reassignment date, DDMMYYYY NHIG, Standard 01/03/2005
Supersedes Waiting time at a census date, version 2, Derived DE, NHDD, NHIMG, Superseded 01/03/2005
Is used in the formation of Elective surgery waiting list episode – overdue patient status, code N NHIG, Standard 01/03/2005
Is used in the formation of Elective surgery waiting list episode – extended wait patient indicator, code N NHIG, Standard 01/03/2005
Elective surgery waiting times (census data) NMDS NHIG, Standard 07/12/2005
Implementation start date: 30/09/2006
Elective surgery waiting times (census data) NMDS NHIG, Superseded 07/12/2005
Implementation start date: 30/09/2002
Implementation end date: 30/06/2006

Implementation in Data Set Specifications:

Waiting time at removal from elective surgery waiting list

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Elective surgery waiting list episode – waiting time (at removal), total days N[NNN]
<i>METeOR identifier:</i>	269960
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The time elapsed (in days) for a patient on the elective surgery waiting list from the date they were added to the waiting list for the procedure to the date they were removed from the waiting list.
<i>Context:</i>	Elective surgery

Data element concept attributes

<i>Data element concept:</i>	Elective surgery waiting list episode – waiting time
<i>Definition:</i>	The time elapsed for a patient waiting for elective surgery from the date they were added to the waiting list for the procedure.
<i>Object class:</i>	Elective surgery waiting list episode
<i>Property:</i>	Waiting time

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	N[NNN]
<i>Maximum character length:</i>	4
<i>Unit of measure:</i>	Day

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	<p>The number of days is calculated by subtracting the listing date for care from the removal date, minus any days when the patient was 'not ready for care', and also minus any days the patient was waiting with a less urgent clinical urgency category than their clinical urgency category at removal.</p> <p>Days when the patient was not ready for care is calculated by subtracting the date(s) the person was recorded as 'not ready for care' from the date(s) the person was subsequently recorded as again being 'ready for care'.</p> <p>If, at any time since being added to the waiting list for the elective surgical procedure, the patient has had a less urgent clinical urgency category than the category at removal, then the number of days waited at the less urgent clinical urgency category should be subtracted from the total number of days waited.</p> <p>In cases where there has been only one category reassignment</p>
-----------------------	--

(i.e. to the more urgent category attached to the patient at removal) the number of days at the less urgent clinical urgency category should be calculated by subtracting the listing date for care from the category reassignment date. If the patient's clinical urgency was reclassified more than once, days spent in each period of less urgent clinical urgency than the one applying at removal should be calculated by subtracting one category reassignment date from the subsequent category reassignment date, and then adding the days together.

When a patient is removed from an elective surgery waiting list, for admission on an elective basis for the procedure they were awaiting, but the surgery is cancelled and the patient remains on or is placed back on the waiting list within the same hospital, the time waited on the list should continue.

Therefore at the removal date, the patient's waiting time includes the number of days waited on an elective surgery waiting list, both before and after any cancelled surgery admission. The time waited before the cancelled surgery should be counted as part of the total time waited by the patient.

Comments:

Elective surgery waiting times data collections include measures of waiting times at removal and at designated census dates. This metadata item is used to measure waiting times at removal whereas the metadata item waiting time at a census date measures waiting times at a designated census date.

The calculation of waiting times for patients, who are transferred from an elective surgery waiting list managed by one public acute hospital to another, will be investigated in the future. In this case, the amount of time waited on previous lists would follow the patient to the next. Therefore when the patient is removed from the waiting list (for admission or other reason), their waiting time would include the total number of days on all lists (less days not ready for care and days in lower urgency categories).

This is a critical elective surgery waiting times metadata item. It is used to determine whether patients were overdue, or had extended waits when they were removed from the waiting list. It is used to assist doctors and patients in making decisions about hospital referral, to assist in the planning and management of hospitals and in health care related research.

Source and reference attributes

Submitting organisation:

Australian Institute of Health and Welfare

Origin:

National Health Data Committee

Relational attributes

Related metadata references:

Supersedes Waiting time at removal from elective surgery waiting list, version 2, Derived DE, NHDD, NHIMG, Superseded 01/03/2005

Is formed using Elective care waiting list episode – category reassignment date, DDMMYYYY NHIG, Standard 01/03/2005

Is formed using Elective surgery waiting list episode – waiting list removal date, DDMMYYYY NHIG, Standard 01/03/2005

Is formed using Episode of admitted patient care – admission date, DDMMYYYY NHIG, Standard 01/03/2005

Is formed using Elective care waiting list episode – listing date

Implementation in Data Set Specifications:

for care, DDMMYYYY NHIG, Standard 01/03/2005
Is used in the formation of Elective surgery waiting list episode – overdue patient status, code N NHIG, Standard 01/03/2005

Is used in the formation of Elective surgery waiting list episode – extended wait patient indicator, code N NHIG, Standard 01/03/2005

Elective surgery waiting times (removals data) NMDS NHIG, Standard 07/12/2005

Implementation start date: 01/07/2006

Elective surgery waiting times (removals data) NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2002

Implementation end date: 30/06/2006

Weight (self-reported)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – weight (self-reported), total kilograms NN[N]
<i>METeOR identifier:</i>	302365
<i>Registration status:</i>	NHIG, Standard 14/07/2005
<i>Definition:</i>	A person's self-reported weight (body mass).

Data element concept attributes

<i>Data element concept:</i>	Person – weight
<i>Definition:</i>	The body mass of a person.
<i>Context:</i>	Public health and health care: Weight is an overall measure of body size that does not distinguish between fat and muscle. Weight is an indicator of nutrition status and health status. Low pre-pregnancy weight is an indicator of poorer gestational outcome in women (Kramer 1988). Low weight is also associated with osteoporosis. In general, change in weight is of interest in adults because it is an indicator of changing health status. Self reported or parentally reported weight for children and adolescents should be used cautiously if at all. It enables the calculation of body mass index which requires the measurement of height and weight (body mass) for adults.
<i>Object class:</i>	Person
<i>Property:</i>	Weight

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	NN[N]
<i>Maximum character length:</i>	3
<i>Supplementary values:</i>	Value Meaning 888 Unknown 999 Not stated
<i>Unit of measure:</i>	Kilogram (Kg)

Collection and usage attributes

<i>Guide for use:</i>	CODE 888 Unknown Use this code if self-reported body mass (weight) is unknown. CODE 999 Not stated Use this code if self-reported body mass (weight) is not responded to.
-----------------------	--

Data element attributes

Collection and usage attributes

Collection methods:

The method of data collection, e.g. face to face interview, telephone interview or self-completion questionnaire, can affect survey estimates and should be reported.

The data collection form should include a question asking the respondent what their weight is. For example, the ABS National Health Survey 1989-90 included the question 'How much do you weigh without clothes and shoes?'. The data collection form should allow for both metric (to the nearest 1 kg) and imperial (to the nearest 1 lb) units to be recorded.

If practical, it is preferable to enter the raw data into the data base before conversion of measures in imperial units to metric. However, if this is not possible, weight reported in imperial units can be converted to metric prior to data entry using a conversion factor of 0.454 kg to the lb.

Rounding to the nearest 1 kg will be required for measures converted to metric prior to data entry, and may be required for data reported in metric units to a greater level of precision than the nearest 1 kg. The following rounding conventions are desirable to reduce systematic over reporting (Armitage and Berry 1994):

nnn.x where x

nnn.x where $x > 5$ - round up, e.g. 72.7 kg would be rounded to 73 kg.

nnn.x where $x = 5$ - round to the nearest even number, e.g. 72.5 kg would be rounded to 72 kg, while 73.5 kg would be rounded to 74 kg.

Comments:

This metadata item is recommended for persons aged 18 years or older. It is recommended for use in population surveys when it is not possible to measure weight.

It is recommended that in population surveys, sociodemographic data including ethnicity should be collected, as well as other risk factors including physiological status (e.g. pregnancy), physical activity, smoking and alcohol consumption. Summary statistics may need to be adjusted for these variables. Metadata items currently exist for sex, date of birth, country of birth, Indigenous status and smoking. Metadata items are being developed for physical activity.

Presentation of data:

Means and 95% confidence intervals, medians and centiles should be reported to one decimal place. Where the sample permits, population estimates should be presented by sex and 5-year age groups. Estimates based on sample surveys may need to take into account sampling weights.

For consistency with conventional practice, and for current comparability with international data sets, recommended centiles are 5, 10, 15, 25, 50, 75, 85, 90 and 95. To estimate the 5th and 95th centiles, a sample size of at least 200 is recommended for each group for which the centiles are being specified.

For some reporting purposes, it may be desirable to present weight data in categories. It is recommended that 5 kg groupings are used for this purpose. Weight data should not be rounded before categorisation. The following categories may be appropriate for describing the weights of Australian men and women, although the range will depend on the population. The

World Health Organization's range for weight is 30-140 kg.

Weight

30 kg = Weight

35 kg = Weight

... in 5 kg categories

135 kg = Weight

Weight => 140 kg

On average, body mass (weight) tends to be underestimated when self-reported by respondents. Data for men and women aged 20-69 years in 1989 indicated that men underestimated by an average of 0.2 kg (sem of 0.05 kg) and women by an average of 0.4 kg (sem of 0.04 kg) (Waters 1993). The extent of underestimation varied with age.

Source and reference attributes

Origin:

National Centre for Monitoring Cardiovascular Disease
Australian Institute of Health and Welfare
National Health Data Committee

Relational attributes

Related metadata references:

Supersedes Weight - self-reported, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Supersedes Adult – weight (self-reported), total kilograms NN[N] NHIG, Superseded 14/07/2005

Is used in the formation of Child – body mass index (self-reported), ratio NN[N].N[N] NHIG, Standard 01/03/2005

Is used in the formation of Adult – body mass index (self-reported), ratio NN[N].N[N] NHIG, Standard 01/03/2005

Implementation in Data Set Specifications:

Acute coronary syndrome (clinical) DSS NHIG, Standard 07/12/2005

Implementation start date: 07/12/2005

Acute coronary syndrome (clinical) DSS NHIG, Superseded 07/12/2005

Weight in grams (measured)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – weight (measured), total grams NNNN
<i>Synonymous names:</i>	Infant weight, neonate, stillborn
<i>METeOR identifier:</i>	310245
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The weight (body mass) of a person measured in grams.

Data element concept attributes

<i>Data element concept:</i>	Person – weight
<i>Definition:</i>	The body mass of a person.
<i>Context:</i>	Public health and health care: Weight is an overall measure of body size that does not distinguish between fat and muscle. Weight is an indicator of nutrition status and health status. Low pre-pregnancy weight is an indicator of poorer gestational outcome in women (Kramer 1988). Low weight is also associated with osteoporosis. In general, change in weight is of interest in adults because it is an indicator of changing health status. Self reported or parentally reported weight for children and adolescents should be used cautiously if at all. It enables the calculation of body mass index which requires the measurement of height and weight (body mass) for adults.
<i>Object class:</i>	Person
<i>Property:</i>	Weight

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total
<i>Data type:</i>	Number
<i>Format:</i>	NNNN
<i>Maximum character length:</i>	4
<i>Unit of measure:</i>	Gram (g)

Data element attributes

Source and reference attributes

<i>Submitting organisation:</i>	Australian Institute of Health and Welfare
---------------------------------	--

Relational attributes

<i>Related metadata references:</i>	Is used in the formation of Episode of admitted patient care – major diagnostic category, code (AR-DRG v5.1) NN NHIG, Standard 01/03/2005
	Is used in the formation of Episode of admitted patient care – diagnosis related group, code (AR-DRG v5.1) ANNA NHIG,

Implementation in Data Set Specifications:

Standard 01/03/2005

Admitted patient care NMDS NHIG, Superseded 07/12/2005

Implementation start date: 01/07/2005

Implementation end date: 30/06/2006

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this metadata item must be consistent with diagnoses and procedure codes for valid grouping.

Weight on the date the infant is admitted should be recorded if the weight is less than or equal to 9000g and age is less than 365 days.

Admitted patient care NMDS 2007-2008 NHIG, Standard 29/11/2006

Implementation start date: 01/07/2007

Information specific to this data set:

For the provision of state and territory hospital data to Commonwealth agencies this metadata item must be consistent with diagnoses and procedure codes for valid grouping.

Weight on the date the infant is admitted should be recorded if the weight is less than or equal to 9000g and age is less than 365 days.

Weight in kilograms (measured)

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person – weight (measured), total kilograms N[NN].N
<i>Synonymous names:</i>	Infant weight, neonate, stillborn
<i>METeOR identifier:</i>	270208
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The weight (body mass) of a person measured in kilograms.

Data element concept attributes

<i>Data element concept:</i>	Person – weight
<i>Definition:</i>	The body mass of a person.
<i>Context:</i>	Public health and health care: Weight is an overall measure of body size that does not distinguish between fat and muscle. Weight is an indicator of nutrition status and health status. Low pre-pregnancy weight is an indicator of poorer gestational outcome in women (Kramer 1988). Low weight is also associated with osteoporosis. In general, change in weight is of interest in adults because it is an indicator of changing health status. Self reported or parentally reported weight for children and adolescents should be used cautiously if at all. It enables the calculation of body mass index which requires the measurement of height and weight (body mass) for adults.
<i>Object class:</i>	Person
<i>Property:</i>	Weight

Value domain attributes

Representational attributes

<i>Representation class:</i>	Total				
<i>Data type:</i>	Number				
<i>Format:</i>	N[NN].N				
<i>Maximum character length:</i>	4				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>999.9</td><td>Not collected</td></tr></tbody></table>	Value	Meaning	999.9	Not collected
Value	Meaning				
999.9	Not collected				
<i>Unit of measure:</i>	Kilogram (Kg)				

Collection and usage attributes

<i>Guide for use:</i>	A continuous variable measured to the nearest 0.1 kg. CODE 999.9 Not collected Use this code if measured weight is not collected.
-----------------------	---

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	In order to ensure consistency in measurement, the
-----------------------	--

Collection methods:

measurement protocol described under Collection methods should be used.

The collection of anthropometric measurements, particularly in those who are overweight or obese or who are concerned about their weight, should be performed with great sensitivity and without drawing attention to an individual's weight.

The measurement protocol described below is that recommended by the WHO Expert Committee (1995).

Measurement protocol:

Equipment used should be described and reported. Scales should have a resolution of at least 0.1kg and should have the capacity to weigh up to at least 200 kg. Measurement intervals and labels should be clearly readable under all conditions of use of the instrument. Scales should be capable of being calibrated across the entire range of measurements. Precision error should be no more than 0.1kg. Scales should be calibrated on each day of use. Manufacturers' guidelines should be followed with regard to the transportation of the scales.

Adults and children who can stand:

The subject stands over the centre of the weighing instrument, with the body weight evenly distributed between both feet.

Heavy jewellery should be removed and pockets emptied.

Light indoor clothing can be worn, excluding shoes, belts, and sweater. Any variations from light indoor clothing (e.g. heavy clothing, such as kaftans or coats worn because of cultural practices) should be noted on the data collection form.

Adjustments for non-standard clothing (i.e. other than light indoor clothing) should only be made in the data checking/cleaning stage prior to data analysis.

If the subject has had one or more limbs amputated, record this on the data collection form and weigh them as they are. If they are wearing an artificial limb, record this on the data collection form but do not ask them to remove it. Similarly, if they are not wearing the limb, record this but do not ask them to put it on.

The measurement is recorded to the nearest 0.1 kg. If the scales do not have a digital readout, take a repeat measurement. If the two measurements disagree by more than 0.5 kg, then take a third measurement. All raw measurements should be recorded on the data collection form. If practical, it is preferable to enter the raw data into the database as this enables intra-observer and, where relevant, inter-observer errors to be assessed. The subject's measured weight is subsequently calculated as the mean of the two observations, or the mean of the two closest measurements if a third is taken, and recorded on the form. If only a mean value is entered into the database then the data collection forms should be retained.

It may be necessary to round the mean value to the nearest 0.1 kg. If so, rounding should be to the nearest even digit to reduce systematic over reporting (Armitage and Berry 1994). For example, a mean value of 72.25 kg would be rounded to 72.2 kg, while a mean value of 72.35 kg would be rounded to 72.4 kg.

Infants:

Birth weight and gender should be recorded with gestational age. During infancy a levelled pan scale with a bean and movable weights or digital scales capable of measuring to two decimal places of a kilogram are acceptable. Birth weight

should be determined within 12 hours of birth. The infant, with or without a nappy or diaper is placed on the scales so that the weight is distributed equally about the centre of the pan. When the infant is lying or suspended quietly, weight is recorded to the nearest 10 grams. If the nappy or diaper is worn, its weight is subtracted from the observed weight i.e. reference data for infants are based on nude weights.

Validation and quality control measures:

If practical, equipment should be checked daily using one or more objects of known weight in the range to be measured. It is recommended that the scale be calibrated at the extremes and in the mid range of the expected weight of the population being studied.

Within- and, if relevant, between-observer variability should be reported. They can be assessed by the same (within-) or different (between-) observers repeating the measurement of weight, on the same subjects, under standard conditions after a short time interval. The standard deviation of replicate measurements (technical error of measurement) between observers should not exceed 0.5 kg and be less than 0.5 kg within observers.

Extreme values at the lower and upper end of the distribution of measured height should be checked both during data collection and after data entry. Individuals should not be excluded on the basis of true biological difference.

Last digit preference, and preference or avoidance of certain values, should be analysed in the total sample and (if relevant) by observer, survey site and over time if the survey period is long.

Comments:

This metadata item applies to persons of all ages. It is recommended for use in population surveys and health care settings.

It is recommended that in population surveys, sociodemographic data including ethnicity should be collected, as well as other risk factors including physiological status (e.g. pregnancy), physical activity, smoking and alcohol consumption. Summary statistics may need to be adjusted for these variables.

Metadata items currently exist for sex, date of birth, country of birth, Indigenous status and smoking. Metadata items are being developed for physical activity.

Presentation of data:

Means and 95% confidence intervals, medians and centiles should be reported to one decimal place. Where the sample permits, population estimates should be presented by sex and 5-year age groups. However 5-year age groups are not generally suitable for children and adolescents. Estimates based on sample surveys may need to take into account sampling weights.

For consistency with conventional practice, and for current comparability with international data sets, recommended centiles are 5, 10, 15, 25, 50, 75, 85, 90 and 95. To estimate the 5th and 95th centiles, a sample size of at least 200 is recommended for each group for which the centiles are being specified.

For some reporting purposes, it may be desirable to present weight data in categories. It is recommended that 5 kg

groupings are used for this purpose. Weight data should not be rounded before categorisation. The following categories may be appropriate for describing the weights of Australian men, women, children and adolescents, although the range will depend on the population.

Weight

10 kg = Weight

15 kg = Weight

... in 5 kg categories

135 kg = Weight

Weight => 140 kg

Source and reference attributes

Submitting organisation:

World Health Organization The consortium to develop standard methods for the collection and collation of anthropometric data in children as part of the National Food and Nutrition Monitoring and Surveillance Project, funded by the Commonwealth Department of Health and Ageing

Reference documents:

Clinical Guidelines on the Identification, Evaluation and Treatment of Overweight and Obesity in Adults (US National Heart, Lung and Blood Institute (NHLBI) in cooperation with the National Institute of Diabetes and Digestive and Kidney Diseases).

Chronic Diseases and Associated Risk Factors in Australia 2001 (AIHW).

Relational attributes

Related metadata references:

Supersedes Weight - measured, version 2, DE, NHDD, NHIMG, Superseded 01/03/2005

Is used in the formation of Child – body mass index (self-reported), ratio NN[N].N[N] NHIG, Standard 01/03/2005

Is used in the formation of Child – body mass index (measured), ratio NN[N].N[N] NHIG, Standard 01/03/2005

Is used in the formation of Adult – body mass index (self-reported), ratio NN[N].N[N] NHIG, Standard 01/03/2005

Is used in the formation of Adult – body mass index (measured), ratio NN[N].N[N] NHIG, Standard 01/03/2005

Implementation in Data Set Specifications:

Cardiovascular disease (clinical) DSS NHIG, Superseded 15/02/2006

Cardiovascular disease (clinical) DSS NHIG, Standard 15/02/2006

Diabetes (clinical) DSS NHIG, Superseded 21/09/2005

Information specific to this data set:

Following Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus, body mass index (BMI) should be below 27 kg/m² for men and women. For adults who suffer from diabetes, the recommendation is to measure weight and calculate BMI on the initial visit and then measure weight every 3 months. If the patient is on a weight reduction program, weight is to be measured more frequently.

Strong evidence exists that weight loss reduces blood pressure in both overweight hypertensive and non-

hypertensive individuals; reduces serum triglycerides and increases high-density lipoprotein (HDL)-cholesterol; and generally produces some reduction in total serum cholesterol and low-density lipoprotein (LDL)-cholesterol. The risk of developing diabetes rises continuously with increasing obesity (DHAC & AIHW 1999:13). An increased central distribution of body fat (when fatness is concentrated in the abdomen) also appears to be associated more often with Type 2 diabetes (Bishop et al. 1998:430-1).

Weight loss reduces blood glucose levels in overweight and obese persons with and without diabetes; and weight loss also reduces blood glucose levels and HbA1c in some patients with type 2 diabetes. Although there have been no prospective trials to show changes in mortality with weight loss in obese patients, reductions in risk factors would suggest that development of type 2 diabetes and CVD would be reduced with weight loss.

Diabetes (clinical) DSS NHIG, Standard 21/09/2005

Information specific to this data set:

Following Principles of Care and Guidelines for the Clinical Management of Diabetes Mellitus, body mass index (BMI) should be below 27 kg/m² for men and women. For adults who suffer from diabetes, the recommendation is to measure weight and calculate BMI on the initial visit and then measure weight every 3 months. If the patient is on a weight reduction program, weight is to be measured more frequently.

Strong evidence exists that weight loss reduces blood pressure in both overweight hypertensive and non-hypertensive individuals; reduces serum triglycerides and increases high-density lipoprotein (HDL)-cholesterol; and generally produces some reduction in total serum cholesterol and low-density lipoprotein (LDL)-cholesterol. The risk of developing diabetes rises continuously with increasing obesity (DHAC & AIHW 1999:13). An increased central distribution of body fat (when fatness is concentrated in the abdomen) also appears to be associated more often with Type 2 diabetes (Bishop et al. 1998:430-1).

Weight loss reduces blood glucose levels in overweight and obese persons with and without diabetes; and weight loss also reduces blood glucose levels and HbA1c in some patients with type 2 diabetes. Although there have been no prospective trials to show changes in mortality with weight loss in obese patients, reductions in risk factors would suggest that development of type 2 diabetes and CVD would be reduced with weight loss.

Weight is an overall measure of body size that does not distinguish between fat and muscle. Weight is an indicator of nutritional and health status. Low pre-pregnancy weight is an indicator of poorer gestational outcome in women (Kramer 1988). Low weight is also associated with osteoporosis. In general, change in weight in adults is of interest because it is an indicator of changing health status,

and in children as it indicates changing health status and growth and development. Self-reported or parentally-reported weight for children and adolescents should be used cautiously if at all. It enables the calculation of body mass index (BMI) which requires the measurement of height and weight for adults as well as sex and date of birth for children and adolescents.

Year insulin started

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Patient – insulin start date, YYYY
<i>METeOR identifier:</i>	269928
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The year the patient started insulin injections.
<i>Context:</i>	Public health, health care and clinical settings.

Data element concept attributes

<i>Data element concept:</i>	Patient – insulin start date
<i>Definition:</i>	The date the patient started insulin injections.
<i>Object class:</i>	Patient
<i>Property:</i>	Insulin start date

Source and reference attributes

<i>Submitting organisation:</i>	National Diabetes Data Working Group
---------------------------------	--------------------------------------

Value domain attributes

Representational attributes

<i>Representation class:</i>	Date
<i>Data type:</i>	Date/Time
<i>Format:</i>	YYYY
<i>Maximum character length:</i>	4

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record the year that insulin injections were started. This data element has to be completed for all patients who use insulin. It is used to cross check diabetes type assignment.
<i>Collection methods:</i>	Ask the individual the year when he/ she started to use insulin. Alternatively obtain this information from appropriate documentation, if available.

Source and reference attributes

<i>Submitting organisation:</i>	National diabetes data working group
<i>Origin:</i>	National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary

Relational attributes

<i>Related metadata references:</i>	Supersedes Year insulin started, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Diabetes (clinical) DSS NHIG, Superseded 21/09/2005 Diabetes (clinical) DSS NHIG, Standard 21/09/2005

Information specific to this data set:

This data element provides information about the duration of diabetes in individual patients.

Insulin is a regulating hormone secreted into the blood in response to a rise in concentration of blood glucose or amino acids. It is a double-chain protein hormone formed from proinsulin in the beta cells of the pancreatic islets of Langerhans. Insulin promotes the storage of glucose and the uptake of amino acids, increases protein and lipid synthesis, and inhibits lipolysis and gluconeogenesis.

Commercially prepared insulin is available in various types, which differ in the speed they act and in the duration of their effectiveness.

Year of arrival in Australia

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Person (overseas born) – year of first arrival in Australia, date YYYY
<i>METeOR identifier:</i>	269929
<i>Registration status:</i>	NHIG, Standard 04/05/2005 NCSIMG, Standard 01/03/2005
<i>Definition:</i>	The year a person (born outside of Australia) first arrived in Australia, from another country, with the intention of staying in Australia for one year or more.

Data element concept attributes

<i>Data element concept:</i>	Person (overseas born) – year of first arrival in Australia
<i>Definition:</i>	The year a person (born outside of Australia) first arrived in Australia, from another country, with the intention of staying in Australia for one year or more.
<i>Context:</i>	Client demographic and support needs: This metadata item is an important indicator of cultural identity, and provides further information about the person's cultural values and practices which is significant in determining service type and support required. May also be used in conjunction with other socio-demographic data to indicate the background of the person, communication skills and their possible needs.
<i>Object class:</i>	Person
<i>Property:</i>	Year of first arrival in Australia

Value domain attributes

Representational attributes

<i>Representation class:</i>	Date
<i>Data type:</i>	Date/Time
<i>Format:</i>	YYYY
<i>Maximum character length:</i>	4

Data element attributes

Collection and usage attributes

<i>Collection methods:</i>	Actual year of arrival in Australia. Recommended question: In what year did you/the person first arrive in Australia to live here for one year or more? (Write in the calendar year of arrival or mark the box if here less than one year) Calendar year of arrival Will be here less than one year It is anticipated that for the majority of people their response to
----------------------------	---

the question will be the year of their only arrival in Australia. However, some respondents may have multiple arrivals in Australia. To deal with these cases in self-enumerated collections, an instruction such as 'Please indicate the year of first arrival only' should be included with the question.

While agencies are encouraged to use the recommended question described above, it is acknowledged that this is not always possible in practice. For example, where the data collection is a by-product of the provision of a health or community service, the information may be ascertained using different means. However, the recommended question should be used wherever practically possible.

Source and reference attributes

Origin: The Australian Bureau of Statistics Standard for Year of Arrival in Australia. (last viewed 05/12/2006)

Reference documents: The ABS standard for the collection of Year of arrival in Australia appears on the ABS Website [http://www.abs.gov.au/ausstats/abs@.nsf/StatsLibrary select Other ABS Statistical Standards/Standards for Social, Labour and Demographic Variables/Cultural Diversity Variable](http://www.abs.gov.au/ausstats/abs@.nsf/StatsLibrary%20select%20Other%20ABS%20Statistical%20Standards/Standards%20for%20Social,%20Labour%20and%20Demographic%20Variables/Cultural%20Diversity%20Variable).

Relational attributes

Related metadata references: Supersedes Year of arrival in Australia, version 2, DE, NCSDD, NCSIMG, Superseded 01/03/2005

Implementation in Data Set Specifications: Computer Assisted Telephone Interview demographic module DSS NHIG, Standard 04/05/2005

Year of diagnosis of diabetes mellitus

Identifying and definitional attributes

<i>Metadata item type:</i>	Data Element
<i>Technical name:</i>	Patient – diagnosis date (diabetes mellitus), YYYY
<i>METeOR identifier:</i>	269930
<i>Registration status:</i>	NHIG, Standard 01/03/2005
<i>Definition:</i>	The year a patient was first diagnosed as having diabetes
<i>Context:</i>	Public health, health care and clinical settings.

Data element concept attributes

<i>Data element concept:</i>	Patient – diagnosis date
<i>Definition:</i>	The date on which a patient is diagnosed with a particular condition or disease.
<i>Object class:</i>	Patient
<i>Property:</i>	Diagnosis date

Value domain attributes

Representational attributes

<i>Representation class:</i>	Date				
<i>Data type:</i>	Date/Time				
<i>Format:</i>	YYYY				
<i>Maximum character length:</i>	4				
<i>Supplementary values:</i>	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>9999</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	9999	Not stated/inadequately described
Value	Meaning				
9999	Not stated/inadequately described				

Data element attributes

Collection and usage attributes

<i>Guide for use:</i>	Record the year that the patient was first diagnosed as having diabetes.
<i>Collection methods:</i>	Ask the individual the year when he/ she was diagnosed with diabetes. Alternatively obtain this information from appropriate documentation, if available.

Source and reference attributes

<i>Submitting organisation:</i>	National diabetes data working group
<i>Origin:</i>	National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary

Relational attributes

<i>Related metadata references:</i>	Supersedes Year of diagnosis of diabetes mellitus, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005
<i>Implementation in Data Set Specifications:</i>	Diabetes (clinical) DSS NHIG, Superseded 21/09/2005 Diabetes (clinical) DSS NHIG, Standard 21/09/2005
	<i>Information specific to this data set:</i>

Long-term complications of diabetes mellitus affect the eyes, kidneys, nerves, and blood vessels.