

Person—timing of angiotensin II receptor blocker prescription, code N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Timing of angiotensin II receptor blocker prescription
METEOR identifier:	350421
Registration status:	<ul style="list-style-type: none">• Health, Standard 01/10/2008
Definition:	The timing of care when an angiotensin II receptor blocker is prescribed to a person, as represented by a code.

Data element concept attributes

Identifying and definitional attributes

Data element concept:	Person—timing of angiotensin II receptor blocker prescription
METEOR identifier:	350416
Registration status:	<ul style="list-style-type: none">• Health, Standard 01/10/2008
Definition:	The timing of care when an angiotensin II receptor blocker is prescribed to a person.
Object class:	Person
Property:	Timing of angiotensin II receptor blocker prescription

Value domain attributes

Identifying and definitional attributes

Value domain:	Timing of prescription N
METEOR identifier:	350350
Registration status:	<ul style="list-style-type: none">• Health, Standard 01/10/2008
Definition:	A code set representing the timing of a prescription.

Representational attributes

Representation class: Code
Data type: Number
Format: N
Maximum character length: 1
Permissible values:

Value	Meaning
1	Prior to presentation at hospital
2	First 24 hours of presentation
3	After 24 hours and before discharge
4	At discharge

Supplementary values:

Value	Meaning
9	Not stated/inadequately described

Data element attributes

Collection and usage attributes

Guide for use:

CODE 1 Prior to presentation at hospital

Use this code when the person has been previously prescribed an angiotensin II receptor blocker prior to presentation at the hospital and the person is still following the prescription.

CODE 2 First 24 hours of presentation

Use this code when an angiotensin II receptor blocker is prescribed within the first 24 hours following presentation to the hospital.

CODE 3 After 24 hours and before discharge

Use this code when an angiotensin II receptor blocker is prescribed following the first 24 hours after presentation to the hospital and before discharge from the hospital.

CODE 4 At discharge

Use this code when an angiotensin II receptor blocker is prescribed at discharge from the hospital.

Collection methods:

Record each time an angiotensin II receptor blocker is prescribed for the person.

Relational attributes

Implementation in Data Set Specifications:

[Acute coronary syndrome pharmacotherapy data cluster Health](#), Standard 01/10/2008

Conditional obligation:

If prescribed, provide a phase for each time angiotensin II receptor blocker therapy is prescribed.

© Australian Institute of Health and Welfare 2015–2022

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 3.0 (CC BY 3.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build upon this work. However, you must attribute the AIHW as the copyright holder of the work in compliance with our attribution policy available at www.aihw.gov.au/copyright. The full terms and conditions of this licence are available at <http://creativecommons.org/licenses/by3.0/au/>.

Enquiries relating to copyright should be addressed to the Head of the Communications, Media and Marketing Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.