

Person—timing of antithrombin therapy prescription, code N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Timing of antithrombin therapy prescription
METEOR identifier:	350510
Registration status:	<ul style="list-style-type: none">• Health, Standard 01/10/2008
Definition:	The timing of care when antithrombin therapy is prescribed to a person, as represented by a code.
Data Element Concept:	Person—timing of antithrombin therapy prescription

Value domain attributes

Representational attributes

Representation class:	Code								
Data type:	Number								
Format:	N								
Maximum character length:	1								
Permissible values:	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>1</td><td>Initial medical management: preceding reperfusion therapy</td></tr><tr><td>2</td><td>During reperfusion therapy</td></tr><tr><td>3</td><td>Following reperfusion therapy</td></tr></tbody></table>	Value	Meaning	1	Initial medical management: preceding reperfusion therapy	2	During reperfusion therapy	3	Following reperfusion therapy
Value	Meaning								
1	Initial medical management: preceding reperfusion therapy								
2	During reperfusion therapy								
3	Following reperfusion therapy								
Supplementary values:	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>9</td><td>Not stated/inadequately described</td></tr></tbody></table>	Value	Meaning	9	Not stated/inadequately described				
Value	Meaning								
9	Not stated/inadequately described								

Source and reference attributes

Reference documents:	National Heart Foundation of Australia and Cardiac Society of Australia and New Zealand, Guidelines for the management of acute coronary syndromes 2006, Med J Aust; 184; S1-S32. © MJA2006.
-----------------------------	--

Data element attributes

Collection and usage attributes

Guide for use:

Reperfusion therapy includes percutaneous coronary intervention and fibrinolytic therapy.

CODE 1 Initial medical management: preceding reperfusion therapy

Use this code when antithrombin therapy is prescribed before reperfusion therapy is to be performed.

CODE 2 During reperfusion therapy

Use this code when antithrombin therapy is prescribed while reperfusion therapy is being performed.

CODE 3 Following reperfusion therapy

Use this code when antithrombin therapy is prescribed after reperfusion therapy has been performed.

Collection methods:

Record for each time antithrombin therapy is prescribed for the person.

Relational attributes

Implementation in Data Set Specifications:

[Acute coronary syndrome pharmacotherapy data cluster Health](#), Standard 01/10/2008

Conditional obligation:

If prescribed, provide a phase for each time antithrombin therapy is prescribed