
# Person—revascularisation percutaneous coronary



#### © Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

# Person—revascularisation percutaneous coronary intervention date, DDMMYYYY

# Identifying and definitional attributes

Metadata item type: Data Element

**Short name:** Date of revascularisation percutaneous coronary intervention

**Synonymous names:** Revascularisation PCI date

METEOR identifier: 359731

Registration status: Health, Standard 01/10/2008

**Definition:** The date when a percutaneous coronary intervention (PCI) is performed for

revascularisation.

Data Element Concept: Person—revascularisation percutaneous coronary intervention date

Value Domain: <u>Date DDMMYYYY</u>

### Value domain attributes

## Representational attributes

Representation class: Date

Data type: Date/Time
Format: DDMMYYYY

Maximum character length: 8

#### Source and reference attributes

**Submitting organisation:** Australian Institute of Health and Welfare

#### Data element attributes

#### Collection and usage attributes

Guide for use: Revascularisation PCI relates to balloon angioplasty inflation and/or stent

implantation performed for subsequent restoration of blood flow.

**Comments:** Routine revascularisation PCI may be performed after ST-segment-elevation

myocardial infarction for people with objective evidence of recurrent myocardial infarction in whom there is spontaneous or inducible ischaemia or haemodynamic instability. Revascularisation PCI may also be performed for treatment of high-risk

non-ST-segment-elevation acute coronary syndrome.

#### Source and reference attributes

**Reference documents:** National Heart Foundation of Australia & Cardiac Society of Australia and New

Zealand. Guidelines for the management of acute coronary syndromes 2006. Med

J Aust 2006; 184; S1-S32. © MJA 2006

#### Relational attributes

Implementation in Data Set Coronary artery cluster

Specifications: Health, Standard 01/10/2008

Conditional obligation: Record when a percutaneous coronary intervention is

performed for revascularisation.