

Person—revascularisation percutaneous coronary intervention date, DDMMYYYY

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Person—revascularisation percutaneous coronary intervention date, DDMMYYYY

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Date of revascularisation percutaneous coronary intervention
Synonymous names:	Revascularisation PCI date
METEOR identifier:	359731
Registration status:	Health , Standard 01/10/2008
Definition:	The date when a percutaneous coronary intervention (PCI) is performed for revascularisation.
Data Element Concept:	Person—revascularisation percutaneous coronary intervention date
Value Domain:	Date DDMMYYYY

Value domain attributes

Representational attributes

Representation class:	Date
Data type:	Date/Time
Format:	DDMMYYYY
Maximum character length:	8

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare

Data element attributes

Collection and usage attributes

Guide for use:	Revascularisation PCI relates to balloon angioplasty inflation and/or stent implantation performed for subsequent restoration of blood flow.
Comments:	Routine revascularisation PCI may be performed after ST-segment-elevation myocardial infarction for people with objective evidence of recurrent myocardial infarction in whom there is spontaneous or inducible ischaemia or haemodynamic instability. Revascularisation PCI may also be performed for treatment of high-risk non-ST-segment-elevation acute coronary syndrome.

Source and reference attributes

Reference documents: National Heart Foundation of Australia & Cardiac Society of Australia and New Zealand. Guidelines for the management of acute coronary syndromes 2006. Med J Aust 2006; 184; S1-S32. © MJA 2006

Relational attributes

Implementation in Data Set Specifications: [Coronary artery cluster](#)
[Health](#), Standard 01/10/2008
Conditional obligation: Record when a percutaneous coronary intervention is performed for revascularisation.

