National Healthcare Agreement: P20-Potentially avoidable deaths, 2010

Exported from METEOR (AIHW's Metadata Online Registry)		
© Australian Institute of Health and Welfare 2024		
This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have		

made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

National Healthcare Agreement: P20-Potentially avoidable deaths, 2010

Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

Short name: Potentially avoidable deaths, 2010

METEOR identifier: 394495

Registration status: Health, Superseded 08/06/2011

Description: Deaths that are potentially avoidable within the present health system:

potentially preventable deaths are those amenable to screening and primary

prevention such as immunisation

• potentially treatable deaths are those amenable to therapeutic interventions

Indicator set: National Healthcare Agreement (2010)

Health, Superseded 08/06/2011

Outcome area: Primary and Community Health

Health, Standard 07/07/2010

Data quality statement: National Healthcare Agreement: P20-Potentially avoidable deaths, 2010 QS

Health, Superseded 08/06/2011

Collection and usage attributes

Population group age to: Less than 75 years

Computation description: Deaths are defined as avoidable in the context of the present health system.

Rates are directly age-standardised to the Australian population as at 30 June

2001.

Presented per 100,000 population.

ICD-10 codes in scope are as specified below:

Table 20A: ICD-10 codes for potentially avoidable deaths, by major cause of death group

Cause of death group	ICD-10 Codes	Limits(age, sex)	Treatable	Preventable
Infections				
Tuberculosis	A15–A19, B90		Х	
Selected invasive bacterial and protozoal infections	A38-A41, A46, A48.1 B50-B54, G00,G03, J02.0, J13- J15, J18, L03		X	
Hepatitis	B15-B19			Х
HIV/AIDS	B20-B24			Х
Viral pneumonia and influenza	J10, J12, J17.1, J21			Х
Enteritis and other diarrhoeal disease	A00–A09			Х

Childhood vaccine- preventable diseases	A35–A37, A49.1, A49.2, A80, B01, B05– B06, J11	0–14		X
Sexually transmitted diseases, except HIV/AIDS	A50-A64, M02.3, N34.1, N70- N73, N75.0, N75.1, N76.4, N76.6, O00			X
Ear infections—otitis media and mastoiditis	H65-H70		X	
Upper respiratory tract infection	J00-J06		X	
Cancer				
Lip, oral cavity and pharynx	C00-C14			X
Oesophagus	C15			Х
Stomach	C16			Х
Colorectal	C18-C21		Х	
Liver	C22			Х
Lung	C33, C34			Х
Melanoma of skin	C43		Х	
Non-melanocytic skin	C44		Х	
Breast	C50	Female	Х	
Cervix	C53		Х	
Uterus	C54, C55		Х	
Bladder	C67		Х	
Thyroid	C73		Х	
Hodgkin's disease	C81		Х	
Lymphoid leukaemia— acute/chronic	C91.0, C91.1		X	
Benign cancers	D10-D36		Х	
Malignant neoplasm of testis	C62			Х
Malignant neoplasm of eye and adnexa	C69			X
Diabetes	E10-E14		X (0.5)	X (0.5)
Drug use disorders				
Alcohol related disease	F10, I42.6, K29.2, K70			X
Illicit drug use disorders	F11–F16, F18, F19			Х
Epilepsy	G40, G41		Х	
Diseases of the circulatory system				
Rheumatic and other valvular heart disease	101–109		Х	

Hypertensive heart disease	l11		Х	
Ischaemic heart disease	I20–I25		X (0.5)	X (0.5)
Cerebrovascular diseases	l60–l69		X (0.5)	X (0.5)
Aortic aneurysm	l71			Х
Phlebitis and thrombophlebitis of other deep vessels of lower extremities	180.2			
Pulmonary embolism	126			X
Diseases of the genitourinary system				
Nephritis and nephrosis	I12, I13, N00–N09, N17–N19		X	
Obstructive uropathy & prostatic hyperplasia	N13, N20, N21, N35, N40, N99.1		Х	
Diseases of the respiratory system				
COPD	J40–J44	45–74 years		X
Asthma	J45, J46	0-44 years	Х	
Diseases of the digestive system				
Peptic ulcer disease	K25–K28		Х	
Diseases of appendix; hernia; disorders of gallbladder, biliary tract, and pancreas	K35–K38, K40–K46, K80–K83, K85, K86, K91.5		X	
Chronic liver disease (excluding alcohol-related disease)	K73, K74			Х
Maternal & infant causes				
Birth defects	H31.1, P00, P04, Q00– Q99		X	
Complications of perinatal period	P03, P05– P95		X	
Unintentional injuries				
Road traffic injuries	V01–V04, V06, V09– V80, V87, V89, V99			Х
Falls	W00-W19			Х
Fires, burns	X00-X09			Х
Accidental poisonings	X40–X49			X
Drownings	W65–W74			X
War	Y36			Х

Intentional injuries			
Suicide and self-inflicted injuries	X60-X84, Y87.0, Y10- Y34		х
Violence	X85–Y09, Y87.1		Х
Other conditions			
Nutritional deficiency anaemia	D50-D53		Х
Thyroid disorders	E00-E07	Х	
Adrenal disorders	E24, E27	Х	
Congenital metabolic disorders	E25, E70.0, E74.2	X	
Osteomyelitis and other osteopathies of bone	M86, M89– M90		X
Complications of pregnancy, labour or the puerperium	O01–O99		Х
Misadventures to patients during surgical and medical care	Y60-Y69		Х
Medical devices associated with adverse incidents in diagnostic & therapeutic use	Y70-Y82		Х
Surgical and other medical procedures as the cause of abnormal reaction of the patient, or of later complication, without mention of misadventure at the time of the procedure	Y83–Y84		X

Computation: 100,000 x (Numerator ÷ Denominator)

Calculated separately for 'Preventable' and 'Treatable' categories, as per the table

above.

Numerator: Number of deaths of persons aged less than 75 years, categorised as potentially

avoidable

Numerator data elements: - Data El

Data Element / Data Set-

Person—underlying cause of death, code (ICD-10 2nd edn) ANN-ANN

Data Source

ABS Causes of Death Collection

Guide for use

Data source type: Registry

ICD-10 codes N99.1 and K91.5 cannot be used as underlying cause of death

Denominator: Population aged less than 75 years

Denominator data elements:

Data Element / Data Set

Data Element

Person—estimated resident population of Australia

Data Source

ABS Estimated resident population (total population)

Guide for use

Data source type: Census-based plus administrative by-product data

Data Element / Data Set-

Data Element

Person—estimated resident population of Australia

Data Source

ABS Indigenous experimental estimates and projections (2001 Censusbased)

Guide for use

Data source type: Census-based plus administrative by-product data

Data Element / Data Set-

Person—age, total years N[NN]

Data Source

ABS Estimated resident population (total population)

Guide for use

Data source type: Census-based plus administrative by-product data

Data Element / Data Set-

Person—age, total years N[NN]

Data Source

ABS Indigenous experimental estimates and projections (2001 Censusbased)

Guide for use

Data source type: Census-based plus administrative by-product data

Disaggregation data elements:

Data Element / Data Set-

Person-Indigenous status, code N

Data Source

ABS Causes of Death Collection

Guide for use

Data source type: Registry

Data Element / Data Set

Person—area of usual residence, geographical location code (ASGC 2007) NNNNN

Data Source

ABS Causes of Death Collection

Guide for use

Data source type: Registry

Used for disaggregation by state/territory, remoteness area and SEIFA of

residence

Comments:

Specified disaggregation: Nationally and by state/territory: by Indigenous status,

remoteness area and SEIFA of residence

Available disaggregation: Nationally and by state/territory: by Indigenous status

Data for 2008-09 will be available mid-2010 (2008 calendar year).

Most recent data available for 2010 CRC baseline report: 2007 (pending release

of ABS data).

Representational attributes

Representation class: Rate

Data type: Real

Unit of measure: Person

Format: NN[N].N

Indicator conceptual framework

Framework and dimensions:

Deaths

Data source attributes

Data sources:

Data Source

ABS Estimated resident population (total population)

Frequency

Quarterly

Data quality statement

ABS Estimated resident population (total population), QS

Data custodian

Australian Bureau of Statistics

Data Source

ABS Causes of Death Collection

Frequency

Annual

Data quality statement

ABS causes of death collection, QS

Data custodian

Australian Bureau of Statistics

Data Source

ABS Indigenous experimental estimates and projections (2001 Censusbased)

Frequency

Periodic

Data quality statement

ABS Indigenous experimental estimates and projections, QS

Data custodian

Australian Bureau of Statistics

Accountability attributes

Reporting requirements: National Healthcare Agreement

Organisation responsible for providing data:

Australian Bureau of Statistics (ABS) to provide the data. Data quality statement to be prepared by ABS in consultation with the Australian Institute of Health and

Welfare.

Further data development / collection required:

Specification: Interim

International work is currently being undertaken to refine and standardise the definition of avoidable mortality.

The specifications for this indicator could be modified following completion of the international work; for example, considering the prevalence of particular conditions, any advances in the effectiveness of healthcare and other interventions to prevent or treat them.

Other issues caveats:

Disaggregations within individual jurisdictions are subject to data quality considerations. Some disaggregations may result in numbers too small for publication.

Theoretically, this indicator could be adjusted for levels of community morbidity and risk factor prevalence. Or, information on risk factor prevalence and community morbidity could be used to assist interpretation of the indicator. There are currently no methods developed to undertake such adjustments.

The 75 years age cut-off is somewhat arbitrary, and will have different effects on different causes of death. For example, most car accident deaths will be included, but many hip fracture deaths will not be included.

Interpreting a change or difference in a mortality rate is problematic, as the change or difference could reflect variation in disease incidence, survival (case fatality) or both.

Relational attributes

Related metadata references:

Has been superseded by National Healthcare Agreement: PI 20-Potentially avoidable deaths, 2011

Health, Superseded 31/10/2011

See also National Healthcare Agreement: P05-Proportion of persons obese, 2010 Health, Superseded 08/06/2011

See also National Healthcare Agreement: P06-Proportion of adults who are daily smokers, 2010

Health, Superseded 08/06/2011

See also <u>National Healthcare Agreement: P07-Proportion of adults at risk of long-term harm from alcohol, 2010</u>

Health, Superseded 08/06/2011

See also National Healthcare Agreement: P09-Immunisation rates for vaccines in the national schedule, 2010

Health, Superseded 08/06/2011

See also National Healthcare Agreement: P18-Life expectancy, 2010 Health, Superseded 08/06/2011

See also National Healthcare Agreement: P19-Infant/young child mortality rate, 2010

Health, Superseded 08/06/2011

See also <u>National Healthcare Agreement</u>: <u>P22-Selected potentially preventable</u> hospitalis<u>ations</u>, <u>2010</u>

Health, Superseded 08/06/2011

See also National Healthcare Agreement: P43-Unplanned/unexpected readmissions within 28 days of selected surgical admissions, 2010

Health, Superseded 08/06/2011

See also <u>National Healthcare Agreement: P59-Age-standardised mortality by major cause of death, 2010</u>

Health, Superseded 08/06/2011