National Indigenous Reform Agreement: P06-Levels of obesity - Body Mass Index, 2010 QS

Exported from METEOR (AIHW's Metadata Online Registry)
© Australian Institute of Health and Welfare 2024
This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have

made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

National Indigenous Reform Agreement: P06-Levels of obesity - Body Mass Index, 2010 QS

Identifying and definitional attributes

Metadata item type: Data Quality Statement

METEOR identifier: 396230

Registration status: Community Services (retired), Superseded 04/04/2011

Data quality

Institutional environment: These surveys are conducted under the Census and Statistics Act 1905. For

information on the institutional environment of the Australian Bureau of Statistics (ABS), including the legislative obligations of the ABS, financing and governance arrangements, and mechanisms for scrutiny of ABS operations, see the

ABS Institutional Environment.

Timeliness: The National Aboriginal and Torres Strait Islander Health Survey (NATSIHS) is

conducted every six years. The 2004-05 survey was conducted between August

2004 and July 2005. First results were released in April 2006.

The National Health Survey (NHS) is conducted every three years and enumerated over a 12-month period to account for seasonal variability in its measures. The 2004-05 survey was conducted concurrently with the 2004-05 NATSIHS. First

results were released in February 2006.

Accessibility: See also National Aboriginal and Torres Strait Islander Health Survey, 2004-05

(4715.0) and *National Health Survey: Summary of Results, 2004-05* (4364.0). Specialised data tables and Confidentialised Unit Record Files (CURFs) are also

available on request.

Interpretability: Data for this indicator have been age-standardised to account for differences

between the age structures of the Indigenous and non-Indigenous populations. Direct age-standardisation to the 2001 total Australian population was used. Age-standardised results provide a measure of relative difference only between

populations.

Information on how to interpret and use the data appropriately is available from the National Health Survey and National Aboriginal and Torres Strait Islander Health Survey 2004-05: Data Reference Package (4363.0.55.002), 2004-05 NATSIHS User's Guide (4715.0.55.004) and 2004-05 NHS User's Guide (4363.0.55.001).

Relevance: While Indigenous status is collected in the NHS, the survey sample and

methodology are not designed to provide output that separately identifies

Indigenous people. The NHS can, however, be used for providing non-Indigenous

comparisons.

The NATSIHS and the NHS are national surveys that collect information on health

status, risk factors and health-related actions.

Body Mass Index (BMI) is calculated in these surveys from height and weight information 'as reported' by respondents and hence may differ from BMI based on

measured height and weight.

Accuracy:

The NATSIHS is conducted in all states and territories and includes remoteand non-remote areas. The 2004-05 sample was 10,000 persons/5200 households, with a response rate of 81 per cent of households. The NHS is conducted in all states and territories excluding very remote areas. This exclusion has a small impact only on national and state estimates, except for the Northern Territory, where such persons comprise over 20 per cent of the population. As a consequence of this exclusion, comparisons between Indigenous and non-Indigenous people in remote areas are not available. The 2004-05 NHS response rate was 91 per cent of households.

Both surveys are weighted to account for non-response. Since it is derived from sample surveys, this indicator is subject to sampling error which occurs because only a small proportion of the population is used to produce estimates that represent the whole population. Sampling error can be reliably estimated and is based on the statistical methods used to design surveys.

Overall, this indicator has a relative standard error (RSE) of less than 25 per cent for all states and territories, except for non-Indigenous estimates for the Northern Territory (29 per cent) which should be used with caution. Finer levels of disaggregation (e.g. by the inclusion of other cross classifying variables) may result in higher levels of sampling error.

Coherence:

The 2004-05 NATSIHS and 2004-05 NHS had similar data content, shared common elements in the questionnaire, and were processed side by side. Data for this indicator are consistent with information presented in the *Aboriginal and Torres Strait Islander Health Performance Framework* (indicator 2.26).

Relational attributes

Related metadata references:

Has been superseded by National Indigenous Reform Agreement: PI 05-

Prevalence of overweight and obesity, 2013 QS Indigenous, Superseded 25/06/2014

Indicators linked to this Data Quality statement:

National Indigenous Reform Agreement: P06-Levels of obesity - Body Mass Index,

<u>2010</u>

Community Services (retired), Superseded 04/04/2011