

Type of dementia code NN

Identifying and definitional attributes

Metadata item type:	Value Domain
METEOR identifier:	423207
Registration status:	<ul style="list-style-type: none">• Health, Recorded 15/11/2021
Definition:	A code set representing the aetiology of dementia .

Representational attributes

Representation class:	Code																																						
Data type:	String																																						
Format:	NN																																						
Maximum character length:	2																																						
Permissible values:	<table><thead><tr><th>Value</th><th>Meaning</th></tr></thead><tbody><tr><td>01</td><td>Alzheimer's disease</td></tr><tr><td>02</td><td>Vascular dementia</td></tr><tr><td>03</td><td>Dementia with Lewy bodies</td></tr><tr><td>04</td><td>Mixed dementia</td></tr><tr><td>05</td><td>Fronto-temporal dementia - Behavioural variant</td></tr><tr><td>06</td><td>Fronto-temporal dementia - Primary progressive aphasia</td></tr><tr><td>07</td><td>Fronto-temporal dementia - Other/unspecified</td></tr><tr><td>08</td><td>Dementia in Creutzfeldt-Jakob disease</td></tr><tr><td>09</td><td>Dementia in Huntington's disease</td></tr><tr><td>10</td><td>Dementia in Parkinson's disease</td></tr><tr><td>11</td><td>Dementia in Human Immunodeficiency Virus</td></tr><tr><td>12</td><td>Delirium superimposed on dementia</td></tr><tr><td>13</td><td>Dementia due to use of alcohol</td></tr><tr><td>14</td><td>Dementia in other substance abuse</td></tr><tr><td>15</td><td>Dementia in other conditions</td></tr><tr><td>16</td><td>Unspecified dementia</td></tr><tr><td>17</td><td>Suspected dementia/diagnosis not confirmed</td></tr><tr><td>18</td><td>Not diagnosed with dementia after clinical assessment</td></tr></tbody></table>	Value	Meaning	01	Alzheimer's disease	02	Vascular dementia	03	Dementia with Lewy bodies	04	Mixed dementia	05	Fronto-temporal dementia - Behavioural variant	06	Fronto-temporal dementia - Primary progressive aphasia	07	Fronto-temporal dementia - Other/unspecified	08	Dementia in Creutzfeldt-Jakob disease	09	Dementia in Huntington's disease	10	Dementia in Parkinson's disease	11	Dementia in Human Immunodeficiency Virus	12	Delirium superimposed on dementia	13	Dementia due to use of alcohol	14	Dementia in other substance abuse	15	Dementia in other conditions	16	Unspecified dementia	17	Suspected dementia/diagnosis not confirmed	18	Not diagnosed with dementia after clinical assessment
Value	Meaning																																						
01	Alzheimer's disease																																						
02	Vascular dementia																																						
03	Dementia with Lewy bodies																																						
04	Mixed dementia																																						
05	Fronto-temporal dementia - Behavioural variant																																						
06	Fronto-temporal dementia - Primary progressive aphasia																																						
07	Fronto-temporal dementia - Other/unspecified																																						
08	Dementia in Creutzfeldt-Jakob disease																																						
09	Dementia in Huntington's disease																																						
10	Dementia in Parkinson's disease																																						
11	Dementia in Human Immunodeficiency Virus																																						
12	Delirium superimposed on dementia																																						
13	Dementia due to use of alcohol																																						
14	Dementia in other substance abuse																																						
15	Dementia in other conditions																																						
16	Unspecified dementia																																						
17	Suspected dementia/diagnosis not confirmed																																						
18	Not diagnosed with dementia after clinical assessment																																						
Supplementary values:	<table><tbody><tr><td>98</td><td>Unknown</td></tr><tr><td>99</td><td>Not stated/inadequately described</td></tr></tbody></table>	98	Unknown	99	Not stated/inadequately described																																		
98	Unknown																																						
99	Not stated/inadequately described																																						

Collection and usage attributes

Guide for use:	There are many forms of dementia as defined by their putative pathologies. The cognitive symptoms of different causes of dementia often overlap, and multiple
-----------------------	---

different causes of dementia can be present in a single person, especially after the age of 80. The codes 01-16 may only be assigned after a diagnosis has been recorded by a clinician.

A diagnosis of dementia should be made only after a clinical assessment, which should include:

- history taking from the person
- history taking from a person who knows the person well, if possible
- cognitive and mental state examination with a validated instrument
- physical examination
- a review of medication in order to identify and minimise use of medications, including over-the-counter products, that may adversely affect cognitive functioning and to simplify medication dosing
- consideration of other causes (including delirium or depression).

A basic dementia screen should be performed at the time of presentation, usually within primary care. It should include the following blood tests:

- routine haematology
- biochemistry tests (including electrolytes, calcium, glucose, and renal and liver function)
- thyroid function tests
- serum vitamin B12.

Structural imaging (with computed tomography (CT) or magnetic resonance imaging (MRI)) should usually be used in the assessment of people with suspected dementia to exclude other cerebral pathologies and to help establish the subtype diagnosis, unless clinical judgement indicates this inappropriate. Structural imaging may rarely be needed in those presenting with moderate-to-severe dementia, if the diagnosis is already clear.

CODE 01 Alzheimer's disease

Alzheimer's disease is a primary degenerative cerebral disease of unknown aetiology with characteristic neuropathological and neuro-chemical features. The disorder is usually insidious in onset and develops slowly but steadily over a period of several years.

This code corresponds to ICD-10-AM code combinations assigning F00.00-F00.01, F00.10-F00.11, F00.20-F00.21, F00.90-F00.91, G30.0, G30.1, G30.8, or G30.9.

Two codes must be recorded i.e. F00.00-F00.01 with G30.0, F00.10-F00.11 with G30.1, F00.20-F00.21 with G30.8, or F00.90-F00.91 with G30.9.

CODE 02 Vascular dementia

Vascular dementia is the result of infarction of the brain due to vascular disease, including hypertensive cerebrovascular disease. The infarcts are usually small but cumulative in their effect. Onset is usually in later life.

This code corresponds to ICD-10-AM codes F01.00-F01.01, F01.10-F01.11, F01.20-F01.21, F01.30-F01.31, F01.80-F01.81, F01.90-F01.91.

CODE 03 Dementia with Lewy bodies

Dementia caused by the degeneration and death of nerve cells in the brain, and characterised by Lewy bodies. Also known as Lewy body dementia.

This code corresponds to ICD-10-AM code combinations assigning F02.80-F02.81 and G31.3. Two codes must be recorded.

CODE 04 Mixed dementia

The person has dementia with atypical or mixed presentation, not distinct to a single dementia type.

This code corresponds to ICD-10-AM code combinations assigning F00.20-F00.21 and G30.8. Two codes must be recorded.

CODE 05 Fronto-temporal dementia - Behavioural variant

A progressive dementia caused by nerve cell damage to the frontal lobes of the brain, often leading to changes in personality, emotion and behaviour, as well as reduced judgement, empathy and motivation. Also known as Pick's disease.

This code corresponds to ICD-10-AM code combinations assigning F02.00-F02.01, and G31.0. Two codes must be recorded.

CODE 06 Fronto-temporal dementia - Primary progressive aphasia

A progressive dementia caused by nerve cell damage to the temporal lobes of the brain, leading to reduced abilities in language, speaking, writing and comprehension. Includes semantic variant and progressive non-fluent aphasia. Also known as Pick's disease.

This code corresponds to ICD-10-AM code combinations assigning F02.00-F02.01, and G31.0. Two codes must be recorded.

CODE 07 Fronto-temporal dementia - Other/unspecified

The person has a confirmed diagnosis of fronto-temporal dementia where the type is not the behavioural variant (code 05) or primary progressive aphasia (code 06), or where the type is unknown or unspecified.

This code corresponds to ICD-10-AM code combinations assigning F02.00-F02.01, and G31.0. Two codes must be recorded.

CODE 08 Dementia in Creutzfeldt-Jakob disease

A progressive dementia with extensive neurological signs, due to specific neuropathological changes that are presumed to be caused by a transmissible agent. Onset is usually in middle or later life, but may be at any adult age. The course is subacute, leading to death within one to two years.

This code corresponds to ICD-10-AM code combinations assigning A81.0 and F02.10-F02.11. Two codes must be recorded.

CODE 09 Dementia in Huntington's disease

A dementia occurring as part of a widespread degeneration of the brain. The disorder is transmitted by a single autosomal dominant gene. Symptoms typically emerge in the third and fourth decade. Progression is slow, leading to death usually within 10 to 15 years.

This code corresponds to ICD-10-AM code combinations assigning F02.20-F02.21 and G10. Two codes must be recorded.

CODE 10 Dementia in Parkinson's disease

A dementia developing in the course of established Parkinson's disease. No particular distinguishing clinical features have yet been demonstrated.

This code corresponds to ICD-10-AM code combinations assigning F02.30-F02.31 and G20. Two codes must be recorded.

CODE 11 Dementia in human immunodeficiency virus disease

Dementia developing in the course of HIV disease, in the absence of a concurrent illness or condition other than HIV infection that could explain the clinical features.

This code corresponds to ICD-10-AM codes B22 and F02.40-F02.41. Two codes must be recorded.

CODE 12 Delirium superimposed on dementia

Delirium superimposed on dementia occurs when a delirium occurs concurrently with a pre-existing dementia.

This code corresponds to ICD-10-AM code F05.1.

CODE 13 Dementia due to use of alcohol

Dementia caused by a disorder in which alcohol-induced changes of cognition, affect, personality, or behaviour persist beyond the period during which alcohol might reasonably be assumed to be operating.

This code corresponds to ICD-10-AM code F10.7.

CODE 14 Dementia in other substance abuse

Dementia caused by a disorder in which substance-induced changes of cognition, affect, personality, or behaviour persist beyond the period during which a direct psychoactive substance-related effect might reasonably be assumed to be operating.

This code corresponds to ICD-10-AM codes F13.70-F13.79, F18.7.

CODE 15 Dementia in other conditions

The person has dementia due to specific disease. Where specific dementia types and another condition are recorded, the specific type assigned to CODE 01–CODE 14 should be used in preference to CODE 15.

This code is equivalent to ICD-10-AM codes F02.80-F02.81 used in conjunction with any of the following codes: E75.0-E75.6, G40.00-G40.91, E83.0, E83.5, E01.0-E01.8, E03.0-E03.9, T36-T65, G35, A52.1, E52, M30.0, M32.0-M32.9, B56.0-B56.9, B57.0-B57.5, N18.5, or E53.8.

Two codes should be recorded, except where G31.3 is also recorded.

CODE 16 Unspecified dementia

The person has a confirmed diagnosis of dementia, and the type is unknown or unspecified.

This code is equivalent to ICD-10-AM codes F03.00-F03.01.

CODE 17 Suspected dementia/diagnosis not confirmed.

The person has undergone or is in the process of assessment for dementia and a diagnosis has not yet been clinically confirmed.

CODE 18 Not diagnosed with dementia after clinical assessment

The person has not received a diagnosis of Alzheimer's disease, vascular dementia, mixed dementia, fronto-temporal dementia, dementia with Lewy bodies, dementia in other conditions or dementia of type unspecified, as recorded in CODE 01–CODE 16 after a clinical assessment.

CODE 98 Unknown

Record if person has not been assessed for dementia in a clinical setting.

CODE 99 Not stated/inadequately described

This code is for use where information is insufficient to assign CODE 01–CODE 18 or information is unclear or unavailable.

Collection Methods:

The diagnosis must be recorded in a clinical setting by a qualified clinician, i.e. Geriatrician, GP, Neurologist etc.

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare

Relational attributes

Data elements implementing this value domain:

[Person—clinical type of dementia, code NN](#)

- [Health](#), Recorded 15/11/2021