

# Indigenous primary health care: PI22b-Proportion of female regular clients who have had a cervical screening, 2013

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Indigenous primary health care: PI22b-Proportion of female regular clients who have had a cervical screening, 2013

## Identifying and definitional attributes

<b>Metadata item type:</b>	Indicator
<b>Indicator type:</b>	Indicator
<b>Short name:</b>	PI22b-Proportion of female regular clients who have had a cervical screening, 2013
<b>METEOR identifier:</b>	438215
<b>Registration status:</b>	<a href="#">Health</a> , Superseded 21/11/2013 <a href="#">Indigenous</a> , Superseded 21/11/2013
<b>Description:</b>	Proportion of female regular clients who are Indigenous, aged 20 to 69 years, who have not had a hysterectomy and who have had a cervical screening within the previous 2 years, 3 years and 5 years.
<b>Rationale:</b>	Cancer mortality rates are higher for Indigenous Australians than for other Australians. National organised programs for cervical screening in Australia have been followed by reductions in mortality rates for these cancers. A pap smear every two years can prevent the most common form of cervical cancer and is the best protection against cervical cancer.
<b>Indicator set:</b>	<a href="#">Indigenous primary health care key performance indicators (2013)</a> <a href="#">Health</a> , Superseded 21/11/2013 <a href="#">Indigenous</a> , Superseded 21/11/2013

## Collection and usage attributes

<b>Population group age from:</b>	20 years
<b>Population group age to:</b>	69 years
<b>Computation description:</b>	<p>Proportion of female regular clients who are Indigenous, aged 20 to 69 years, who have not had a hysterectomy and who have had a cervical screening within the previous 2 years, 3 years and 5 years.</p> <p>'Regular client' refers to a client of an OATSIH-funded primary health care service (that is required to report against the Indigenous primary health care key performance indicators) who has an active medical record; that is, a client who has attended the OATSIH-funded primary health care service at least 3 times in 2 years.</p> <p>Presented as a percentage.</p> <p>Calculated separately for women who have had a cervical screening within the last 2 years, the last 3 years and the last 5 years.</p>
<b>Computation:</b>	$(\text{Numerator} \div \text{Denominator}) \times 100$
<b>Numerator:</b>	<p>Calculation A: Number of female regular clients who are Indigenous, aged 20 to 69 years, who have not had a hysterectomy and who have had a cervical screening within the previous 2 years.</p> <p>Calculation B: Number of female regular clients who are Indigenous, aged 20 to 69 years, who have not had a hysterectomy and who have had a cervical screening within the previous 3 years.</p> <p>Calculation C: Number of female regular clients who are Indigenous, aged 20 to 69 years, who have not had a hysterectomy and who have had a cervical screening within the previous 5 years.</p>

**Numerator data elements:**

[Person—sex, code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Guide for use**

Female only.

**Data Element / Data Set**

[Person—Indigenous status, code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Data Element / Data Set**

[Person—age, total years N\[NN\]](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Data Element / Data Set**

[Female—cervical screening indicator, yes/no/not stated/inadequately described code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Data Element / Data Set**

[Person—regular client indicator, yes/no code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Data Element / Data Set**

[Female—hysterectomy indicator, yes/no code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**Denominator:**

Calculation A, B and C: Total number of female regular clients who are Indigenous, aged 20-69 years who have not had a hysterectomy.

**Denominator data elements:**

**Data Element / Data Set**

[Person—sex, code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Guide for use**

Only female.

**Data Element / Data Set**

[Person—Indigenous status, code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Data Element / Data Set**

[Person—age, total years N\[NN\]](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Data Element / Data Set**

[Person—regular client indicator, yes/no code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Data Element / Data Set**

[Female—hysterectomy indicator, yes/no code N](#)

**Data Source**

[Indigenous primary health care data collection](#)

**NMDS / DSS**

[Indigenous primary health care DSS 2012-14](#)

**Representational attributes**

**Representation class:** Percentage

**Data type:** Real

Unit of measure: Person

## Indicator conceptual framework

Framework and dimensions: [Effective/Appropriate/Efficient](#)

## Data source attributes

Data sources:

### Data Source

[Indigenous primary health care data collection](#)

### Frequency

6 monthly

### Data custodian

Australian Institute of Health and Welfare.

## Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare (AIHW)

Department of Health and Ageing (DoHA)

## Relational attributes

Related metadata references:

Has been superseded by [Indigenous primary health care: PI22b-Proportion of regular clients who have had a cervical screening, 2014](#)

[Health](#), Superseded 13/03/2015

[Indigenous](#), Superseded 13/03/2015

See also [Indigenous primary health care: PI22a-Number of female regular clients who have had a cervical screening, 2013](#)

[Health](#), Superseded 21/11/2013

[Indigenous](#), Superseded 21/11/2013