

Indigenous primary health care: PI06a-Number of regular clients with Type II diabetes whose HbA1c measurement result was within a specified level, 2014

Identifying and definitional attributes

| | |
|-----------------------------|---|
| Metadata item type: | Indicator |
| Indicator type: | Output measure |
| Short name: | PI06a-Number of regular clients with Type II diabetes whose HbA1c measurement result was within a specified level, 2014 |
| METEOR identifier: | 504704 |
| Registration status: | <ul style="list-style-type: none">• Health, Superseded 13/03/2015• Indigenous, Superseded 13/03/2015 |

| | |
|---------------------|--|
| Description: | <p>Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded within either the previous 6 months or 12 months, was categorised as one of the following:</p> <ul style="list-style-type: none">• less than or equal to 7% (less than or equal to 53 mmol/mol);• greater than 7% but less than or equal to 8% (greater than 53 mmol/mol but less than or equal to 64 mmol/mol);• greater than 8% but less than 10% (greater than 64 mmol/mol but less than 86 mmol/mol) or;• greater than or equal to 10% (greater than or equal to 86 mmol/mol) |
|---------------------|--|

| | |
|-------------------|--|
| Rationale: | Chronic diseases, such as diabetes, are a major cause of morbidity and mortality among Aboriginal and Torres Strait Islander people. Monitoring glycosylated haemoglobin (HbA1c) levels plays a key role in the management of the disease. |
|-------------------|--|

| | |
|-----------------------|--|
| Indicator set: | Indigenous primary health care key performance indicators (2014) Health , Superseded 13/03/2015 Indigenous , Superseded 13/03/2015 |
|-----------------------|--|

Collection and usage attributes

Computation description: Count of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded within either the previous 6 months or 12 months, was categorised as one of the following:

- less than or equal to 7% (less than or equal to 53 mmol/mol);
- greater than 7% but less than or equal to 8% (greater than 53 mmol/mol but less than or equal to 64 mmol/mol);
- greater than 8% but less than 10% (greater than 64 mmol/mol but less than 86 mmol/mol) or;
- greater than or equal to 10% (greater than or equal to 86 mmol/mol)

'Regular client' refers to a client of an OATSIH-funded primary health care service (that is required to report against the Indigenous primary health care key performance indicators) who has an active medical record; that is, a client who has attended the OATSIH-funded primary health care service at least 3 times in 2 years.

HbA1c: glycosylated haemoglobin.

Exclude Type I diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, impaired fasting glucose, impaired glucose tolerance.

Presented as a number.

Calculated separately for each HbA1c range, for each time period.

Include only the most recent HbA1c measurement result from each time period.

Computation:

Numerator only

Numerator:

Calculation A: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 6 months, was less than or equal to 7% (less than or equal to 53 mmol/mol).

Calculation B: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 6 months, was greater than 7% but less than or equal to 8% (greater than 53 mmol/mol but less than or equal to 64 mmol/mol).

Calculation C: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 6 months, was greater than 8% but less than 10% (greater than 64 mmol/mol but less than 86 mmol/mol).

Calculation D: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 6 months, was greater than or equal to 10% (greater than or equal to 86 mmol/mol).

Calculation E: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 12 months, was less than or equal to 7% (less than or equal to 53 mmol/mol).

Calculation F: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 12 months, was greater than 7% but less than or equal to 8% (greater than 53 mmol/mol but less than or equal to 64 mmol/mol).

Calculation G: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 12 months, was greater than 8% but less than 10% (greater than 64 mmol/mol but less than 86 mmol/mol).

Calculation H: Number of regular clients who are Indigenous, have Type II diabetes and whose HbA1c measurement result, recorded in the previous 12 months, was greater than or equal to 10% (greater than or equal to 86 mmol/mol).

Numerator data elements:

Data Element / Data Set

[Person—diabetes mellitus status, code NN](#)

Data Source

[Indigenous primary health care data collection](#)

NMDS / DSS

[Indigenous primary health care DSS 2014-15](#)

Guide for use

Type II diabetes only.

Data Element / Data Set

[Person—Indigenous status, code N](#)

Data Source

[Indigenous primary health care data collection](#)

NMDS / DSS

[Indigenous primary health care DSS 2014-15](#)

Data Element / Data Set

[Person—regular client indicator, yes/no code N](#)

Data Source

[Indigenous primary health care data collection](#)

NMDS / DSS

[Indigenous primary health care DSS 2014-15](#)

Data Element / Data Set

[Person—glycosylated haemoglobin level, code N](#)

Data Source

[Indigenous primary health care data collection](#)

NMDS / DSS

[Indigenous primary health care DSS 2014-15](#)

Guide for use

Response to this is conditional on responding 'yes' to having had an HbA1c measurement result recorded in either the previous 6 months or 12 months.

- Disaggregation:**
1. Sex:
 - a) Male
 - b) Female
 2. Age:
 - a) 0-4 years
 - b) 5-14 years
 - c) 15-24 years
 - d) 25-34 years
 - e) 35-44 years
 - f) 45-54 years
 - g) 55-64 years
 - h) 65 years and over

Disaggregation data elements:

| |
|---|
| <p>Data Element / Data Set</p> <p>Person—sex, code N</p> <p>Data Source</p> <p>Indigenous primary health care data collection</p> <p>NMDS / DSS</p> <p>Indigenous primary health care DSS 2014-15</p> |
| <p>Data Element / Data Set</p> <p>Person—age, total years N[NN]</p> <p>Data Source</p> <p>Indigenous primary health care data collection</p> <p>NMDS / DSS</p> <p>Indigenous primary health care DSS 2014-15</p> |

Representational attributes

- Representation class:** Count
- Data type:** Real
- Unit of measure:** Person

Indicator conceptual framework

- Framework and dimensions:** [Effective/Appropriate/Efficient](#)

Data source attributes

- Data sources:**
- Data Source**
- [Indigenous primary health care data collection](#)
- Frequency**
- 6 monthly
- Data custodian**
- Australian Institute of Health and Welfare.

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare
Department of Health

Relational attributes

Related metadata references:

Supersedes [Indigenous primary health care: PI06a-Number of regular clients with Type II diabetes whose HbA1c measurement result was within a specified level, 2013](#)

- [Health](#), Superseded 21/11/2013
- [Indigenous](#), Superseded 21/11/2013

Has been superseded by [Indigenous primary health care: PI06a-Number of regular clients with Type II diabetes whose HbA1c measurement result was within a specified level, 2015](#)

- [Health](#), Superseded 05/10/2016
- [Indigenous](#), Superseded 20/01/2017

See also [Indigenous primary health care: PI06b-Proportion of regular clients with Type II diabetes whose HbA1c measurement result was within a specified level, 2014](#)

- [Health](#), Superseded 13/03/2015
- [Indigenous](#), Superseded 13/03/2015