

Primary postpartum haemorrhage indicator

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Primary postpartum haemorrhage indicator

Identifying and definitional attributes

Metadata item type:	Property
METEOR identifier:	504953
Registration status:	Health , Standard 07/03/2014
Definition:	An indicator of whether a primary postpartum haemorrhage occurred. Primary postpartum haemorrhage is characterised by abnormal or excessive bleeding from the genital tract within the first 24 hours postpartum.
Property group:	Birth event

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Data Element Concepts implementing this property: [Female—primary postpartum haemorrhage indicator](#)
[Health](#), Standard 07/03/2014