

National Healthcare Agreement: PI 16- Potentially avoidable deaths, 2013 QS

Identifying and definitional attributes

Metadata item type:	Quality Statement
METEOR identifier:	511943
Registration status:	<ul style="list-style-type: none">• Health, Superseded 14/01/2015

Relational attributes

Indicators linked to this Quality statement:	National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2013 Health , Superseded 30/04/2014
--	--

Data quality

Institutional environment:	These collections are conducted under the <i>Census and Statistics Act 1905</i> . For information on the institutional environment of the Australian Bureau of Statistics (ABS), including the legislative obligations of the ABS, financing and governance arrangements, and mechanisms for scrutiny of ABS operations, see ABS Institutional Environment.
----------------------------	---

Timeliness: Causes of death data is published on an annual basis. Death records are provided electronically to the ABS by individual Registrars on a monthly basis for compilation into aggregate statistics on a quarterly and annual basis. One dimension of timeliness in death registrations data is the interval between the occurrence and registration of a death. As a result, a small number of deaths occurring in one year are not registered until the following year or later.

Preliminary estimated resident population (ERP) data is compiled and published quarterly and is generally made available five to six months after the end of each reference quarter.

Every year, the 30 June ERP is further disaggregated by sex and single year of age, and is made available five to six months after the end of the reference quarter. Commencing with data for September quarter 2006, revised estimates are released annually and made available 21 months after the end of the reference period for the previous financial year, once more accurate births, deaths and net overseas migration data becomes available. In the case of births and deaths, the revised data is compiled on a date of occurrence basis. In the case of net overseas migration, final data is based on actual traveller behaviour. Final estimates are made available every 5 years after a census and revisions are made to the previous intercensal period. ERP data is not changed once it has been finalised. Releasing preliminary, revised and final ERP involves a balance between timeliness and accuracy.

For further information on ABS Estimated Resident Population, see the relevant Data Quality Statement.

Accessibility: Causes of death data are available in a variety of formats on the ABS website under the 3303.0 product family. ERP data is available in a variety of formats on the ABS website under the 3101.0 and 3201.0 product families. Further information on deaths and mortality may be available on request. The ABS observes strict confidentiality protocols as required by the *Census and Statistics Act 1905*. This may restrict access to data at a very detailed level.

Interpretability: Data for this indicator have been age-standardised, using the direct method, to 'under 75 years' of age. Direct age-standardisation to the 2001 total Australian population was used. Age-standardised results provide a measure of relative difference only between populations.

Relevance: The ABS Causes of Death collection includes all deaths that occurred and were registered in Australia, including deaths of persons whose usual residence is overseas. Deaths of Australian residents that occurred outside Australia may be registered by individual Registrars, but are not included in ABS deaths or causes of death statistics.

Data in the Causes of Death collection include demographic items, as well as causes of death information, which is coded according to the International Statistical Classification of Diseases and Related Health Problems (ICD). ICD is the international standard classification for epidemiological purposes and is designed to promote international comparability in the collection, processing, classification, and presentation of causes of death statistics. The classification is used to classify diseases and causes of disease or injury as recorded on many types of medical records as well as death records. The ICD has been revised periodically to incorporate changes in the medical field. The 10th revision of ICD (ICD-10) has been used by the ABS to code cause of death since 1997.

For further information on the ABS Causes of Death collection, see the relevant Data Quality Statement.

Accuracy: Information on causes of death is obtained from a complete enumeration of deaths registered during a specified period and is not subject to sampling error. However, deaths data sources are subject to nonsampling error which can arise from inaccuracies in collecting, recording and processing the data.

Although it is considered likely that most deaths of Aboriginal and Torres Strait Islander (Indigenous) Australians are registered, a proportion of these deaths are not registered as Indigenous. Information about the deceased is supplied by a relative or other person acquainted with the deceased, or by an official of the institution where the death occurred and may differ from the self-identified Indigenous origin of the deceased.

Forms are often not subject to the same best practice design principles as statistical questionnaires, and respondent and/or interviewer understanding is rarely tested. Over-precise analysis of Indigenous deaths and mortality should be avoided.

Previous COAG reporting and *Causes of Death, Australia* (cat. no. 3303.0) publications prior to the 2010 edition indicated that all coroner certified deaths registered after 1 January 2007 are now subject to a revisions process. In order to improve the quality of historical data, the 2006 reference year data has also been revised. Therefore, in this round of COAG reporting, 2006, 2007 and 2008 data is final, 2009 data is revised and 2010 data is preliminary. Data for 2009 and 2010 is subject to further revisions. This is a change from previous years (up to the 2005 reference year) where all ABS processing of causes of death data for a particular reference period was finalised approximately 13 months after the end of the reference period. Where insufficient information was available to code a cause of death (e.g. a coroner certified death was yet to be finalised by the Coroner), less specific ICD codes were assigned as required by the ICD coding rules. The revision process enables the use of additional information relating to coroner certified deaths, as it becomes available over time. This results in increased specificity of the assigned ICD-10 codes.

Revisions will only impact on coroner certified deaths, as further information becomes available to the ABS about the causes of these deaths. See *Technical Note: Causes of Death Revisions 2006 and Causes of Death Revisions 2008 and 2009* and in *Causes of Death, Australia, 2010* (cat. no. 3303.0).

In November 2010, the Queensland Registrar of Births, Deaths and Marriages advised the ABS of an outstanding deaths registration initiative undertaken by the Registry. This initiative resulted in the November 2010 registration of 374 previously unregistered deaths which occurred between 1992 and 2006 (including a few for which a date of death was unknown).

Of these, around three-quarters (284) were deaths of Aboriginal and Torres Strait Islander Australians.

The ABS discussed different methods of adjustment of Queensland death registrations data for 2010 with key stakeholders. Following the discussion, a decision was made by the ABS and key stakeholders to use an adjustment method that added together deaths registered in 2010 for usual residents of Queensland which occurred in 2007, 2008, 2009 and 2010. This method minimises the impact on mortality indicators used in various government reports. However, care should still be taken when interpreting Aboriginal and Torres Strait Islander death data for Queensland for 2010. Please note that there are differences between data output in the *Causes of Death, Australia, 2010* publication (cat. No. 3303.0) and 2010 data reported for COAG, as this adjustment was not applied in the publication. For further details see *Technical Note: Registration of outstanding deaths, Queensland 2010*, from the *Deaths, Australia, 2010* publication (cat. no. 3302.0) and *Explanatory Note 103* in the *Causes of Death, Australia, 2010* publication (cat. no. 3303.0).

Investigation conducted by the WA Registrar of Births, Deaths and Marriages indicated that some deaths of non-Indigenous people were wrongly recorded as deaths of Indigenous people in WA for 2007, 2008 and 2009. The ABS discussed this issue with a range of key stakeholders and users of Aboriginal and Torres Strait Islander deaths statistics.

Following this discussion, the ABS did not release WA Aboriginal and Torres Strait Islander deaths data for the years 2007, 2008 and 2009 in the 2010 issue of *Deaths, Australia* publication, or in the 2011 COAG data supply. The WA Registry corrected the data and resupplied the corrected data to the ABS. These corrected data were then released by the ABS in spreadsheets attached to *Deaths, Australia, 2010* (ABS, 2011) publication on 24 May 2012, and are now included in this round of COAG reporting.

All ERP data sources are subject to non-sampling error. Non-sampling error can arise from inaccuracies in collecting, recording and processing the data. In the case of Census and Post Enumeration Survey (PES) data, every effort is made to minimise reporting error by the careful design of questionnaires, intensive training and supervision of interviewers, and efficient data processing procedures. The

ABS does not have control over any non-sampling error associated with births, deaths and migration data.

For more information see the *Demography Working Paper 1998/2 - Quarterly birth and death estimates, 1998* (cat. no. 3114.0) and *Australian Demographic Statistics* (cat. no. 3101.0).

Non-Indigenous estimates are available for census years only. In the intervening years, Indigenous population projections are based on assumptions about past and future levels of fertility, mortality and migration. In the absence of non-Indigenous population figures for these years, it is possible to derive denominators for calculating non-Indigenous rates by subtracting the projected Indigenous population from the total population. Such figures have a degree of uncertainty and should be used with caution, particularly as the time from the base year of the projection series increases.

Non-Indigenous data from the Causes of Death collection do not include death registrations with a 'not stated' Indigenous status.

Some rates are unreliable due to small numbers of deaths over the reference period. Resultant rates could be misleading, for example, where the non-Indigenous mortality rate is higher than the indigenous mortality rate. As such, age-standardised death rates based on a very low death count have been deemed unpublishable. Some cells have also not been published to prevent back-calculation of these suppressed cells.

Caution should be used when interpreting rates for this indicator.

Data for 2007 was final in both the 2011 and 2012 supply of COAG data.

Despite this, there are very slight differences between the 2007 data reported in Table 20.4, compared with the same raw figures that were supplied in 2011 (Table 20.6). This is due to age at death 'not stated' being included in the raw figures for preventable and treatable causes of death in the 2011 reporting, but not in the current data supply. It was also identified that ICD-10 code J02.0 is included in the definition of two treatable variables: Selected invasive bacterial and protozoal infections and Upper respiratory tract infection. The Productivity Commission identified their preference to have J02.0 included only in data for Selected invasive bacterial and protozoal infections (thus avoiding deaths for this code being double-counted in the total figures). This, and the inclusion of age 'not stated' last year, account for all differences between this year and last year's 2007 data for Indicator 20.

Coherence:

The methods used to construct the indicator are consistent and comparable with other collections and with international practice.

Relational attributes

Related metadata references:

Supersedes [National Healthcare Agreement: PI 20-Potentially avoidable deaths, 2012 QS](#)

- [Health](#), Superseded 14/01/2015

Has been superseded by [National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2014 QS](#)

- [Health](#), Superseded 14/01/2015

© Australian Institute of Health and Welfare 2015–2022

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 3.0 (CC BY 3.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build upon this work. However, you must attribute the AIHW as the copyright holder of the work in compliance with our attribution policy available at www.aihw.gov.au/copyright. The full terms and conditions of this licence are available at <http://creativecommons.org/licenses/by3.0/au/>.

Enquiries relating to copyright should be addressed to the Head of the Communications, Media and Marketing Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.