

Female—type of diabetes mellitus therapy during pregnancy, code N

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Female—type of diabetes mellitus therapy during pregnancy, code N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Diabetes therapy type during pregnancy
METEOR identifier:	516185
Registration status:	Health , Superseded 02/08/2017
Definition:	The type of diabetes mellitus treatment which a female is prescribed during pregnancy, as represented by a code.
Data Element Concept:	Female—type of diabetes mellitus therapy during pregnancy
Value Domain:	Type of diabetes mellitus therapy during pregnancy code N

Value domain attributes

Representational attributes

Representation class:	Code
Data type:	Number
Format:	N
Maximum character length:	1

	Value	Meaning
Permissible values:	1	Insulin
	2	Oral hypoglycaemic
	3	Diet and exercise
Supplementary values:	9	Not stated/inadequately described

Collection and usage attributes

Guide for use:	<p>All therapies prescribed during pregnancy should be recorded. Therefore more than one code can be selected when reporting this item.</p> <p>CODE 2 Oral hypoglycaemic</p> <p>This code includes the options of sulphonylurea, biguanide (e.g. metformin), alpha-glucosidase inhibitor, thiazolidinedione, meglitinide, combination (e.g. biguanide and sulphonylurea), or other.</p> <p>CODE 3 Diet and exercise</p> <p>This code includes the options of generalised prescribed diet; avoid added sugar/simple carbohydrates (CHOs); low joule diet; portion exchange diet and uses glycaemic index and a recommendation for increased exercise.</p> <p>CODE 9 Not stated/inadequately described</p> <p>To be recorded by data entry personnel (state/territory health authority) if the data field is left blank or is inadequately completed in the perinatal data collection form or extract. Clinicians should not record Code 9.</p>
-----------------------	---

Data element attributes

Collection and usage attributes

Collection methods: Jurisdictions that record perinatal data using the ICD-10-AM should apply the following codes:

'Code 1 Insulin' is equivalent in the ICD-10-AM to a code in the range O24.1–24.9 used in conjunction with the fifth character '2' (insulin treated).

'Code 2 Oral hypoglycaemic' is equivalent in the ICD-10-AM to a code in the range O24.1–24.9 used in conjunction with the fifth character '3' (oral hypoglycaemic therapy).

'Code 3 Diet and exercise' is equivalent in the ICD-10-AM to a code in the range O24.1–24.9 used in conjunction with the fifth character '4' (other; diet; exercise; lifestyle management).

For example, for a mother who has pre-existing Type 2 diabetes mellitus and uses oral hypoglycaemic therapy and insulin, this would be coded in the ICD-10-AM as O2412 and O2413 and would be reported against this data item using Codes 1 and 2.

Source and reference attributes

Reference documents: National Casemix and Classification Centre 2013. The International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Australian Modification (ICD-10-AM), Australian Classification of Health Interventions (ACHI) and Australian Coding Standards (ACS), Eighth edition. National Casemix and Classification Centre, Australian Health Services Research Institute: University of Wollongong

Relational attributes

Related metadata references: Has been superseded by [Female—type of diabetes mellitus therapy during pregnancy, code N](#)
[Health](#), Superseded 12/12/2018

See also [Female—diabetes mellitus during pregnancy indicator, yes/no/not stated/inadequately described code N](#)
[Health](#), Superseded 02/08/2017

See also [Female—type of diabetes mellitus during pregnancy, code N](#)
[Health](#), Superseded 02/08/2017

See also [Person—diabetes therapy type, code NN](#)
[Health](#), Standard 01/03/2005

Implementation in Data Set Specifications:

[Perinatal DSS 2014-15](#)

[Health](#), Superseded 13/11/2014

Implementation start date: 01/07/2014

Implementation end date: 30/06/2015

Conditional obligation:

Conditional on diabetes mellitus during pregnancy indicator being coded as yes.

[Perinatal DSS 2015-16](#)

[Health](#), Superseded 04/09/2015

Implementation start date: 01/07/2015

Implementation end date: 30/06/2016

Conditional obligation:

This data element is conditional on [Female—diabetes mellitus during pregnancy indicator, yes/no/not stated/inadequately described code N](#) being coded to Yes.

[Perinatal NBEDS 2016-17](#)

[Health](#), Superseded 05/10/2016

Implementation start date: 01/07/2016

Implementation end date: 30/06/2017

Conditional obligation:

This data element is conditional on [Female—diabetes mellitus during pregnancy indicator, yes/no/not stated/inadequately described code N](#) being coded to Yes.

[Perinatal NBEDS 2017-18](#)

[Health](#), Superseded 02/08/2017

Implementation start date: 01/07/2017

Implementation end date: 30/06/2018

Conditional obligation:

This data element is conditional on [Female—diabetes mellitus during pregnancy indicator, yes/no/not stated/inadequately described code N](#) being coded to Yes.