

Indications for caesarean section code NN

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Indications for caesarean section code NN

Identifying and definitional attributes

Metadata item type:	Value Domain
METEOR identifier:	516638
Registration status:	Health , Superseded 13/11/2014
Definition:	A code set representing indications for a caesarean section.

Representational attributes

Representation class:	Code
Data type:	String
Format:	NN
Maximum character length:	2

	Value	Meaning
Permissible values:	01	Fetal compromise
	02	Suspected fetal macrosomia
	03	Malpresentation
	04	Lack of progress; less than or equal to 3 cm cervical dilatation
	05	Lack of progress in the first stage; 4 cm to less than 10 cm cervical dilatation
	06	Lack of progress in the second stage
	07	Placenta praevia
	08	Placental abruption
	09	Vasa praevia
	10	Antepartum/intrapartum haemorrhage
	11	Multiple pregnancy
	12	Unsuccessful attempt at assisted delivery
	13	Unsuccessful induction
	14	Cord prolapse
	15	Previous caesarean section
	16	Previous shoulder dystocia
	17	Previous perineal trauma/4th degree tear
	18	Previous adverse fetal/neonatal outcome
	19	Other obstetric, medical, surgical, psychological indications
	20	Maternal choice in the absence of any obstetric, medical, surgical, psychological indications
Supplementary values:	99	Not stated/inadequately described

Collection and usage attributes

Guide for use:**CODE 01 Fetal compromise**

This includes suspected or actual fetal compromise and intra uterine growth restriction (IUGR).

CODE 04 Lack of progress; less than or equal to 3 cm cervical dilatation

Lack of progress includes slow or no progress.

If there has been an attempted induction of labour and then a lack of progress leading to a caesarean section use Code 13 as the main indication and Code 04 as an additional indication.

CODE 05 Lack of progress in the first stage; 4 cm to less than 10 cm cervical dilatation

Lack of progress includes slow or no progress.

If there has been an attempted induction of labour and then a lack of progress leading to a caesarean section use Code 13 as the main indication and Code 05 as an additional indication.

CODE 06 Lack of progress in the second stage

Lack of progress includes slow or no progress.

CODE 07 Placenta praevia

Record placenta praevia as the indication for caesarean section if there is ultrasound or clinical evidence that the edge of the placenta covers the internal cervical os, or encroaches into the lower segment less than 2 cm away from the internal cervical os.

CODE 08 Placental abruption

Record placental abruption as the indication for caesarean section if there is ultrasound or clinical evidence antenatally of abruption of the placenta prior to onset or during labour.

CODE 09 Vasa praevia

Record vasa praevia as the indication for caesarean section if there is ultrasound or visual evidence of exposed fetal blood vessels running across the fetal membrane below or at the level of the fetal presenting part in the lower segment of the uterus. This code is to be used when the caesarean section is planned or in the case of an emergency when the vessels may have ruptured.

CODE 10 Antepartum/intrapartum haemorrhage

Record antepartum/intrapartum haemorrhage as the indication for caesarean section if there has been any antenatal or intrapartum vaginal bleeding that leads to the immediate delivery of the baby by caesarean section. This code should only be used as a main indication if a more specific cause of the antepartum/intrapartum haemorrhage is not known.

Where there is a vasa praevia and an antepartum/intrapartum haemorrhage, Code 09 is to be recorded as the main indication and Code 10 as an additional indication.

CODE 19 Other obstetric, medical, surgical, psychological indications

Where a woman has a psychopathological indication for caesarean section, e.g. extreme fear of natural childbirth, this code should be used. It is not to be used for psychosocial indications which should be coded under Code 20.

CODE 20 Maternal choice in the absence of any obstetric, medical, surgical, psychological indications

This includes psychosocial indications.

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Related metadata references: Has been superseded by [Indication for caesarean section code N\[N\] Health](#), Superseded 02/08/2017

Data elements implementing this value domain: [Birth event—additional indications for caesarean section, code NN Health](#), Superseded 13/11/2014
[Birth event—main indication for caesarean section, code NN Health](#), Superseded 13/11/2014