## Hypertensive disorder during pregnancy indicator



parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

# Hypertensive disorder during pregnancy indicator

#### Identifying and definitional attributes

Metadata item type: Property
METEOR identifier: 516795

**Registration status:** <u>Health</u>, Superseded 05/10/2016

**Definition:** An indicator of whether a <u>hypertensive disorder during pregnancy</u> is present.

Hypertensive disorders in pregnancy include eclampsia, preeclampsia, gestational

hypertension, chronic hypertension (essential and secondary) and preeclampsia/eclampsia superimposed on a hypertensive disorder.

Property group: <u>Illness and/or Injury event</u>

### Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Origin: National Perinatal Data Development Committee

Australian Institute of Health and Welfare

**Reference documents:** SOMANZ Guidelines for the Management of Hypertensive Disorders of Pregnancy

2008

#### Relational attributes

Related metadata

references:

Has been superseded by Hypertensive disorder during pregnancy indicator

Health, Superseded 03/12/2020

Data Element Concepts

implementing this

property:

Female—hypertensive disorder during pregnancy indicator

Health, Superseded 05/10/2016