Blood transfusion due to primary postpartum haemorrhage indicator

Exported from METEOR (AIHW's Metadata Online Registry)
© Australian Institute of Health and Welfare 2024
This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY4.0 (CC BY4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Blood transfusion due to primary postpartum haemorrhage indicator

Identifying and definitional attributes

Metadata item type: Property
METEOR identifier: 522206

Registration status: Health, Superseded 17/12/2021

Definition: An indicator of whether blood or blood products or substitutes were received

intravenously due to abnormal or excessive bleeding from the genital tract within the

first 24 hours postpartum.

Property group: Birth event

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Related metadata Has been superseded by <u>Blood transfusion indicator</u>

references: Health, Standard 17/12/2021

Data Element Concepts Female—blood transfusion due to primary postpartum haemorrhage indicator

implementing this Health, Superseded 17/12/2021

property: