

Hypertensive disorder during pregnancy

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Hypertensive disorder during pregnancy

Identifying and definitional attributes

Metadata item type: Glossary Item

METEOR identifier: 523104

Registration status: [Health](#), Superseded 05/10/2016
[Tasmanian Health](#), Standard 20/12/2016

Definition: Hypertensive disorder during pregnancy includes pre-existing hypertensive disorders, hypertension arising in pregnancy and associated disorders such as eclampsia and preeclampsia.

Hypertension in pregnancy is defined as:

1. Systolic blood pressure greater than or equal to 140 mmHg and/or
2. Diastolic blood pressure greater than or equal to 90 mmHg.

Measurements should be confirmed by repeated readings over several hours. Elevations of both systolic and diastolic blood pressures have been associated with adverse fetal outcome and therefore both are important.

Disorders associated with hypertension such as eclampsia and preeclampsia are further characterised by symptoms such as proteinuria, oedema or high body temperature.

There are several reasons to support the blood pressure readings defined above as diagnostic of hypertension in pregnancy:

- Perinatal mortality rises with diastolic blood pressures above 90 mmHg
- Readings above this level were beyond two standard deviations of mean blood pressure in a New Zealand cohort of normal pregnant women
- The chosen levels are consistent with international guidelines and correspond with the current diagnosis of hypertension outside of pregnancy.

This definition of hypertensive disorder in pregnancy from the Society of Obstetric Medicine in Australia and New Zealand (SOMANZ) aligns with the definition of the International Society for the Study of Hypertension in Pregnancy (ISSHP).

Source and reference attributes

Reference documents: Lowe S, Brown M, Dekker G, Gatt S, McLintock C, McMahon L et al 2008. Guidelines for the Management of Hypertensive Disorders of Pregnancy. Society of Obstetric Medicine of Australia and New Zealand (SOMANZ).

Brown M, Lindheimer M, Swiet M, Assche A and Moutquin J-M 2001. The classification and diagnosis of the hypertensive disorders of pregnancy: statement from the International Society for the Study of Hypertension in Pregnancy (ISSHP). Hypertension in pregnancy 20(1), ix-xiv.

Relational attributes

Related metadata references: Has been superseded by [Hypertensive disorder during pregnancy](#) [Health](#), Superseded 03/12/2020

**Metadata items which use
this glossary item:**

[Female—hypertensive disorder during pregnancy indicator](#)

[Health](#), Superseded 05/10/2016

[Female—hypertensive disorder during pregnancy indicator, yes/no/not
stated/inadequately described code N](#)

[Health](#), Superseded 05/10/2016

[Female—type of hypertensive disorder during pregnancy](#)

[Health](#), Superseded 05/10/2016

[Female—type of hypertensive disorder during pregnancy, code N](#)

[Health](#), Superseded 05/10/2016

[Hypertensive disorder during pregnancy indicator](#)

[Health](#), Superseded 05/10/2016

[Perinatal DSS 2014-15](#)

[Health](#), Superseded 13/11/2014

[Perinatal DSS 2015-16](#)

[Health](#), Superseded 04/09/2015

[Perinatal NBEDS 2016-17](#)

[Health](#), Superseded 05/10/2016

[Type of hypertensive disorder during pregnancy](#)

[Health](#), Superseded 05/10/2016

[Type of hypertensive disorder during pregnancy code N](#)

[Health](#), Superseded 05/10/2016