## Estimated blood loss indicating primary postpartum haemorrhage



made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

## Estimated blood loss indicating primary postpartum haemorrhage

## Identifying and definitional attributes

Metadata item type: Property
METEOR identifier: 524208

Registration status: Health, Standard 07/03/2014

**Definition:** Estimated volume of abnormal or excessive blood loss from the genital tract within

the first 24 hours postpartum.

A blood loss of 500 mls is the usual minimum amount for identification of postpartum haemorrhage however a woman's haemodynamic instability is also taken into account, meaning that a smaller blood loss may be significant in a severely compromised woman. A loss of 1,000 mls or more is considered major or

severe although definitions of severity vary.

Property group: Birth event

## Source and reference attributes

**Submitting organisation:** National Perinatal Data Development Committee

**Relational attributes** 

Data Element Concepts implementing this

property:

Female—estimated blood loss indicating primary postpartum haemorrhage

Health, Standard 07/03/2014