

Patient—colonoscope withdrawal time

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Patient—colonoscopy withdrawal time

Identifying and definitional attributes

Metadata item type:	Data Element Concept
METEOR identifier:	530140
Registration status:	Health , Standard 29/08/2014
Definition:	The time taken to withdraw the colonoscopy from the patient during a medical test or evaluation.

Object Class attributes

Identifying and definitional attributes

Object class:	Patient
Definition:	A person for whom a health service accepts responsibility for treatment and/or care.
Specialisation of:	Person/group of persons

Source and reference attributes

Submitting organisation:	Australian Institute of Health and Welfare
--------------------------	--

Property attributes

Identifying and definitional attributes

Property:	Colonoscopy withdrawal time
Definition:	<p>The time taken to withdraw the colonoscopy during a medical test or evaluation. This is the duration it takes to remove the colonoscopy from the caecum (considered to be the start of the colon) to the rectum.</p> <p>Longer colonoscopy withdrawal times have been demonstrated to improve polyp detection rates, and, conversely, rapid withdrawal of the colonoscopy may miss lesions and reduce the effectiveness of colon cancer prevention.</p>
Property group:	Service provision event

Source and reference attributes

Reference documents:	<p>ASGE & ACG (American Society for Gastrointestinal Endoscopy & American College of Gastroenterology) 2006. Taskforce on Quality in Endoscopy quality indicators for gastrointestinal endoscopic procedures: an introduction, ASGE/ACG Taskforce on Quality in Endoscopy. Gastrointestinal Endoscopy 63(4).</p> <p>NBCSP-QWG (National Bowel Cancer Screening Program—Quality Working Group) 2009. Improving Colonoscopy Services in Australia. Canberra: DoHA. Viewed 23 October 2013, www.cancerscreening.gov.au/internet/screening/publishing.nsf/Content/nbcs-imp-col-ser-0709-cnt</p>
----------------------	---

Data element concept attributes

Relational attributes

Data Elements implementing this Data Element Concept:	Patient—colonoscopy withdrawal time, total minutes NNN Health , Standard 29/08/2014
---	--

