

Indication for caesarean section code N[N]

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Indication for caesarean section code N[N]

Identifying and definitional attributes

Metadata item type:	Value Domain
METEOR identifier:	587034
Registration status:	Health , Superseded 02/08/2017
Definition:	A code set representing indications for a caesarean section.

Representational attributes

Representation class:	Code
Data type:	Number
Format:	N[N]
Maximum character length:	2

	Value	Meaning
Permissible values:	1	Fetal compromise
	2	Suspected fetal macrosomia
	3	Malpresentation
	10	Lack of progress; less than or equal to 3 cm cervical dilatation
	11	Lack of progress in the first stage; greater than 3 cm to less than 10 cm cervical dilatation
	12	Lack of progress in the second stage
	13	Placenta praevia
	14	Placental abruption
	15	Vasa praevia
	16	Antepartum/intrapartum haemorrhage
	17	Multiple pregnancy
	18	Unsuccessful attempt at assisted delivery
	19	Cord prolapse
	20	Previous adverse perinatal outcome
	21	Previous caesarean section
	22	Previous severe perineal trauma
	23	Previous shoulder dystocia
	29	Other obstetric, medical, surgical, psychological indications
	30	Maternal choice in the absence of any obstetric, medical, surgical, psychological indications
Supplementary values:	99	Not stated/inadequately described

Collection and usage attributes

Guide for use:**CODE 1 Fetal compromise**

This includes suspected or actual fetal compromise and intra uterine growth restriction (IUGR).

CODE 10 Lack of progress; less than or equal to 3 cm cervical dilatation

Lack of progress includes slow or no progress.

This code may be appropriate for cases of unsuccessful induction.

CODE 11 Lack of progress in the first stage; greater than 3 cm to less than 10 cm cervical dilatation

Lack of progress includes slow or no progress.

CODE 12 Lack of progress in the second stage

Lack of progress includes slow or no progress.

CODE 13 Placenta praevia

Record placenta praevia as the indication for caesarean section if there is ultrasound or clinical evidence that the edge of the placenta covers the internal cervical os, or encroaches into the lower segment less than 2 cm away from the internal cervical os.

CODE 14 Placental abruption

Record placental abruption as the indication for caesarean section if there is ultrasound or clinical evidence antenatally of abruption of the placenta prior to onset or during labour.

CODE 15 Vasa praevia

Record vasa praevia as the indication for caesarean section if there is ultrasound or visual evidence of exposed fetal blood vessels running across the fetal membrane below or at the level of the fetal presenting part in the lower segment of the uterus. This code is to be used when the caesarean section is planned or in the case of an emergency when the vessels may have ruptured.

CODE 16 Antepartum/intrapartum haemorrhage

Record antepartum/intrapartum haemorrhage as the indication for caesarean section if there has been any antenatal or intrapartum vaginal bleeding that leads to the immediate delivery of the baby by caesarean section. This code should only be used as a main indication if a more specific cause of the antepartum/intrapartum haemorrhage is not known.

Where there is a vasa praevia and an antepartum/intrapartum haemorrhage, Code 15 is to be recorded as the main indication and Code 16 as an additional indication.

CODE 22 Previous severe perineal trauma

Includes previous 4th degree tears and other perineal damage or injury where a woman is symptomatic and the clinician determines the caesarean section is justified.

CODE 29 Other obstetric, medical, surgical, psychological indications

Where a woman has a psychopathological indication for caesarean section, e.g. extreme fear of natural childbirth, this code should be used. It is not to be used for psychosocial indications which should be coded under Code 30.

CODE 30 Maternal choice in the absence of any obstetric, medical, surgical, psychological indications

This includes psychosocial indications.

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Related metadata references:	Supersedes Indications for caesarean section code NN Health , Superseded 13/11/2014
	Has been superseded by Indication for caesarean section code N[N] Health , Superseded 12/12/2018
Data elements implementing this value domain:	Birth event—additional indication for caesarean section, code N[N] Health , Superseded 02/08/2017
	Birth event—main indication for caesarean section, code N[N] Health , Superseded 02/08/2017