

Pelvic lymph node dissection indicator

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Pelvic lymph node dissection indicator

Identifying and definitional attributes

Metadata item type:	Property
METEOR identifier:	587594
Registration status:	Health , Standard 14/05/2015
Definition:	An indicator of the occurrence of a pelvic lymph node dissection. Pelvic lymph node dissection is completed for the purpose of determining lymph node metastasis in cancer. This procedure is most commonly used in prostate and bladder cancers.

Source and reference attributes

Submitting organisation:	Cancer Australia
---------------------------------	------------------

Relational attributes

Data Element Concepts implementing this property:	Person with cancer—pelvic lymph node dissection indicator Health , Standard 14/05/2015
--	---