Clinical care standard indicators: antimicrobial stewardship

Exported from METEOR (AIHW's Metadata Online Registry)
© Australian Institute of Health and Welfare 2024
This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.
You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our

Metadata 612216

meteor@aihw.gov.au.

attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Clinical care standard indicators: antimicrobial stewardship

Identifying and definitional attributes

Metadata item type: Indicator Set

Indicator set type: Other

METEOR identifier: 612216

Registration status: Australian Commission on Safety and Quality in Health Care, Retired 25/01/2022

Health, Standard 12/09/2016

Description: The Australian Commission on Safety and Quality in Health Care has produced the

Antimicrobial stewardship clinical care standard indicators to assist with local implementation of the Antimicrobial stewardship clinical care standard (ACSQHC 2014). The Antimicrobial stewardship clinical care standard aims to ensure that a patient with a bacterial infection receives optimal treatment with antibiotics. 'Optimal treatment' means treating patients with the right antibiotic to treat their condition, the right dose, by the right route, at the right time and for the right duration based on accurate assessment and timely review. Clinicians and health services can use the Antimicrobial stewardship clinical care standard and indicators to monitor the implementation of the quality statements, and support improvements as

needed.

Relational attributes

Related metadata references:

Has been superseded by <u>Clinical care standard indicators: antimicrobial</u> stewardship 2020

Australian Commission on Safety and Quality in Health Care, Standard 24/12/2020

See also Clinical care standard indicators: acute coronary syndromes

Health, Standard 12/09/2016

See also Clinical care standard indicators: delirium

Health, Standard 12/09/2016

See also Clinical care standard indicators: heavy menstrual bleeding

Health, Standard 17/10/2018

See also Clinical care standard indicators: hip fracture

Australian Commission on Safety and Quality in Health Care, Superseded

18/06/2018

Health, Standard 12/09/2016

See also Clinical care standard indicators: hip fracture 2018

Australian Commission on Safety and Quality in Health Care, Standard

15/05/2018

See also Clinical care standard indicators: osteoarthritis of the knee

Health, Standard 02/08/2017

Outcome areas linked to this Indicator set:

Documentation

Health, Standard 12/09/2016

Life-threatening conditions

Health, Standard 12/09/2016

Surgical prophylaxis

Health, Standard 12/09/2016

Use of broad-spectrum antibiotics

Health, Standard 12/09/2016

Use of guidelines and clinical condition

Health, Standard 12/09/2016

Indicators linked to this Indicator set:

Antimicrobial stewardship: 1a-Median time to first dose of antibiotics for patients with suspected bacterial meningitis, or for actual or suspected severe sepsis

<u>Australian Commission on Safety and Quality in Health Care</u>, Retired 25/01/2022

Health, Standard 12/09/2016

Antimicrobial stewardship: 4a-Antibiotic prescribing in accordance with current and peer reviewed clinical guidelines

Australian Commission on Safety and Quality in Health Care, Retired 25/01/2022

Health, Standard 12/09/2016

Antimicrobial stewardship: 4b-Antibiotic-allergy mismatch in prescribing
Australian Commission on Safety and Quality in Health Care, Retired
25/01/2022

Health, Standard 12/09/2016

Antimicrobial stewardship: 6a-Documentation of reason for prescribing antibiotics

Australian Commission on Safety and Quality in Health Care, Retired

25/01/2022

Health, Standard 12/09/2016

Antimicrobial stewardship: 7a-Review of patients prescribed broad-spectrum antibiotics

Australian Commission on Safety and Quality in Health Care, Retired 25/01/2022

Health, Standard 12/09/2016

Antimicrobial stewardship: 9a-Surgical antibiotic prophylaxis in accordance with guidelines

Australian Commission on Safety and Quality in Health Care, Retired 25/01/2022

Health, Standard 12/09/2016

Antimicrobial stewardship: 9b-Timely administration of prophylactic antibiotics prior to surgery

Australian Commission on Safety and Quality in Health Care, Retired 25/01/2022

Health, Standard 12/09/2016

Antimicrobial stewardship: 9c-Cessation of prophylactic antibiotics following surgery

Australian Commission on Safety and Quality in Health Care, Retired 25/01/2022

Health, Standard 12/09/2016

Collection and usage attributes

National reporting arrangement:

The Indicator specification: antimicrobial stewardship clinical care standard has been developed to assist with local implementation of the Antimicrobial stewardship clinical care standard (ACSQHC 2014). The indicators are intended for local use by hospitals and **Local Hospital Networks** (LHNs). There are no benchmarks set for any of the indicators in the Antimicrobial stewardship clinical care standard indicators set.

Hospitals using Antimicrobial stewardship clinical care standard indicators can compare their results against themselves during a previous period, amongst other hospitals in the LHN, or other external hospitals with whom they have made such arrangements. Hospitals may also look to the literature for the experiences of other hospitals.

Comments:

The Australian Commission on Safety and Quality in Health Care has produced the Antimicrobial stewardship clinical care standard (ACSQHC 2014) to support optimal treatment for people with bacterial infection requiring antibiotics and/or preventing bacterial infection, and is based on the best evidence available at the time of development. Health care professionals are advised to use clinical discretion and consideration of the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian when applying information contained within the clinical care standard. Consumers should use the information in the clinical care standard as a guide to inform discussions with their health care professional about the applicability of the standard to their individual condition.

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care