Antimicrobial stewardship: 9a-Surgical antibiotic prophylaxis in accordance with guidelines
Exported from METEOR (AIHW's Metadata Online Registry)

#### © Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

# Antimicrobial stewardship: 9a-Surgical antibiotic prophylaxis in accordance with guidelines

# Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

**Short name:** Indicator 9a-Surgical antibiotic prophylaxis in accordance with guidelines

**METEOR identifier:** 612228

Registration status: Australian Commission on Safety and Quality in Health Care, Retired 25/01/2022

Health, Standard 12/09/2016

**Description:** Proportion of patients for whom surgical prophylactic antibiotics were prescribed in

accordance with Therapeutic guidelines: antibiotic (Antibiotic Expert Group 2014)

or locally-endorsed antibiotic prescribing guidelines.

Rationale: Health services should regularly audit surgical prophylaxis practices to ensure that:

surgical patients should receive timely prophylaxis when indicated;
correct antibiotics, route of administration and dosage are used;

• the duration of prophylaxis is appropriate.

Indicator set: Clinical care standard indicators: antimicrobial stewardship

Australian Commission on Safety and Quality in Health Care, Retired

25/01/2022

Health, Standard 12/09/2016

Outcome area: Surgical prophylaxis

Health, Standard 12/09/2016

## Collection and usage attributes

Computation description: Prophylaxis is the use of antibiotics to prevent surgical site infection and, in some

circumstances, bacteraemia.

The numerator and denominator include patients undergoing surgery for which there are documented guidelines for the administration of prophylactic antibiotics, that is, the *Therapeutic guidelines: antibiotic* or locally-endorsed antibiotic

prescribing guidelines.

The numerator and denominator exclude patients undergoing surgery for which there are no documented guidelines for the administration of prophylactic antibiotics, that is, the *Therapeutic guidelines: antibiotic* or locally-endorsed antibiotic prescribing guidelines, or where prophylaxis is not indicated due to the

patient's current antibiotic therapy.

For the numerator, 'in accordance' with guidelines means that the choice of antimicrobial is in accordance with guidelines. Timing of dose is dealt with in Antimicrobial stewardship: 9b-Timely administration of prophylactic antibiotics prior to surgery, and duration of antibiotic therapy is dealt with in Antimicrobial stewardship: 9c-Cessation of prophylactic antibiotics following surgery.

Presented as a percentage.

**Computation:** (Numerator ÷ denominator) x 100

**Numerator:** Number of admitted patients undergoing surgery who received prophylactic

antibiotics in accordance with Therapeutic guidelines: antibiotic or locally-

endorsed antibiotic prescribing guidelines.

**Denominator:** Number of admitted patients undergoing surgery for which there are documented

guidelines for the administration of prophylactic antibiotics.

Comments:

The *Therapeutic guidelines: antibiotic* provides guidance on administration of prophylactic antibiotics for the following types of surgery:

- · abdominal surgery, including:
  - colorectal surgery, appendicectomy, and upper gastrointestinal tract or biliary surgery (including laparoscopic approaches)
  - o endoscopic procedures
  - o hernia repair
- burns, extensive skin loss
- cardiac surgery
- · head, neck and thoracic surgery
- lower limb amputation
- neurosurgery
- · obstetric and gynaecological surgery, including:
  - hysterectomy and termination of pregnancy
  - o caesarean section
- · orthopaedic surgery
- urological surgery, including:
  - prostatectomy
  - transrectal prostatic biopsy
- vascular surgery.

Services may select specific procedures to focus on as part of the audit process, based on their own mix of patients and priorities.

This indicator is based on surgical antibiotic prophylaxis indicators developed by the then Victorian Hospital Acquired Infection Surveillance System Coordinating Centre (VICNISS) (Department of Human Services 2007) (now the Healthcare Associated Infection Surveillance Coordinating Centre), and the Western Australian Safety and Quality Investment for Reform (SQuIRe) Program (Department of Health WA 2009).

## Representational attributes

Representation class: Percentage

Data type: Real

Unit of measure: Service event

Format: N[NN]

#### Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

**Reference documents:** Antibiotic Expert Group 2014. Therapeutic guidelines: antibiotic. Version 15.

Melbourne: Therapeutic Guidelines Limited.

Department of Health WA 2009. SQulRe 2 CPI guide: Surgical site infection

prevention. Perth: DHWA.

Department of Human Services 2007. Type 2 surveillance manual version 10. Melbourne: Victorian Hospital Acquired Infection Surveillance System Coordinating

Centre, DHS.