# Minimising use of antipsychotic medicines

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY4.0 (CC BY4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

## Minimising use of antipsychotic medicines

## Identifying and definitional attributes

| Metadata item type:  | Outcome Area   |
|----------------------|--|
| METEOR identifier:   | 624408   |
| Registration status: | Health, Standard 12/09/2016  |
| Description:         | Treatment with an antipsychotic medicine is only considered if a patient with delirium is distressed and the cause of their distress cannot be addressed and non-drug strategies have failed to ease their symptoms. |

#### **Relational attributes**

| Indicator sets linked to this outcome area: | <u>Clinical care standard indicators: delirium</u><br><u>Health</u> , Standard 12/09/2016   |
|---|---|
| Indicators linked to this outcome area:     | Delirium clinical care standard indicators: 6a-Evidence of local arrangements to<br>ensure that patients with delirium are not routinely prescribed antipsychotic<br>medicines<br>Health, Standard 12/09/2016 |
|   | Delirium clinical care standard indicators: 6b-Proportion of patients with delirium<br>prescribed antipsychotic medicines in hospital<br>Health, Standard 12/09/2016  |

### Source and reference attributes

| Submitting organisation: | Australian Commission on Safety and Quality in Health Care   |
|--------------------------|--|
| Reference documents:     | ACSQHC (Australian Commission on Safety and Quality in Health Care) 2015. Delirium clinical care standard. Sydney: ACSQHC. |