

Pain management

Identifying and definitional attributes

Metadata item type: Outcome Area

METEOR identifier: 628050

Registration status:

- [Health](#), Standard 12/09/2016

Description: A patient with a hip fracture is assessed for pain at the time of presentation and regularly throughout their hospital stay, and receives pain management including the use of multimodal analgesia, if clinically appropriate.

Relational attributes

Indicator sets linked to this outcome area: [Clinical care standard indicators: hip fractureHealth](#), Standard 12/09/2016
[Australian Commission on Safety and Quality in Health Care](#), Superseded 18/06/2018

[Clinical care standard indicators: hip fracture 2018Australian Commission on Safety and Quality in Health Care](#), Standard 15/05/2018

Indicators linked to this outcome area: [Hip fracture care clinical care standard indicators: 2a-Evidence of local arrangements for timely and effective pain management for hip fractureHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 2a-Evidence of local arrangements for timely and effective pain management for hip fractureHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require itHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require itHealth](#), Standard 12/09/2016

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care