

Mobilisation and weight-bearing

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Mobilisation and weight-bearing

Identifying and definitional attributes

Metadata item type:	Outcome Area
METEOR identifier:	628056
Registration status:	Health , Standard 12/09/2016
Description:	A patient with a hip fracture is offered mobilisation without restrictions on weight-bearing the day after surgery and at least once a day thereafter, depending on the patient's clinical condition and agreed goals of care.

Relational attributes

Indicator sets linked to this outcome area:	Clinical care standard indicators: hip fracture Australian Commission on Safety and Quality in Health Care , Superseded 18/06/2018 Health , Standard 12/09/2016 Clinical care standard indicators: hip fracture 2018 Australian Commission on Safety and Quality in Health Care , Standard 15/05/2018
Indicators linked to this outcome area:	Hip fracture care clinical care standard indicators: 5a-Proportion of patients with a hip fracture who are mobilised on day one post hip fracture surgery Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 5a-Proportion of patients with a hip fracture who are mobilised on day one post hip fracture surgery Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 5c-Proportion of patients with a hip fracture experiencing a new Stage II or higher pressure injury during their hospital stay Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 5c-Proportion of patients with a hip fracture experiencing a new Stage II or higher pressure injury during their hospital stay Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 5d-Proportion of patients with a hip fracture returning to pre-fracture mobility Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 5d-Proportion of patients with a hip fracture returning to pre-fracture mobility Health , Standard 12/09/2016

Source and reference attributes

Submitting organisation:	Australian Commission on Safety and Quality in Health Care
---------------------------------	--