

Mobilisation and weight-bearing

Identifying and definitional attributes

Metadata item type: Outcome Area

METEOR identifier: 628056

Registration status:

- [Health](#), Standard 12/09/2016

Description: A patient with a hip fracture is offered mobilisation without restrictions on weight-bearing the day after surgery and at least once a day thereafter, depending on the patient's clinical condition and agreed goals of care.

Relational attributes

Indicator sets linked to this outcome area: [Clinical care standard indicators: hip fractureHealth](#), Standard 12/09/2016
[Australian Commission on Safety and Quality in Health Care](#), Superseded 18/06/2018

[Clinical care standard indicators: hip fracture 2018Australian Commission on Safety and Quality in Health Care](#), Standard 15/05/2018

Indicators linked to this outcome area: [Hip fracture care clinical care standard indicators: 5a-Proportion of patients with a hip fracture who are mobilised on day one post hip fracture surgeryHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5a-Proportion of patients with a hip fracture who are mobilised on day one post hip fracture surgeryHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgeryHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgeryHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5c-Proportion of patients with a hip fracture experiencing a new Stage II or higher pressure injury during their hospital stayHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5c-Proportion of patients with a hip fracture experiencing a new Stage II or higher pressure injury during their hospital stayHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5d-Proportion of patients with a hip fracture returning to pre-fracture mobilityHealth](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5d-Proportion of patients with a hip fracture returning to pre-fracture mobilityHealth](#), Standard 12/09/2016

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care