

Minimising risk of another fracture

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Minimising risk of another fracture

Identifying and definitional attributes

Metadata item type:	Outcome Area
METEOR identifier:	628058
Registration status:	Health , Standard 12/09/2016
Description:	Before a patient with a hip fracture leaves hospital, they are offered a falls and bone health assessment, and a management plan based on this assessment to reduce the risk of another fracture.

Relational attributes

Indicator sets linked to this outcome area:	Clinical care standard indicators: hip fracture Australian Commission on Safety and Quality in Health Care , Superseded 18/06/2018 Health , Standard 12/09/2016 Clinical care standard indicators: hip fracture 2018 Australian Commission on Safety and Quality in Health Care , Standard 15/05/2018
Indicators linked to this outcome area:	Hip fracture care clinical care standard indicators: 6a-Proportion of patients with a hip fracture receiving bone protection medicine prior to separation from the hospital at which they underwent surgery Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 6a-Proportion of patients with a hip fracture receiving bone protection medicine prior to separation from the hospital at which they underwent surgery Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 6b-Proportion of patients with a hip fracture readmitted to hospital with another femoral fracture within 12 months of admission from initial hip fracture Health , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 6b-Proportion of patients with a hip fracture readmitted to hospital with another femoral fracture within 12 months of admission from initial hip fracture Health , Standard 12/09/2016

Source and reference attributes

Submitting organisation:	Australian Commission on Safety and Quality in Health Care
---------------------------------	--