

# **Hip fracture care clinical care standard indicators: 2b- Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it**

**Exported from METEOR (AIHW's Metadata Online Registry)**

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it

## Identifying and definitional attributes

<b>Metadata item type:</b>	Indicator
<b>Indicator type:</b>	Indicator
<b>Short name:</b>	Indicator 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it
<b>METEOR identifier:</b>	628070
<b>Registration status:</b>	<a href="#">Health</a> , Standard 12/09/2016
<b>Description:</b>	Proportion of patients with a <a href="#">hip fracture</a> who have a documented assessment of pain within 30 minutes of presentation to the emergency department and either receive analgesia within this time or do not require it according to the assessment.
<b>Rationale:</b>	<p>Acute pain and injury of various types are inevitably interrelated and if severe and prolonged, the injury response invoked in the patient becomes counterproductive and can have adverse effects on outcome (Macintyre et al. 2010).</p> <p>Specific early analgesic interventions may reduce the incidence of chronic pain after surgery (Macintyre et al. 2010).</p>
<b>Indicator set:</b>	<a href="#">Clinical care standard indicators: hip fracture</a> <a href="#">Australian Commission on Safety and Quality in Health Care</a> , Superseded 18/06/2018 <a href="#">Health</a> , Standard 12/09/2016
<b>Outcome area:</b>	<a href="#">Pain management</a> <a href="#">Health</a> , Standard 12/09/2016

## Collection and usage attributes

<b>Computation description:</b>	Both the numerator and the denominator include patients admitted to hospital for a hip fracture, initially presenting to the emergency department with the fracture.  Presented as a percentage.
<b>Computation:</b>	$(\text{Numerator} \div \text{denominator}) \times 100$
<b>Numerator:</b>	Number of patients admitted to hospital with a hip fracture, initially presenting to the emergency department, who have a documented assessment of pain within 30 minutes of presentation to the emergency department and either receive analgesia within this time frame or have documented in their medical record that they do not require analgesia at this point according to the assessment.
<b>Denominator:</b>	Number of patients admitted to hospital with a hip fracture, initially presenting to the emergency department.

## Representational attributes

<b>Representation class:</b>	Percentage
<b>Data type:</b>	Real

**Unit of measure:** Service event

**Format:** N[NN]

## Source and reference attributes

**Submitting organisation:** Australian Commission on Safety and Quality in Health Care

**Reference documents:** Macintyre PE, Scott DA, Schug SA, Visser EJ, Walker SM; Working Group of the Australian and New Zealand College of Anaesthetists and Faculty of Pain Medicine 2010. Acute Pain Management: Scientific Evidence 3rd edn, Melbourne: ANZCA & FPM.